

NEUMANN PROBLEMS FOR p -HARMONIC FUNCTIONS, AND INDUCED NONLOCAL OPERATORS IN METRIC MEASURE SPACES

LUCA CAPOGNA, JOSH KLINE, RIIKKA KORTE,
NAGESWARI SHANMUGALINGAM, AND MARIE SNIPES

ABSTRACT. Following ideas of Caffarelli and Silvestre in [20], and using recent progress in hyperbolic fillings, we define fractional p -Laplacians $(-\Delta_p)^\theta$ with $0 < \theta < 1$ on any compact, doubling metric measure space (Z, d, ν) , and prove existence, regularity and stability for the non-homogenous non-local equation $(-\Delta_p)^\theta u = f$. These results, in turn, rest on the new existence, global Hölder regularity and stability theorems that we prove for the Neumann problem for p -Laplacians Δ_p , $1 < p < \infty$, in bounded domains of measure metric spaces endowed with a doubling measure that supports a Poincaré inequality. Existence of solutions for the Neumann boundary value problem in the metric setting was first demonstrated in [55] for the case that the Neumann data f is bounded, and here we relax the boundedness requirement of f with a more general integrability requirement. Our work also includes as special cases much of the previous results by other authors in the context of compact subsets in the Euclidean, Riemannian and Carnot group settings. Unlike other recent contributions in the metric measure spaces context, our work does not rely on the assumption that (Z, d, ν) supports a Poincaré inequality.

CONTENTS

1. Introduction	2
2. Background	10
2.1. John and uniform domains	10
2.2. Measures	11
2.3. Newton-Sobolev functions and Poincaré inequalities	11
2.4. Besov spaces	12
2.5. Cheeger differential structures	15
2.6. Neumann boundary value problem	16
2.7. Hyperbolic fillings	16

The research of N.S. and J.K. are partially funded by the NSF grants #DMS-1800161 and #DMS-2054960. The research of M.S. is partially funded by a supplemental grant from NSF #DMS-1800161. The research of L.C. is partly funded by NSF #DMS-195599 and NSF #2348806. R.K. was supported by Academy of Finland, projects 308063 and 360184. The authors thank Mathav Murugan and Zhen-Qing Chen for illuminating discussion on jump processes and for pointing out the references [27, 28, 29] to us when N.S. and they were visiting at MSRI in Spring 2022, and Yannick Sire for pointing out the reference [22]. We also thank the anonymous referee for the thoughtful comments that helped us improve the exposition of the paper. **MRC Classification (2020):** 30L15, 31E05, 35R11, 35B65, 46E35.

3. Equivalent Formulations of the Neumann problem for the p -Laplacian	17
4. Boundary Regularity for $1 < p \leq Q$	19
5. Stability of solutions under perturbation of the Neumann data	25
6. Constructing an induced non-local fractional Laplacian for compact doubling metric measure spaces	29
6.1. The general case $1 < p < \infty$	29
6.2. The case $p = 2$	33
7. Reconciling construction of fractional Laplacian with [33]	34
7.1. Doubling property of μ_ω .	37
7.2. Modification of the norm on Cheeger differential structure	40
7.3. 2-Poincaré inequality with respect to d_ρ and μ_ω	40
7.4. John domain property	43
7.5. Conclusion: the reconciliation	44
Appendix A. Removing boundedness condition on f in [55] and establishing existence and boundedness of solutions	45
References	52

1. INTRODUCTION

The goal of this paper is to construct and study a notion of the fractional p -Laplacian in the setting of compact doubling metric measure spaces, by extending Caffarelli and Silvestre's approach in [20]. Our strategy consists of two main steps: First, we continue the study begun in [55], of the Neumann boundary value problem for the p -Laplacian, $1 < p < \infty$,

$$\begin{aligned}\Delta_p u &:= \operatorname{div}(|\nabla u|^{p-2} \nabla u) = 0 \text{ in } \Omega, \\ |\nabla u|^{p-2} \partial_\eta u &= f \text{ on } \partial\Omega,\end{aligned}$$

expressed in the weak form

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla \phi \, d\mu = \int_{\partial\Omega} \phi f \, d\nu \quad \text{for every Sobolev function } \phi,$$

where Ω is an open domain in a metric measure space (X, d, μ) equipped with a doubling measure supporting a Poincaré inequality. In this general setting ∇u indicates either the gradient in a Cheeger differential structure (see [25]) or in an infinitesimally Hilbertian structure in the sense of Gigli [2, 3, 41, 42], and hence provides a unified framework for a wide class of operators. We prove global Hölder regularity and stability results with respect to the Neumann data in suitable Lebesgue classes.

Second, using our new results for the Neumann problem, we provide a construction of a family of fractional p -Laplacian operators on the boundary of the domain. In fact, thanks to Theorem 1.5, we can construct an analog of the fractional Laplacian for *any* compact doubling metric measure space (Z, d_Z, ν) even if it does not support a 2-Poincaré inequality. We do this by constructing a locally compact but non-compact doubling metric measure space (X, d, μ) that supports a 1-Poincaré inequality such that Z

is biLipschitz equivalent to $\partial X = \overline{X} \setminus X$, and considering solutions to the corresponding Neumann boundary value problem in X . Such a construction of a metric measure space X as a uniformization of a hyperbolic filling of Z can be found in [14]. We will discuss this in detail in Section 2.7 below. As a consequence, we prove existence, regularity, Harnack inequality and stability for solutions of the associated non-homogenous equation

$$(-\Delta_p)^\theta v = f \text{ in } Z,$$

with $\theta \in (0, 1)$. We also prove that for $p = 2$, the fractional operators defined here are the same as those appearing in the literature, for example Euclidean setting [20, 61], Riemannian manifolds [8], Carnot groups [34], and metric measure spaces [33]. A recently completed companion paper, [23], provides an extension of our results to the unbounded setting, thus generalizing the results found in [20, 33, 34, 61].

We remark explicitly that even in the special case $p = 2$, part of the novelty of our approach is that it allows one to define and study fractional Laplacians in metric measure spaces that support neither a Poincaré inequality nor a corresponding regular Dirichlet form, such as the Von Koch snowflake K , or the Rickman rug $[0, 1] \times K$.

Next, we proceed by outlining the relevant assumptions of the paper, before stating our main results.

Structure hypotheses: Throughout the paper we let $1 < p < \infty$, and Ω be an open, connected domain in a complete metric measure space (X, d, μ) such that:

- (H0) Ω is a John domain as defined in Subsection 2.1.
- (H1) $(\Omega, d, \mu|_{\overline{\Omega}})$ is doubling and supports a p -Poincaré inequality as in Subsection 2.2 and Subsection 2.3.
- (H2) The boundary $\partial\Omega$ is equipped with a Radon measure ν for which there are constants $C \geq 1$ and $0 < \Theta < p$ such that for all $x \in Z$ and $0 < r < 2 \operatorname{diam}(\partial\Omega)$,

$$(1.1) \quad \frac{1}{C} \frac{\mu(B(x, r) \cap \Omega)}{r^\Theta} \leq \nu(B(x, r)) \leq C \frac{\mu(B(x, r) \cap \Omega)}{r^\Theta};$$

that is, ν is a Θ -codimensional Hausdorff measure with respect to $\mu|_{\Omega}$.

Remark 1.2. The constants associated with p , the John domain condition, and the above two conditions will be referred to as the structural constants.

Recall that John domains are necessarily bounded domains. Condition (H0) can be waived for Theorem 1.5 and Theorem 1.6. The boundedness of Ω is used by us only to know that solutions to the Neumann boundary value problem exist (see [55]), and Condition (H0) is used only to know the existence of traces of Sobolev functions on Ω and ensure that the trace lies in the suitable Besov class of functions on the boundary of Ω . For unbounded domains, John condition can be replaced with the assumption that Ω is a uniform domain in order to obtain local trace estimates as in [54], and this is sufficient for Theorem 1.5 and Theorem 1.6 in the event that Ω is not bounded. The recent paper [23] implements this extension in the unbounded setting. Note

that [20, 34] establish neither existence nor stability results for the fractional Laplacian, but focus only on local regularity such as Hölder continuity and Harnack inequality, under the assumption that the solution to the fractional Laplacian problem exists.

Remark 1.3. Our choice of $0 < \Theta < p$ in (H2) corresponds to the choice of a in [20]. In [20], for each $0 < \theta < 1$ the choice $-1 < a = 1 - 2\theta < 1$ is made in considering the weight y^a imposed on the domain $\Omega = Z \times (0, \infty)$, and in this paper, we need $\Theta = 2(1 - \theta)$ (with $p = 2$) correspondingly.

Remark 1.4. The structure hypotheses are more general than those assumed in the literature so far. In fact, domains with fractal boundaries such as in the vonKoch snowflake domain satisfy these conditions for some $\Theta \neq 1$. Moreover, such a flexibility allows us to consider all powers $\theta \in (0, 1)$ rather than just $\theta = 1/2$. These structural assumptions are satisfied by the contexts studied in [20, 34, 61], where $\Omega = X \times (0, \infty)$ with X either a Euclidean space or a Carnot group, and Ω equipped with a weighted product measure. Moreover, *every* compact doubling metric measure space X is the boundary of a John domain (and in fact, the boundary of a uniform domain) satisfying our structural conditions, see [14].

The first main theorem of this paper, stated next, is the main tool we use to construct the fractional operator on the metric space $(\partial\Omega, d, \nu)$ by considering analogs of the Dirichlet-to-Neumann transformation. Condition (c) of this theorem tells us that the trace of the solution u on $\partial\Omega$ should belong to the domain of the fractional operator.

Theorem 1.5. *Suppose that conditions (H0), (H1) and (H2) hold for the metric measure space (X, d, μ) and Ω . Fix $1 < p < \infty$, and suppose that $f : \partial\Omega \rightarrow \mathbb{R}$ be in the class $L^{p'}(\partial\Omega)$ where p' is the Hölder conjugate of p with $\int_{\partial\Omega} f d\nu = 0$. Let $u \in N^{1,p}(\Omega) = N^{1,p}(\overline{\Omega})$. Then the following are equivalent.*

- (a) *u is a solution to the Neumann boundary value problem with data f in the domain $\Omega \subset X$; that is, for all $\phi \in N^{1,p}(\overline{\Omega})$,*

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla \phi d\mu = \int_{\partial\Omega} \phi f d\nu$$

- (b) *u minimizes the energy functional*

$$I(v) := \int_{\overline{\Omega}} |\nabla v|^p d\mu - p \int_{\partial\Omega} v f d\nu$$

among all functions $v \in N^{1,p}(\overline{\Omega})$. Here we extend ν to a measure on the closure $\overline{\Omega}$ by zero outside of $\partial\Omega$.

- (c) *u is p -harmonic in Ω and*

$$|\nabla u|^{p-2} \nabla u \cdot \nabla \eta_\epsilon d\mu \rightharpoonup -f d\nu.$$

Here the convergence is that of weak convergence of signed Radon measures on $\overline{\Omega}$, and the function $\eta_\epsilon : \overline{\Omega} \rightarrow \mathbb{R}$ is given by $\eta_\epsilon(x) = \min\{1, \text{dist}(x, \overline{\Omega} \setminus \Omega)/\epsilon\}$, and again we extend ν to a measure on the closure $\overline{\Omega}$ by zero outside of $\partial\Omega$.

The proof of Theorem 1.5 is in Section 3. Note that from our structural assumption (H2), we have that $\int_{B(\zeta, r)} |\nabla \eta_\varepsilon| d\mu$ is at most a constant multiple of $\varepsilon^{-1+\Theta} \nu(B(\zeta, 2r))$ whenever $\zeta \in \partial\Omega$ and $r > 0$.

Next we turn our attention to global properties of solutions of the Neumann boundary value problem, namely we obtain Hölder regularity up to the boundary, and stability with respect to the data. Here, the index Q is the lower mass bound exponent associated with the measure μ as in (2.1). Notice that we can increase the value of Q in (2.1), and so, if we are willing to pay the price of changing the estimates in the following theorem, the restriction $p \leq Q$ should not be considered to be a restrictive one. Note that when p increases, its Hölder dual p' decreases, and so we have a wider class of admissible Neumann data $f \in L^{p'}(\partial\Omega)$.

Theorem 1.6. *Assume that (H0), (H1) and (H2) hold, $1 < p \leq Q$, and let u be a solution of the Neumann problem as in Theorem 1.5, for the boundary data $f \in L^q(\partial\Omega, \nu)$, with $\int_{\partial\Omega} f d\nu = 0$. Denote by $B_{R_0} \subset \bar{\Omega}$ the intersection of $\bar{\Omega}$ with a ball of radius $R_0 > 0$ centered at a point in $\partial\Omega$. If the boundary data satisfies the additional integrability assumption $f \in L^q(B_{2R_0} \cap \partial\Omega, d\nu)$ for some q with*

$$q_0 := \frac{Q-\Theta}{p-\Theta} < q \leq \infty,$$

then any solution of the Neumann problem u is ε -Hölder continuous in B_{R_0} with

$$\varepsilon = \min \left\{ \beta_0, \left(1 - \frac{\Theta}{p} \right) \left(1 - \frac{q_0}{q} \right) \right\},$$

where $\beta_0 > 0$ is the Hölder exponent for the interior regularity estimates in Proposition 4.1. More specifically, one has the estimate

$$(1.7) \quad \sup_{x, y \in B_{R_0}, x \neq y} \frac{|u(x) - u(y)|}{d(x, y)^\varepsilon} \leq C,$$

where $C > 0$ is a constant depending on the structure constants, on the choice of q , on $\|f\|_{L^q(B_{2R_0} \cap \partial\Omega)}$ and on R_0 . Furthermore, there is a constant $C > 1$ such that if $u \geq 0$ on Ω and $W \subset \partial\Omega$ is a non-empty relatively open subset of $\partial\Omega$ with $f = 0$ on W , then whenever $x \in \Omega \cup W$ and $r > 0$ such that $B(x, 2r) \cap \bar{\Omega} \subset W \cup \Omega$, we have the Harnack inequality

$$\sup_{B(x, r) \cap \bar{\Omega}} u \leq C \inf_{B(x, r) \cap \bar{\Omega}} u.$$

Remark 1.8. In subregions where f is non-negative, the results of [53] yield a better Hölder exponent for the solutions than the one above, i.e.

$$\epsilon = \left(1 - \frac{q_0}{q} \right) \frac{p - \Theta}{p - 1},$$

(see [23, Section 9.4]). The argument in [53] relies on Caccioppoli-type estimates which are not available in regions where f changes sign. In the Euclidean setting, the results in [57] lead to the same (optimal) Hölder exponent without any restrictions on the sign of f . We believe the techniques of [57] can be extended to our more general setting, yielding the same optimal exponent, and plan to return to this issue in a future paper.

This theorem will be proved in Section 4, but an explanation regarding the proof is warranted here. Even in the Euclidean setting, the perspective of metric spaces gives a new viewpoint of the Neumann problem; we can see the domain Ω as an open subset of an ambient metric measure space while also viewing $\bar{\Omega}$ as a metric measure space in its own right. This allows us to see the solutions for Neumann boundary value problem also as solutions to the inhomogeneous problem $-\Delta_p u = \nu_f$ on the metric measure space $\bar{\Omega}$, where the measure ν_f is a singular measure, supported on the boundary $\partial\Omega = \bar{\Omega} \setminus \Omega$, and is associated with the Neumann data of the original Neumann boundary value problem. This point of view allows us to adapt a Morrey-Campanato type argument on the whole “open set” that is $\bar{\Omega}$. To do so, we took inspiration from [58], but as we rely on the results from [55] and [49], our proof is more direct. The idea is to prove a version of the Morrey inequality for the metric measure space setting. Note that with our assumption on $\partial\Omega$, we have that $\mu(\partial\Omega) = 0$. For $p = 2$, the argument in the proof of Hölder continuity on regions of $\partial\Omega$ where $f = 0$, in the Euclidean, Riemann manifold setting, and Carnot setting as established by [8, 20, 24, 34, 61] is based on a Harnack inequality, from which the Hölder continuity follows. Here we give a more direct proof of the theorem, and in doing so, we obtain Hölder continuity even in regions of the boundary where the Neumann data f does not vanish. Hölder regularity results for the non-homogeneous equation were obtained for bounded Euclidean domains in [22], and it is interesting to note that the integrability conditions $f \in L^q_{loc}(\partial\Omega, d\nu)$ placed on the data f in [22] is exactly the condition

$$q > \frac{Q - \Theta}{p - \Theta},$$

on q that we find in the coarser setting of nonsmooth metric spaces. We also point out that, unlike the above-mentioned papers, our discussion also includes the nonlinear setting $p \neq 2$ of the (fractional) p -Laplacian, and that the Harnack inequality for the homogenous fractional PDE follows by virtue of the global Hölder regularity of the Neumann problem and by the results in [49].

In terms of continuity with respect to boundary data, we prove the following,

Theorem 1.9. *Let $1 < p$ and $p' = \frac{p}{p-1}$, and suppose that (H0), (H1) and (H2) hold. There exists a constant $C > 0$ depending on the structural constants in (H0), (H1), and (H2) such that the following holds. For boundary data $f, g \in L^{p'}(\partial\Omega, \nu)$ with*

$$\int_{\partial\Omega} f \, d\nu = 0 = \int_{\partial\Omega} g \, d\nu,$$

denote by $u, v \in N^{1,p}(\bar{\Omega})$ respectively the solutions to the corresponding p -Neumann problems such that

$$\int_{\Omega} u \, d\mu = 0 = \int_{\Omega} v \, d\mu.$$

Then, when $p \geq 2$ we have

$$\|\nabla u - \nabla v\|_{L^p(\Omega)} \leq C \left(\|f\|_{L^{p'}(\partial\Omega)} + \|g\|_{L^{p'}(\partial\Omega)} \right)^{\frac{1}{p(p-1)}} \|f - g\|_{L^{p'}(\partial\Omega)}^{\frac{1}{p}},$$

and when $1 < p < 2$ we have

$$\|\nabla u - \nabla v\|_{L^p(\Omega)} \leq \left(\|f\|_{L^{p'}(\partial\Omega)} + \|g\|_{L^{p'}(\partial\Omega)} \right)^{\frac{3-p}{2(p-1)}} \|f - g\|_{L^{p'}(\partial\Omega)}^{\frac{1}{2}}.$$

As a consequence of this theorem together with Lemma A.3 and the Poincaré inequality, we see that if the boundary data converges in $L^{p'}$ to a function $f \in L^{p'}$, then the solutions converge to a solution of the Neumann boundary value problem with boundary data f . We will prove this theorem in Section 5 below. In addition, in Theorem 5.4 we also prove a weaker form of stability that applies to the Neumann problem with respect to the upper gradient formulation. Theorem 5.4 is independent of the above theorem as it considers a variant Neumann boundary value problem that arises from an energy minimization principle that may not correspond to an Euler-Lagrange equation unless the metric measure space is infinitesimally Hilbertian in the sense of Gigli [2, 3, 41, 42]. Such an Euler-Lagrange equation is essential for our proof of Theorem 1.9. In the setting of hyperbolic filling as in [14] and in Section 6, there is a Cheeger differential structure that is available to us, and so from the point of view of studying nonlocal minimization problems, we do not lose much by considering the Cheeger differential formulation.

We are now ready to introduce and discuss the fractional (non-local) operators we are interested in.

Definition 1.10. Let (Z, d, ν) be a metric measure space. For any $1 < p < \infty$ and $0 < \theta < 1$, consider a form $\mathcal{E} : L^p(Z) \times L^p(Z) \rightarrow [-\infty, \infty]$, that is linear in the second component and with $\mathcal{E}(\alpha u, \beta v) = |\alpha|^{p-2} \alpha \beta \mathcal{E}(u, v)$, such that $\mathcal{E}(u, u) \approx \|u\|_{B_{p,p}^\theta(Z)}^\theta$ whenever u is in the Besov class $B_{p,p}^\theta(Z)$. We say that a function $u \in B_{p,p}^\theta(Z)$ is in the domain of the fractional p -Laplacian operator $(-\Delta_p)^\theta$ if there is a function $f \in L^{p'}(Z)$ such that the integral identity

$$\mathcal{E}(u, \varphi) = \int_Z \varphi f d\nu$$

holds for every $\varphi \in B_{p,p}^\theta(Z)$. We then denote

$$(-\Delta_p)^\theta u = f \in L^{p'}(Z).$$

The above definition is concomitant with the notion of Laplacian in the theory of Dirichlet forms, see for example the comprehensive book [36]. The fractional operators we consider in this paper are associated to the form (1.11) below. Its definition is based on a process reminiscent of the hyperbolic filling technique: We will show in Section 6 that, given $1 < p < \infty$ and $0 < \theta < 1$, every compact doubling metric measure space Z arises as the boundary of a uniform domain Ω that is equipped with a measure μ so that the metric measure space $X = \overline{\Omega} = \Omega \cup Z$, together with $Z = \partial\Omega$, satisfies conditions (H0), (H1) and (H2), with $\theta = 1 - \Theta/p$. We fix a Cheeger differential structure ∇ on Ω . For each $u \in B_{p,p}^\theta(Z)$ we consider \hat{u} to be

the unique function in $N^{1,p}(\Omega)$ such that \widehat{u} is Cheeger p -harmonic in Ω and has trace $Tr(\widehat{u}) = u$ ν -almost everywhere on Z . We will show that $\int_{\Omega} |\nabla \widehat{u}|^p d\mu \approx \|u\|_{B_{p,p}^{\theta}(Z)}^p$ (see Section 6 below). We then set

$$(1.11) \quad \mathcal{E}_T(u, v) := \int_{\Omega} |\nabla \widehat{u}|^{p-2} \nabla \widehat{u} \cdot \nabla \widehat{v} d\mu.$$

This construction gives us a way of analyzing a wide array of fractional operators $(-\Delta_p)^{\theta}$ on the Besov class $B_{p,p}^{\theta}(Z)$, for we have a broad choice of Ω , and for each choice of Ω we then have the flexibility of choosing a desired Cheeger differential structure as outlined in [25] (see [46] for further exposition on Cheeger differential structure). We may, instead of a Cheeger differential structure, consider the Γ -limit of discrete differential structures as discussed in [32]. Should Ω be infinitesimally Hilbertian in the sense of [41, 42], we may use the differential structure associated with the infinitesimal Hilbertianity. Additional choices of structures are available in the Euclidean setting, and a related family of non-local fractional operators were studied by Caffarelli and Soria-Carro in [21] using lower dimensional slices.

Another approach to the notion of Laplacian is from the theory of Dirichlet forms, see for example [36]. To emphasize this connection we introduce a new form $\mathcal{E}_p(u, v)$, given by

$$\mathcal{E}_p(u, v) = \int_Z \int_Z \frac{|u(y) - u(x)|^{p-2} (u(y) - u(x))(v(y) - v(x))}{\nu(B(x, d(x, y))) d(x, y)^{\theta p}} d\nu(y) d\nu(x),$$

with $u, v \in L^p(Z)$. Note that \mathcal{E}_2 is a Dirichlet form in the sense of [36], and is associated with a jump process, see also [9, 27, 28, 29]. In the following theorem, Q_Z is the lower mass bound exponent corresponding to Q from (2.1) for the doubling measure ν on Z , and $p' = p/(p-1)$ is the Hölder conjugate of p .

Theorem 1.12. *Let (Z, d, ν) be a compact doubling metric measure space, $1 < p < \infty$, and $0 < \theta < 1$. Then the form \mathcal{E}_T on $B_{p,p}^{\theta}(Z)$ given by (1.11) satisfies $\mathcal{E}_T(u, u) \approx \mathcal{E}_p(u, u)$ for each $u \in B_{p,p}^{\theta}(Z)$ with the comparison constant depending solely on the doubling constant of ν and the indices p, θ . Denote by $(-\Delta_p)^{\theta}$ the fractional p -Laplacian associated to the form $\mathcal{E}_T(u, u)$. Moreover,*

- (i) *For each $f \in L^{p'}(Z)$ with $\int_Z f d\nu = 0$ there exists a function $u_f \in B_{p,p}^{\theta}(Z)$ such that $(-\Delta_p)^{\theta} u_f = f$ on Z . If \widetilde{u}_f is any other such function, then $u_f - \widetilde{u}_f$ is constant ν -a.e. in Z . If in addition $f \in L^q(Z)$ for some $q > \max\{1, Q_Z/\theta\}$, then u_f is Hölder continuous on Z .*
- (ii) *There exists a constant $C > 0$, depending only on the structure constants, such that if $f_1, f_2 \in L^{p'}(Z)$ with $\int_Z f_1 d\nu = 0 = \int_Z f_2 d\nu$ and u_{f_1}, u_{f_2} are the functions in $B_{p,p}^{\theta}(Z)$ corresponding to f_1, f_2 as above, then*

$$\|u_{f_1} - u_{f_2}\|_{L^p(Z)} \leq C \max\{\|f_1\|_{L^{p'}(Z)}, \|f_2\|_{L^{p'}(Z)}\}^{\kappa} \|f_1 - f_2\|_{L^{p'}(Z)}^{\tau},$$

with $\kappa = 1/(p(p-1))$, $\tau = 1/p$ when $p \geq 2$ and $\kappa = (3-p)/(2(p-1))$, $\tau = 1/2$ when $1 < p < 2$.

- (iii) Let $W \subset Z$ be an open (nonempty) subset such that $f = 0$ on W . There exists a constant $C > 0$ depending only on the structure constants, such that if $u \geq 0$ is a solution of $(-\Delta_p)^\theta u = f$ in Z , then u satisfies the Harnack inequality $\sup_B u \leq C \inf_B u$ for all balls $B = B_R$ such that $B_{4R} \subset W$.

This theorem will be proved in Section 6.

Remark 1.13. We can weaken the hypotheses of the above theorem by replacing the compactness requirement of Z with the condition that Z is biLipschitz equivalent to the boundary of a uniform domain that satisfies the conditions (H0), (H1), and (H2). Thus the above theorem also includes the cases when Z is the entire Euclidean space as in [20] or even a Carnot groups as in [34].

In [33] an alternate construction, based on spectral theory, for a fractional Laplacian $(-\Delta_2)^\theta$ corresponding to a Cheeger differential structure on a complete doubling metric measure space (Z, d_Z, ν) supporting a 2-Poincaré inequality was constructed and studied. The methods used there depended strongly on the availability of the Poincaré inequality on the metric space itself, and moreover, it is not adaptable to fractional powers of the nonlinear p -Laplacian operator $(-\Delta_p)^\theta$. In [33], the metric space (Z, d_Z, ν) is naturally seen as the boundary of the unbounded domain $Z \times (0, \infty)$, where $X = Z \times [0, \infty)$ is equipped with the ℓ_2 -product metric and the product measure. In contrast, in our present study, we consider fractional Laplacians on the boundary of a *bounded* domain. Therefore, to show that the two approaches lead to the same notion of nonlocal Laplacian, we show that we can conformally transform X into a bounded doubling metric measure space Ω so that it supports a 2-Poincaré inequality and that functions that are 2-harmonic in $Z \times (0, \infty) = X$ are also 2-harmonic in this modified space, and finally that Z is isometric to the boundary of this modified space.

Theorem 1.14. Let (Z, d, ν) be a compact doubling metric measure space supporting a 2-Poincaré inequality, and let $X = Z \times (0, \infty)$. There is a conformal transformation of X to a metric space Ω with transformed metric d_ρ , together with a natural transformation ν_ω of the measure ν , so that $(\Omega, d_\rho, \nu_\omega)$ is a John domain in its completion $\bar{\Omega}$ and so that $(\bar{\Omega}, d_\rho, \nu_\omega)$ satisfies our hypotheses (H0), (H1) and (H2). Moreover, Z is isometric to $\partial\Omega$, and the fractional Laplacian $(-\Delta_2)^\theta$ as constructed in Theorem 1.12 above agrees with the construction given in [33].

This theorem will be proved in Section 7, see Subsection 7.5.

Remark 1.15. The fractional Neumann boundary value problem considered in [22, (1.6)] for bounded Euclidean domains U correspond to the solutions constructed in [33] for the domain $\Omega = \bar{U} \times (0, \infty)$. Theorem 1.14 shows then that the problem studied in [22, (1.6)] corresponds to the problem studied in the present paper, in the special setting of $p = 2$ and in the Euclidean setting. We point out that the Hölder regularity result [22, Theorem 1.2] requires the same integrability condition on the nonhomogenous data f as that of Theorem 1.6.

In the smooth setting of Euclidean spaces, Carnot groups and asymptotically hyperbolic Riemannian manifolds, there is now a vast literature on fractional powers of the Laplacian operator, and [8, 20, 21, 22, 24, 61] is merely a small sampling of the current literature; these all are associated with the fractional powers of the *linear* operator (i.e., $p = 2$). The literature in more general non-smooth setting is more limited; we refer interested readers to [6, 7, 10, 29, 33, 34, 41] as well as the references listed therein for the linear setting. For symmetry results related to some other problems involving fractional p -Laplacian in the setting of Euclidean and Heisenberg group settings and for the use of the method of moving planes in those settings, we refer the interested reader to [5, 26, 62] and the references listed therein. Analysis of non-local (linear, that is, $p = 2$) operators was undertaken in the papers [27, 28, 29] from the point of view of jump processes and more general α -stable processes (recall that for us, $\alpha = 2\theta$) using the language of Dirichlet forms. In [27], Chen and Kumagai consider such processes on Ahlfors regular complete metric measure spaces, and prove Hölder regularity, see for example [27, Theorem 4.14]. In [28] they extend this study to doubling metric measure spaces where a notion of uniform doubling property is also assumed; this additional property is removed in the recent paper [29]. Our approach is more aligned with the approach of [20], as that approach is adaptable also to the nonlinear ($p \neq 2$) setting.

2. BACKGROUND

In this section we gather together the needed background used in the paper. The triple (X, d, μ) denotes a complete metric measure space with μ a Radon measure. We now list some basic notions associated with the theory of analysis on metric measure spaces. For $x \in X$ and $r > 0$, the ball centered at x and with radius r is denoted $B(x, r)$, that is, $B(x, r)$ consists of all the points $y \in X$ for which $d(x, y) < r$. The closed ball $\overline{B}(x, r)$ consists of all the points $y \in X$ for which $d(x, y) \leq r$. Note that in general $\overline{B}(x, r)$ could be a larger set than the topological closure of the open ball $B(x, r)$, but if X is a length space these two sets are the same.

2.1. John and uniform domains. Recall that we assume X to be a complete metric space. We say that a domain $\Omega \subset X$ is a *John domain* if there is a point $x_0 \in \Omega$, called a John center, and a John constant $C_J \geq 1$ such that whenever $x \in \Omega$, there is a rectifiable curve γ_x in Ω with end points x_0 and x such that for each point z in the trajectory of γ_x we have that

$$\text{dist}_{X \setminus \Omega}(z) \geq C_J^{-1} \ell(\gamma_x[x, z]),$$

where $\gamma_x[x, z]$ denotes the segments of γ_x with end points z and x . Clearly a John domain is a connected open set, and moreover, if $\Omega \neq X$ then Ω is bounded. In this paper we also refer to a narrower class of domains, called *uniform domains*, characterized by the existence of a constant $C_U \geq 1$ such that for every pair $x, y \in \Omega$ there exists a rectifiable curve γ_{xy} joining them,

with the property

$$\text{dist}_{X \setminus \Omega}(z) \geq C_U^{-1} \min \left(\ell(\gamma_{xy}[x, z]), \ell(\gamma_{xy}[z, y]) \right) \quad \text{and} \quad \ell(\gamma_{xy}) \leq C_U d(x, y),$$

for all $z \in \gamma_{xy}$. Clearly a bounded uniform domain is also John, but the converse is false.

2.2. Measures. We say that the measure μ is *doubling* if there is a constant $C_d \geq 1$ such that

$$0 < \mu(B(x, 2r)) \leq C_d \mu(B(x, r)) < \infty$$

for each $x \in X$ and $r > 0$. Doubling measures satisfy the following lower mass bound property: there are constants $c > 0$ and $Q > 0$ such that for each $x \in X$, $0 < r < R$, and for each $y \in B(x, R)$,

$$(2.1) \quad c \left(\frac{r}{R} \right)^Q \leq \frac{\mu(B(y, r))}{\mu(B(x, R))},$$

see for example [46, page 76]. The constant c depends solely on the doubling constant C_d .

The measure μ is *Ahlfors Q -regular* for some $Q > 0$ if there is a constant $C \geq 1$ such that for all $x \in X$ and $r > 0$ we have $C^{-1}r^Q \leq \mu(B(x, r)) \leq Cr^Q$. It is now well-known that parts of harmonic analysis can be conducted on doubling metric measure spaces, as described in [60, page 8], though currently there is significant headway in extending the theory of singular integrals beyond doubling spaces. However, much of the theory of quasiconformal maps seems to require μ to be Ahlfors regular, see for instance [45]. If μ is Ahlfors Q -regular, then μ is comparable to the Q -dimensional Hausdorff measure on X .

2.3. Newton-Sobolev functions and Poincaré inequalities. We are interested in using a first-order calculus in metric measure spaces; such a first order calculus was first developed by Heinonen and Koskela in their seminal paper [45] in the process of investigating quasiconformal mappings between Ahlfors regular metric spaces; see also [12, 43, 46, 59]. The idea here is that one needs only the information encoded in magnitude of the gradient of a function in order to conduct much of first-order calculus. Given a measurable function $u : X \rightarrow \mathbb{R}$, we say that a non-negative Borel measurable function g on X is an *upper gradient* of u if

$$|u(x) - u(y)| \leq \int_{\gamma} g \, ds$$

for every non-constant compact rectifiable curve γ in X ; here, x and y denote the terminal points of γ . The function u is said to be in the Dirichlet-Sobolev class $D^{1,p}(X)$ (also known as the homogeneous Sobolev class) if u has an upper gradient that belongs to $L^p(X)$; and u is said to be in the Newton-Sobolev class $N^{1,p}(X)$ if it is in $D^{1,p}(X)$ and in addition, u itself belongs

to $L^p(X)$. Given that upper gradients are not unique, we set the energy semi-norm on $D^{1,p}(X)$ by

$$\mathcal{E}_p(u)^p := \inf_g \int_X g^p d\mu,$$

where the infimum is over all upper gradients g of u . The norm on $N^{1,p}(X)$ is given by

$$\|u\|_{N^{1,p}(X)} := \|u\|_{L^p(X)} + \mathcal{E}_p(u).$$

Indeed, if $1 \leq p < \infty$, for each $u \in D^{1,p}(X)$ there is a unique (up to sets of μ -measure zero) non-negative function g_u that is the L^p -limit of a sequence of upper gradients of u from $L^p(X)$ and so that for each upper gradient g of u we have that $\|g_u\|_{L^p(X)} \leq \|g\|_{L^p(X)}$. The functions g_u belong to a larger class of "gradients" of u , called p -weak upper gradients, see for example [43, 46, 59] or [12].

For $1 \leq p < \infty$, the metric measure space (X, d, μ) is said to support a p -Poincaré inequality if there are constants $C_P > 0$ and $\lambda \geq 1$ such that for all $u \in D^{1,p}(X)$ and balls $B = B(x, r)$ in X , we have

$$\int_{B(x,r)} |u - u_B| d\mu \leq C_P r \left(\int_{B(x,\lambda r)} g_u^p d\mu \right)^{1/p}.$$

It was shown in [43] that if X is a length space, then we can take $\lambda = 1$ at the expense of increasing the constant C_P .

For the rest of this section, we consider Ω to be a domain in a complete metric space X . From [1, Section 7] we know that if a metric measure space (Z, d_Z, μ_Z) is locally compact with μ_Z doubling and supports a p -Poincaré inequality for some $1 \leq p < \infty$, then the zero-extension of μ_Z to the completion \overline{Z} of Z also is doubling and supports a p -Poincaré inequality. Moreover, in this case, $N^{1,p}(Z) = N^{1,p}(\overline{Z})$. We will exploit this property in this paper by using the identity $N^{1,p}(\Omega) = N^{1,p}(\overline{\Omega})$ when the restriction of μ to Ω is also doubling and supports a p -Poincaré inequality.

2.4. Besov spaces. Consider a metric measure space (Z, d, ν) . For $0 < \theta < 1$ and $1 < p < \infty$ we will consider the following Besov energy:

$$\|u\|_{\theta,p}^p := \int_Z \int_Z \frac{|u(y) - u(x)|^p}{d(x,y)^{\theta p} \nu(B(x,d(x,y)))} d\nu(y) d\nu(x),$$

and set $B_{p,p}^\theta(Z)$ to be the space of all L^p -functions for which this energy is finite. If ν is Ahlfors Q -regular, then we can replace $\nu(B(x,d(x,y)))$ with $d(x,y)^Q$ to obtain an equivalent energy. The following lemma holds also for this modified norm. The homogeneous Besov space $HB_{p,p}^\theta(\partial\Omega)$ is the collection of equivalence classes of functions from $B_{p,p}^\theta(\partial\Omega)$, where two functions $u, v \in B_{p,p}^\theta(\partial\Omega)$ are said to be equivalent if $\|u - v\|_{\theta,p} = 0$, that is, $u - v$ is ν -a.e. constant on $\partial\Omega$. By a slight abuse of notation, we conflate equivalence classes that are elements of $HB_{p,p}^\theta(\partial\Omega)$ with representative functions in those classes, but are careful to remember that then there is an ambiguity up to additive constants here.

Lemma 2.2. *Let Z be a bounded metric space equipped with a measure ν with $\nu(Z) < \infty$. If $f \in HB_{p,p}^\theta(Z)$, then $f \in L^p(Z)$. Moreover, there exists $C > 0$ such that for all $f \in HB_{p,p}^\theta(Z)$ we have that*

$$(2.3) \quad \|f - f_Z\|_{L^p(Z)} \leq C \|f\|_{\theta,p}.$$

Here $f_Z = \int_Z f d\nu$. In particular, $HB_{p,p}^\theta(\partial\Omega)$ is a reflexive Banach space under the norm $\|\cdot\|_{\theta,p}$, and is a Hilbert space when $p = 2$.

Proof. Since

$$\|f\|_{\theta,p}^p = \int_Z \int_Z \frac{|f(y) - f(x)|^p}{\nu(B(y, d(x, y))) d(x, y)^{\theta p}} d\nu(y) d\nu(x) < \infty,$$

there exists $x_0 \in Z$ such that $|f(x_0)| < \infty$ and

$$C := \int_Z \frac{|f(y) - f(x_0)|^p}{\nu(B(y, d(x_0, y))) d(x_0, y)^{\theta p}} d\nu(y) < \infty.$$

Thus, we have that

$$\frac{1}{\nu(Z) \text{diam}(Z)^{\theta p}} \int_Z |f(y) - f(x_0)|^p d\nu(y) \leq C.$$

Since Z is bounded and $\nu(Z) < \infty$, it follows that

$$\begin{aligned} \int_Z |f(y)|^p &\leq \int_Z (|f(y) - f(x_0)| + |f(x_0)|)^p d\nu(y) \\ &\leq 2^p \left(\int_Z |f(y) - f(x_0)|^p d\nu(y) + |f(x_0)|^p \nu(Z) \right) \\ &\leq 2^p \left(C \nu(Z) \text{diam}(Z)^{\theta p} + |f(x_0)|^p \nu(Z) \right) < \infty. \end{aligned}$$

Next, by the Hölder inequality and by the boundedness of Z ,

$$\begin{aligned} \int_Z |f(y) - f_Z|^p d\nu(y) &= \int_Z \left| f(y) - \nu(Z)^{-1} \int_Z f(x) d\nu(x) \right|^p d\nu(y) \\ &\leq \nu(Z)^{-1} \int_Z \int_Z |f(y) - f(x)|^p d\nu(x) d\nu(y) \\ &\leq \text{diam}(Z)^{\theta p} \int_Z \int_Z \frac{|f(y) - f(x)|^p}{d(x, y)^{\theta p} \nu(B(x, d(x, y)))} d\nu(x) d\nu(y) \\ &= \text{diam}(Z)^{\theta p} \|f\|_{\theta,p}^p. \end{aligned}$$

Thus we have that $\|f - f_Z\|_{L^p(Z)} \leq \text{diam}(Z)^\theta \|f\|_{\theta,p}$, and so the choice of $C = \text{diam}(Z)^\theta$ proves (2.3).

To see that the homogeneous space is a Banach space, it suffices to show that it is complete under the given norm. Let $\{u_k\}$ be a Cauchy sequence in $HB_{p,p}^\theta(Z)$. Replacing u_k with $u_k - \int_Z u_k d\nu$ if needed, we may assume that $(u_k)_Z := \int_Z u_k d\nu = 0$ for each k . Then by (2.3) we know that $\{u_k\}$ is a Cauchy sequence also in $L^p(Z)$, and hence converges to some function $u \in L^p(Z)$. By passing to a subsequence if necessary, we may also assume that this convergence also occurs pointwise ν -a.e. in Z . Note that $\|u_k\|_{\theta,p} =$

$\|v_k\|_{L^p(Z \times Z, \nu_0)}$ where ν_0 is the weighted measure on $Z \times Z$ given by [Changed the next line according to report. Nages]

$$\nu_0(A) = \iint_A \frac{1}{d(x, y)^{\theta p} \nu(B(x, d(x, y)))} d(\nu \times \nu)(x, y),$$

and

$$v_k(x, y) = u_k(x) - u_k(y).$$

It follows that $\{v_k\}$ is also a Cauchy sequence, and hence converges to some function $v \in L^p(Z \times Z, \nu_0)$. By considering a further subsequence, we can also ensure that this convergence is also ν_0 -a.e. in $Z \times Z$. Now, by applying Fubini's theorem, we also know that v_k converges pointwise ν -a.e. in Z to the function $(x, y) \mapsto u(x) - u(y)$. That is, the $L^p(Z \times Z)$ -limit v of v_k is the function given by $v(x, y) = u(x) - u(y)$. Hence we have the desired conclusion that $\|u_k - u\|_{\theta, p} \rightarrow 0$ as $k \rightarrow \infty$ and $u \in HB_{p,p}^\theta(Z)$ as desired. Since $1 < p < \infty$, it is clear from the reflexivity of $L^p(Z \times Z, \nu_0)$ that $HB_{p,p}^\theta(Z)$ is also reflexive.

Since the norm on $HB_{2,2}^\theta(Z)$ is given via an inner product \mathcal{E}_2 , and as the above argument tells us that $HB_{2,2}^\theta(Z)$ is complete, we conclude that it is a Hilbert space. \square

We will show below that any doubling, compact metric measure space (Z, d, ν) arises as boundary $Z = \partial\Omega$, with Ω a space satisfying the structure conditions (H0), (H1), (H2), and such that ν satisfies the comparison (1.1). Such co-dimension condition arises in the study of traces of the $N^{1,p}$ and $D^{1,p}$ -classes on Ω to $\partial\Omega$ (see for instance [14, 54, 55]), and is a natural condition that arises from considering Ω to be a uniformization of a hyperbolic filling of a doubling compact metric measure space as in [14]. Besov spaces on $\partial\Omega$ arise as trace classes of Newton-Sobolev spaces on Ω . The following result is from [54], but in the setting of Ω such a trace result can also be found in [14].

We equip the Besov space $B_{p,p}^\theta(\partial\Omega)$ with the norm

$$\|u\|_{B_{p,p}^\theta(\partial\Omega)} := \|u\|_{L^p(\partial\Omega)} + \|u\|_{\theta, p}.$$

Theorem 2.4 ([54, Theorem 1.1]). *Let Ω be a John domain in X , with $\overline{\Omega}$ compact such that the restriction of μ to $\overline{\Omega}$ is doubling and supports a p -Poincaré inequality. Suppose in addition that $\partial\Omega$ is equipped with a measure ν that satisfies (1.1). Then there is a bounded linear trace operator $T : N^{1,p}(\Omega) \rightarrow B_{p,p}^{1-\theta/p}(\partial\Omega)$ and a bounded linear extension operator $E : B_{p,p}^{1-\theta/p}(\partial\Omega) \rightarrow N^{1,p}(\Omega)$ such that*

- (i) $T \circ Eu = u$ for $u \in B_{p,p}^{1-\theta/p}(\partial\Omega)$,
- (ii) for each $u \in N^{1,p}(\Omega)$, for \mathcal{H} -a.e. $x \in \partial\Omega$ we have

$$\lim_{r \rightarrow 0^+} \int_{B(x,r) \cap \Omega} |u - Tu(x)| d\mu = 0.$$

The domains Ω considered in this paper satisfy the hypotheses given in Theorem 2.4. Note that if Ω is a uniform domain in X and (X, d, μ) is doubling and supports a p -Poincaré inequality, then the restrictions of μ

to Ω and to $\overline{\Omega}$ are doubling and support a p -Poincaré inequality, see for example [15, Theorem 4.4].

2.5. Cheeger differential structures. We now describe Cheeger (linear) differential structures in doubling metric measure spaces (X, d, μ) ; these structures play a key role in the definition of Cheeger p -harmonic functions.

We say that a metric measure space (X, d, μ) supports a Cheeger differential structure if there is a positive integer N and a collection $\{X_\alpha\}_{\alpha \in A}$, with each X_α a measurable subset of X , such that $\mu(X_\alpha) > 0$ and $\mu(X \setminus \bigcup_\alpha X_\alpha) = 0$, an inner-product structure $\langle \cdot, \cdot \rangle_x$, $x \in X_\alpha$, on \mathbb{R}^N that is μ -measurable, and for each $\alpha \in A$ a Lipschitz map $\varphi_\alpha : X_\alpha \rightarrow \mathbb{R}^N$ satisfying the following condition for each Lipschitz function $f : X \rightarrow \mathbb{R}$: for μ -a.e. $x \in X_\alpha$ there is a vector $\nabla f(x) \in \mathbb{R}^N$ such that

$$\operatorname{ess\,lim\,sup}_{X_\alpha \ni y \rightarrow x} \frac{|f(y) - f(x) - \langle \nabla f(x), \varphi_\alpha(y) - \varphi_\alpha(x) \rangle_x|}{d(y, x)} = 0.$$

For doubling spaces we can assume that A is a countable set and that the collection $(X_\alpha)_{\alpha \in A}$ is pairwise disjoint. For doubling metric measure spaces X supporting a p -Poincaré inequality, such a differential structure was constructed by Cheeger in [25]; the structure constructed there satisfies the additional property that there is a constant $C \geq 1$ such that whenever g_f is a minimal p -weak upper gradient of a Lipschitz function f , then $C^{-1}g_u(x) \leq \langle \nabla f(x), \nabla f(x) \rangle_x \leq Cg_u(x)$ for μ -a.e. $x \in X$. In this paper we will consider such a differential structure. From the discussion in [25] we know that the notion of ∇f extends from the class of locally Lipschitz functions in X to functions in $D^{1,p}(X)$; see also the discussion in [35]. We set $|\nabla f(x)| = \langle \nabla f(x), \nabla f(x) \rangle_x^{1/2}$.

We note that there is more than one possible Cheeger differential structure on X , leading to us considering a wide range of differential operators, one for each such structure. We say that a function $u \in D^{1,p}(\Omega)$ is a *Cheeger p -harmonic function* in Ω if, whenever $v \in D^{1,p}(\Omega)$ has compact support in Ω , we have

$$\int_{\operatorname{supt}(v)} |\nabla u|^p d\mu \leq \int_{\operatorname{supt}(v)} |\nabla(u+v)|^p d\mu.$$

Equivalently, we have the following corresponding Euler-Lagrange equation:

$$\int_{\Omega} |\nabla u(x)|^{p-2} \langle \nabla u(x), \nabla v(x) \rangle_x d\mu(x) = 0.$$

For brevity, in our exposition we will suppress the dependence of x on the inner product structure, and denote

$$\langle \nabla u(x), \nabla v(x) \rangle_x = \nabla u(x) \cdot \nabla v(x)$$

when this will not lead to confusion. Cheeger p -harmonic functions are quasiminimizers in the sense of Giaquinta, and hence we can avail ourselves of the properties derived in [49].

2.6. Neumann boundary value problem. In [55] a generalization of the Neumann boundary value problem

$$\begin{aligned}\Delta_p u &= 0 \text{ on } \Omega, \\ |\nabla u|^{p-2} \partial_\eta u &= f \text{ on } \partial\Omega\end{aligned}$$

was constructed and analyzed. There it was assumed that the boundary data f is a bounded measurable function on $\partial\Omega$ such that $\int_{\partial\Omega} f d\nu = 0$. Strictly speaking, the version considered in [55] is the problem of minimizing the energy operator

$$N^{1,p}(\Omega) \ni v \mapsto I_f(v) := \int_{\overline{\Omega}} g_v^p d\mu - p \int_{\partial\Omega} v f d\nu,$$

with g_v the minimal p -weak upper gradient of v and $f \in L^\infty(\overline{\Omega}, \nu)$ in addition. However, the proofs given in [55] are robust and apply also to the Cheeger differential formulation considered in the current paper, and we will use the results from [55] here, namely the boundedness property of solutions to the Neumann problem. For bounded Neumann data f , the existence of solutions was established in [55], and it was also shown that solutions are bounded. In the appendix, we show that the boundedness of the Neumann data f is not needed in order to obtain existence and boundedness of solutions. In fact, we have the following result as a consequence of the discussion in Lemma A.12 from the appendix together with the results from [55] (for existence of the minimizers).

Theorem 2.5. *Let Ω satisfy the structural assumptions (H0), (H1), and (H2). Then for each $f \in L^p(\partial\Omega)$ with $\int_{\partial\Omega} f d\nu = 0$ there is a bounded function $u \in N^{1,p}(\Omega)$ such that whenever $v \in N^{1,p}(\Omega)$, we have*

$$\int_{\Omega} |\nabla u|^p d\mu - p \int_{\partial\Omega} u f d\nu \leq \int_{\Omega} |\nabla v|^p d\mu - p \int_{\partial\Omega} v f d\nu.$$

Moreover, we can choose u so that $\int_{\Omega} u d\mu = 0$.

2.7. Hyperbolic fillings. In this subsection we give a brief description of the hyperbolic filling of a compact doubling metric measure space as given in [14]. While this construction differs somewhat from the constructions given in earlier literature, in essence the metric portion of the construction is similar to that of [16, 17, 18, 19].

With (Z, d) a compact metric space equipped with a doubling measure ν , we fix $\alpha > 1$, $\tau > 1$, and, for each non-negative integer n we choose a maximal α^{-n} -separated set $S_n \subset Z$. By scaling the metric if need be, we can always assume that the diameter of Z is smaller than 1; hence S_0 contains only one point x_0 . We can also ensure that $S_n \subset S_{n+1}$, and consider the vertex set $V = \bigcup_{n=0}^{\infty} S_n \times \{n\}$. Two points (x, n) and (y, m) are declared to be neighbors if either $B_Z(x, \alpha^{-n})$ intersects $B_Z(y, \alpha^{-m})$ with $|n - m| = 1$, or if $B_Z(x, \tau\alpha^{-n}) \cap B_Z(y, \tau\alpha^{-m})$ is non-empty with $n = m$; here B_Z denotes a ball in the metric space Z . This converts V into a graph X , with each edge assigned a unit length interval. As shown in [14], the graph X , equipped

with the path metric, is a roughly star-like Gromov-hyperbolic space, which, when uniformized via the metric

$$d_\varepsilon(v, w) = \inf_{\gamma} \int_{\gamma} e^{-\varepsilon d(\gamma(t), v_0)} ds(t), \quad v, w \in X,$$

where $v_0 = (x_0, 0)$ and the infimum is over all curves in X with end points v, w , turns the metric graph (X, d) into a uniform domain $X_\varepsilon := (X, d_\varepsilon)$ when $\varepsilon = \log(\alpha)$. The metric graph has a natural measure on it, given by considering the one-dimensional Hausdorff measure \mathcal{H}^1 on the edges of the graph. Next, following [14], for each $\beta > 0$, one can lift the measure ν on Z up to a measure μ_β in X by setting for each Borel set $A \subset X$,

$$\mu_\beta(A) := \int_A e^{-\beta d(x, v_0)} \nu(B_Z(\zeta_x, \alpha^{-n_x})) d\mathcal{H}^1(x),$$

where $(\zeta_x, n_x) \in V$ is a nearest vertex to the point $x \in X$. It is shown that with $\varepsilon = \log(\alpha)$ and $\beta > 0$, the metric measure space $(X, d_\varepsilon, \mu_\beta)$ is doubling and supports a 1-Poincaré inequality; moreover, Z is biLipschitz to the boundary $\partial_\varepsilon X$ of the uniform domain (X, d_ε) , and in addition, $\Omega := X$ equipped with the metric d_ε and measure μ_β satisfies our structural conditions (H0), (H1), and (H2) with $\Theta = \beta/\varepsilon$. As shown in [14] the trace class of the Sobolev space $N^{1,p}(X, d_\varepsilon, \mu_\beta)$ is the Besov class $B_{p,p}^\theta(Z)$, where $\theta = 1 - \beta/(\varepsilon p) = 1 - \Theta/p$. Moreover, as X is a graph with edges of varying length (as determined by the above uniformization metric d_ε), the upper gradient of a function in $N^{1,p}(X, d_\varepsilon, \mu_\beta)$ is exactly the magnitude of the derivative of the function along each edge. In this setting, we also can choose a natural Cheeger differential structure – differentiation along each edge. Note that the orientation of each edge can be chosen as we please, for the key quantities that are the magnitude of the derivative of a function on the edge and the inner product between the Cheeger derivative of two functions are *independent* of the choice of orientation.

Thus, with our primary object (Z, d, ν) , for each $1 < p < \infty$ and $0 < \theta < 1$, we can choose $\beta = p\varepsilon(1 - \theta)$ and $\varepsilon = \log(\alpha)$ in the above construction to obtain a uniform domain $\Omega = (X, d_\varepsilon)$ equipped with the measure μ_β that satisfies our structural assumptions and yields the fractional Laplacian $(-\Delta_p)^\theta$.

3. EQUIVALENT FORMULATIONS OF THE NEUMANN PROBLEM FOR THE p -LAPLACIAN

In this section we prove Theorem 1.5. Formulation (a), when seen as a problem on the independent metric measure space $\overline{\Omega}$ corresponds to the inhomogeneous problem $-\Delta_p u = \nu$ with ν a (signed) Radon measure on the space of interest such that ν is in the dual of the Sobolev space $W^{1,p}(\Omega)$; see for example [47, 56, 58, 63]. The formulation (b) (with $|\nabla v|$ and with $|\nabla v|$ replaced by the minimal p -weak upper gradient g_v) was considered in [55]. Formulation (c) was motivated by the study of nonlocal minimization problems considered in [20] (Euclidean setting), [34] (Carnot group setting), [8, 24] (manifold setting with $\gamma = 1/2$) and [33] (metric spaces of controlled geometry).

Proof of Theorem 1.5. We first show that (a) is equivalent to (b). To do so, first suppose that (b) holds for u , and let $\phi \in N^{1,p}(\overline{\Omega})$. Then setting for each $\varepsilon \in \mathbb{R}$,

$$I(\varepsilon) := \int_{\Omega} |\nabla(u + \varepsilon\phi)|^p d\mu - p \int_{\partial\Omega} (u + \varepsilon\phi) f d\nu,$$

we know that $I(\varepsilon)$ has a minimum at $\varepsilon = 0$. Therefore $\frac{d}{d\varepsilon} I|_{\varepsilon=0} = 0$. Note that

$$\frac{d}{d\varepsilon} I(\varepsilon) = \frac{p}{2} \int_{\Omega} |\nabla(u + \varepsilon\phi)|^{p-2} 2 \nabla(u + \varepsilon\phi) \cdot \nabla\phi d\mu - p \int_{\partial\Omega} \phi f d\nu.$$

Letting $\varepsilon = 0$ gives

$$\frac{d}{d\varepsilon} I(\varepsilon)|_{\varepsilon=0} = p \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla\phi d\mu - p \int_{\partial\Omega} \phi f d\nu.$$

Setting this equal to zero yields (a).

Next, suppose that (a) holds for u . We will use the following convexity of $|\cdot|^p$ for vectors when $p > 1$: when $\xi, \eta \in \mathbb{R}^N$, and $\langle \cdot, \cdot \rangle$ is an inner product on \mathbb{R}^N , we have

$$|\xi|^p - |\eta|^p \geq p|\eta|^{p-2} \langle \eta, \xi - \eta \rangle$$

where $|\cdot|$ is the norm corresponding to this inner product. Therefore when $v \in N^{1,p}(\overline{\Omega})$,

$$\int_{\Omega} (|\nabla v|^p - |\nabla u|^p) d\mu \geq p \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla(v - u) d\mu.$$

Applying (a) to the function $\phi = v - u$ gives the desired inequality $I(u) \leq I(v)$.

It now only remains to show that (a)+(b) is equivalent to (c). This is a Morrey-type argument. For each $\varepsilon > 0$ let η_ε be as in the statement of (c). Suppose that u satisfies (a). Let ϕ be a Lipschitz function on $\overline{\Omega}$. Then by (a) we have that

$$\int_{\partial\Omega} \phi f d\nu = \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla\phi d\mu = \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla((1 - \eta_\varepsilon)\phi) d\mu,$$

where the latter equality follows from the fact that $\eta_\varepsilon\phi = 0$ on $\partial\Omega$ and by (a) applied to $\eta_\varepsilon\phi$. Note that the Cheeger differential structure also follows the Leibniz rule (see for example [25, (4.43)] or [41]), and so $\nabla(1 - \eta_\varepsilon)\phi = (1 - \eta_\varepsilon)\nabla\phi - \phi\nabla\eta_\varepsilon$. Since $|\nabla u| \in L^p(\Omega)$, we have that

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\Omega} (1 - \eta_\varepsilon) |\nabla u|^{p-2} \nabla u \cdot \nabla\phi d\mu = 0.$$

It follows that

$$\int_{\partial\Omega} \phi f d\nu = - \lim_{\varepsilon \rightarrow 0^+} \int_{\Omega} \phi |\nabla u|^{p-2} \nabla u \cdot \nabla\eta_\varepsilon d\mu.$$

As the above holds for all Lipschitz ϕ on $\overline{\Omega}$, the claim (c) follows.

Finally, suppose that u satisfies (c). This means that

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\Omega} \psi |\nabla u|^{p-2} \nabla u \cdot \nabla\eta_\varepsilon d\mu = - \int_{\partial\Omega} \psi f d\nu$$

whenever ψ is a continuous function on $\overline{\Omega}$. For $\phi \in N^{1,p}(\overline{\Omega})$, the above argument with $\phi = \eta_\varepsilon \phi + (1 - \eta_\varepsilon)\phi$ gives that

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla (\eta_\varepsilon \phi) d\mu = 0$$

because of the p -harmonicity of u in Ω , and so for Lipschitz ϕ on $\overline{\Omega}$,

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla \phi d\mu = - \lim_{\varepsilon \rightarrow 0^+} \int_{\Omega} \phi |\nabla u|^{p-2} \nabla u \cdot \nabla \eta_\varepsilon d\mu = \int_{\partial\Omega} \phi f d\nu,$$

that is, (a) holds true for Lipschitz ϕ . Now, if $v \in N^{1,p}(\overline{\Omega})$, then by the fact that $\overline{\Omega}$ supports a p -Poincaré inequality, there is a sequence of Lipschitz functions ϕ_k on $\overline{\Omega}$ such that $\phi_k \rightarrow v$ in $N^{1,p}(\overline{\Omega})$, see [46, Theorem 8.2.1]. Then as $\nabla \phi_k \rightarrow \nabla v$ in $L^p(\Omega; \mathbb{R}^N)$, we have

$$\lim_{k \rightarrow \infty} \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla \phi_k d\mu = \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla v d\mu.$$

Moreover, as $\phi_k \rightarrow v$ in $N^{1,p}(\overline{\Omega})$ and trace $T : N^{1,p}(\overline{\Omega}) \rightarrow B_{p,p}^\theta(\partial\Omega) \subset L^p(\partial\Omega, \nu)$ is a bounded linear map with f a bounded Borel function on $\partial\Omega$, we obtain also that

$$\lim_{k \rightarrow \infty} \int_{\partial\Omega} \phi_k f d\nu = \int_{\partial\Omega} v f d\nu.$$

Thus (a) follows for all $v \in N^{1,p}(\overline{\Omega})$. \square

4. BOUNDARY REGULARITY FOR $1 < p \leq Q$

The goal of this section is to prove Theorem 1.6. We will only consider the range $1 < p \leq Q$, since in the range $p > Q$ we already know that u is $1 - Q/p$ -Hölder continuous on $\overline{\Omega}$ in view of the Morrey embedding theorem, see [46, Lemma 9.2.12] for instance.

Let $(\overline{\Omega}, d, \mu)$ be a metric measure space satisfying the structural assumptions (H1), (H2) in the introduction. In the following, for $x \in \overline{\Omega}$ and $r > 0$ the balls in the induced metric are $B(x, r) = \{y \in \overline{\Omega} \mid d(y, x) < r\}$. We recall from Subsection 2.4 that in view of [54, Theorem 1.3], there exists a bounded linear trace operator $T : N^{1,p}(\Omega) \rightarrow B_{p,p}^{1-1/p}(\partial\Omega) \subset L^p(\partial\Omega)$. For ease of notation, we will denote Tu , which is a function on $\partial\Omega$, also by u . We recall from the discussion at the end of Subsection 2.3 that because $(\overline{\Omega}, d, \mu|_{\Omega})$ supports a p -Poincaré inequality and $\mu|_{\Omega}$ is doubling, it follows that the extension of u by Tu to $\partial\Omega$ is in $N^{1,p}(\overline{\Omega})$, see for example [48] or [46, Theorem 9.2.8].

Let $f : \partial\Omega \rightarrow \mathbb{R}$ be bounded and Borel measurable, with $\int_{\partial\Omega} f d\nu = 0$ and $f \in L^{p'}(\partial\Omega)$. Suppose that $u \in N^{1,p}(\overline{\Omega})$ is p -harmonic with Neumann boundary conditions; that is, for all $\phi \in N^{1,p}(\overline{\Omega})$,

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla \phi d\mu = \int_{\partial\Omega} \phi f d\nu.$$

Then by [55] or by Theorem 1.5, u is a minimizer of $\int_{\Omega} |\nabla u|^p d\mu - p \int_{\partial\Omega} u f d\nu$. We also require that $\int_{\Omega} u d\mu = 0$. This is always possible by subtracting a constant from any given solution. Note that as we deal with a choice of

Cheeger differential structure, such a solution is unique, see [55, Lemma 4.5 and the subsequent comment].

For $x \in \overline{\Omega}$ and $r > 0$ the boundary of $B(x, r)$ relative to the topology induced by the metric on $\overline{\Omega}$, is $\partial B(x, r) \subset \{y \in \Omega \mid d(x, y) = r\}$.

We begin by recalling the notion of p -harmonic extension and an immediate application of some interior Hölder estimates established in [49, Theorem 5.2].

Proposition 4.1. *Let $u \in N^{1,p}(\overline{\Omega})$, and x, r as above. There exists a unique function $v \in N^{1,p}(\overline{\Omega})$ such that v is p -harmonic in $B(x, r)$ and $v = u$ on $\overline{\Omega} \setminus B(x, r)$. Moreover such v is Hölder continuous in $B(x, r/2)$, and it satisfies the estimate*

$$\sup_{B(x, r/2)} |v - v(x)| \leq C \|u\|_{L^\infty} r^\alpha$$

for some constants C, α depending only on the structure conditions of Ω .

Remark 4.2. We want to emphasize that the function v is the solution to the Dirichlet problem on the domain $\overline{\Omega} \cap B(x, r)$, whose boundary is $\overline{\Omega} \cap \partial B(x, r)$. In other words, for all $\phi \in N_0^{1,p}(\overline{\Omega} \cap B(x, r))$ one has

$$\int_{\overline{\Omega} \cap B(x, r)} |\nabla v|^{p-2} \nabla v \cdot \nabla \phi \, d\mu = 0,$$

whereas for the solution of the Neumann problem u we have instead

$$\int_{\overline{\Omega} \cap B(x, r)} |\nabla u|^{p-2} \nabla u \cdot \nabla \phi \, d\mu = \int_{\partial \Omega \cap B(x, r)} \phi f \, d\nu.$$

Because of this consideration, the interior Hölder estimates in [49] can be applied to v in the metric space $X = \overline{\Omega}$ and the domain $B(x, r) \cap \overline{\Omega}$, thus establishing Hölder continuity of v in $B(x, r) \cap \overline{\Omega}$.

Invoking interior estimates to prove regularity up to the boundary may give pause to readers familiar with the smooth setting. The issue here is that the Dirichlet problem for v in $B(x, r) \cap \overline{\Omega}$ is not the same as a Dirichlet problem in $B(x, r) \cap \Omega$. The test functions used in the former are in $N_0^{1,p}(B(x, r) \cap \overline{\Omega})$ and thus do not need to have zero trace on $\partial \Omega \cap B(x, r)$. In fact the function v does not satisfy the Dirichlet problem (with boundary data u) in $\Omega \cap B(x, r)$, since there is an extra part of the boundary, namely, $B(x, r) \cap \partial \Omega$. On this extra part of the boundary, the interested reader should note that, in the smooth Riemannian setting, v satisfies zero Neumann boundary condition. A similar perspective can be found in the discussion on orbifolds [11].

The following proposition gives Morrey type bounds on the growth of the averages of $|\nabla u|^p$ on balls near points on the boundary $\partial \Omega$. This bound will be used in Theorem 1.6 to show Hölder continuity of u near these points. Recall that the measure ν is Θ -codimensional with respect to μ for some $0 < \Theta \leq 1$, see (1.1) above.

Proposition 4.3. *Suppose that $f \in L^q(\partial \Omega) \cap L^{p'}(\partial \Omega)$ for some $q > 1$. There exists a constant \mathbb{M} , depending only on q and the structure conditions (H1),*

(H2), such that when u is a solution to the Neumann boundary value problem with boundary data f , then for every $x_0 \in \partial\Omega$, and every $x \in \bar{\Omega}$ and $r > 0$ such that $B(x, 2r) \subset B(x_0, R)$ with $R < \text{diam}(\Omega)/2$, we have

$$\int_{B(x, r/4) \cap \Omega} |\nabla u|^p d\mu \leq \mathbb{M} \left[M^p \frac{\mu(B(x, r))}{r^{(1-\alpha)p}} + M \left(\int_{B(x_0, R) \cap \partial\Omega} |f|^q d\nu \right)^{1/q} \frac{\mu(B(x, r))^{1/q'}}{r^{\Theta/q'}} \right].$$

Here, $q' = q/(q-1)$, α is as in Proposition 4.1, and $M = \sup_{\Omega} |u|$.

For ease, we adopt the following convention in the proof: when notating the integral over a ball B with respect to μ and ν , we mean integration over $B \cap \Omega$ and $B \cap \partial\Omega$ respectively.

Proof. Consider a p -harmonic extension v of u from $\bar{\Omega} \setminus B(x, r)$ to $B(x, r) \cap \Omega$, as in Proposition 4.1. Then $v \in N^{1,p}(\bar{\Omega})$ with $u - v \in N_0^{1,p}(B(x, r))$.

First, from the convexity of the function $t \mapsto t^p$ for $p > 1$,

$$(4.4) \quad \int_{B(x, r/4)} |\nabla u|^p d\mu \leq 2^{p-1} \left(\int_{B(x, r/4)} |\nabla(u-v)|^p d\mu + \int_{B(x, r/4)} |\nabla v|^p d\mu \right).$$

Let us first estimate the first integral on the right hand side. The p -Laplace operator satisfies the following structural conditions:

$$(4.5) \quad (|z|^{p-2}z - |w|^{p-2}w) \cdot (z - w) \geq \begin{cases} C'|z - w|^p, & p \geq 2 \\ C''(|z| + |w|)^{p-2}|z - w|^2, & p \leq 2. \end{cases}$$

Here the constants C' and C'' only depend on p .

When $p \geq 2$, the use of (4.5) gives us

$$\int_{B(x, r/4)} |\nabla(u-v)|^p d\mu \leq C \int_{B(x, r/4)} (|\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v) \cdot (\nabla u - \nabla v) d\mu.$$

Combining this with (4.4) gives

$$(4.6) \quad \begin{aligned} \int_{B(x, r/4)} |\nabla u|^p d\mu &\leq C \int_{B(x, r/4)} (|\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v) \cdot (\nabla u - \nabla v) d\mu \\ &\quad + C \int_{B(x, r/4)} |\nabla v|^p d\mu. \end{aligned}$$

For $p < 2$, we first use Young's inequality with exponents $2/p$ and $2/(2-p)$ and then (4.5) to obtain

$$\begin{aligned}
& \int_{B(x,r/4)} |\nabla(u-v)|^p d\mu \\
&= \int_{B(x,r/4)} \tau^{(p-2)/2} |\nabla(u-v)|^p (|\nabla u| + |\nabla v|)^{p(p-2)/2} \cdot \tau^{(2-p)/2} (|\nabla u| + |\nabla v|)^{p(2-p)/2} d\mu \\
&\leq \int_{B(x,r/4)} \frac{p\tau^{(p-2)/p}}{2} |\nabla(u-v)|^2 (|\nabla u| + |\nabla v|)^{p-2} + \frac{\tau(2-p)}{p} (|\nabla u| + |\nabla v|)^p d\mu \\
&\leq C\tau^{(p-2)/p} \int_{B(x,r/4)} (|\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v) \cdot (\nabla u - \nabla v) d\mu \\
&\quad + C\tau \int_{B(x,r/4)} (|\nabla u|^p + |\nabla v|^p) d\mu.
\end{aligned} \tag{4.7}$$

By choosing τ small enough so that $C\tau \leq 1/2$ we can absorb the integral of $|\nabla u|^p$ to the left hand side of (4.4) to obtain (4.6) even for the case $1 < p < 2$, that is,

$$\begin{aligned}
& \int_{B(x,r/4)} |\nabla u|^p d\mu \leq C \int_{B(x,r)} (|\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v) \cdot (\nabla u - \nabla v) d\mu \\
&\quad + C \int_{B(x,r/4)} |\nabla v|^p d\mu.
\end{aligned} \tag{4.8}$$

Notice that all the integrands are nonnegative and thus integrating over $B(x,r)$ instead of $B(x,r/4)$ can only increase the integrals.

Let us first consider the first term on the right hand side. Using $u-v$ as a test function, we have from the weak formulation of the p -Laplacian equation 1.5(a) (with Ω replaced by $B(x,r)$ and f replaced by the constant function 0 on $B(x,r)$; recall that v is a solution to the Dirichlet problem on $B(x,r) \cap \overline{\Omega}$ with boundary data u on $\Omega \cap \partial B(x,r)$) that

$$\int_{B(x,r)} |\nabla v|^{p-2} \nabla v \cdot \nabla(u-v) d\mu = 0.$$

Hence, integrating over the ball $B(x,r)$ we get

$$\begin{aligned}
& \int_{B(x,r)} (|\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v) \cdot (\nabla u - \nabla v) d\mu \\
&= \int_{B(x,r)} |\nabla u|^{p-2} \nabla u \cdot (\nabla u - \nabla v) - |\nabla v|^{p-2} \nabla v \cdot (\nabla u - \nabla v) d\mu \\
&= \int_{B(x,r)} |\nabla u|^{p-2} \nabla u \cdot (\nabla u - \nabla v) d\mu.
\end{aligned}$$

Now, since u is a solution to the Neumann problem with boundary data f , we obtain

$$\int_{B(x,r)} |\nabla u|^{p-2} \nabla u \cdot \nabla(u-v) d\mu = \int_{B(x,r)} (u-v) f dv.$$

Hence, invoking the L^∞ bounds on solutions, and the regularity assumption (H2) of the measure ν , one has

$$\begin{aligned}
 \int_{B(x,r)} (u-v)f \, d\nu &\leq \int_{B(x,r)} (|u| + |v|)|f| \, d\nu \\
 &\leq 2CM \int_{B(x,r)} |f| \, d\nu \\
 &\leq 2CM \nu(B(x,r))^{1/q'} \left(\int_{B(x,r) \cap \partial\Omega} |f|^q \, d\nu \right)^{1/q} \\
 (4.9) \quad &\leq 2CM \left(\int_{B(x_0,R)} |f|^q \, d\nu \right)^{1/q} \frac{\mu(B(x,r))^{1/q'}}{r^{\Theta/q'}}.
 \end{aligned}$$

Note that the constant C above changes from line to line, but only depends on p and the regularity constant of the measure ν . Here we have used the fact that u , and hence by maximum principle, v are bounded in $B(x_0, R)$, see [55]. An upper bound for $|u| + |v|$ in $B(x_0, R)$ is denoted by M .

We now consider the second term of the sum on the right hand side of equation (4.8).

Since v is p -harmonic in $B(x, r)$, then it is in the De Giorgi class $DG_p(B(x, r))$, see [49, Proposition 3.3]. Hence for all $k \in \mathbb{R}$,

$$\int_{B(x,r/4)} |\nabla(v-k)_\pm|^p \, d\mu \leq \frac{C}{(r/2 - r/4)^p} \int_{B(x,r/2)} (v-k)_\pm^p \, d\mu.$$

Here, for a function h , the function $h_+ = \max\{h, 0\}$ is the positive part of h and $h_- = \max\{-h, 0\}$ is the negative part of h . Hence, choosing $k = v(x)$, we have

$$\int_{B(x,r/4)} |\nabla(v-v(x))_+|^p \, d\mu \leq \frac{C}{r^p} \int_{B(x,r/2)} (v-v(x))_+^p \, d\mu$$

and

$$\int_{B(x,r/4)} |\nabla(v-v(x))_-|^p \, d\mu \leq \frac{C}{r^p} \int_{B(x,r/2)} (v-v(x))_-^p \, d\mu.$$

Summing these two inequalities, and recalling that $B(x, r/2) \subset B(x, \frac{3}{2}r) \subset \Omega$, we can then invoke the scale invariant *local* α -Hölder continuity estimates for p -harmonic functions from Proposition 4.1 (originally in [49, Theorem 5.2]), obtaining

$$\begin{aligned}
 \int_{B(x,r/4)} |\nabla v|^p \, d\mu &\leq \frac{C}{r^p} \int_{B(x,r/2)} |v-v(x)|^p \, d\mu \leq \frac{C M^p}{r^p} \int_{B(x,r/2)} r^{\alpha p} \, d\mu \\
 (4.10) \quad &\leq C M^p \frac{\mu(B(x,r))}{r^{(1-\alpha)p}}.
 \end{aligned}$$

Combining inequalities (4.4), (4.9), and (4.10), we finally conclude

$$\begin{aligned} \int_{B(x, r/4)} |\nabla u|^p d\mu &\leq C M^p \frac{\mu(B(x, r))}{r^{(1-\alpha)p}} \\ &\quad + 2CM \left(\int_{B(x_0, R) \cap \partial\Omega} |f|^q d\nu \right)^{1/q} \frac{\mu(B(x, r))^{1/q'}}{r^{\Theta/q'}}. \quad \square \end{aligned}$$

Next, we establish the global Hölder continuity of u in $\bar{\Omega}$. The argument is along the lines of the standard proof of Morrey embedding theorem found in [43], and is a streamlined version of the classical regularity proof found in PDE texts such as [38, 39]. This classical proof is in two parts: the first part is to show that functions whose gradient exhibit a decay property in the spirit of Proposition 4.3 belong to a Campanato space (see [39, page 43]) by using Poincaré inequalities, and the second part is to show that functions in the Campanato space are locally Hölder continuous (see [39, page 41]) by using a telescoping sequence of balls and a Lebesgue point argument. Both parts are combined into one seamless argument in the proof given below for the convenience of the reader.

Now we are ready to prove Theorem 1.6.

Proof of Theorem 1.6. Choose R_0 with $0 < R_0 < \text{diam}(\Omega)/2$. For $x_0 \in \partial\Omega$, let $x, y \in B(x_0, R_0/64) \cap \Omega$ be μ -Lebesgue points of u . Then $d(x, y) < R_0/32$, $B(x, R_0/16) \subset B(x_0, R_0)$, and $B(y, R_0/16) \subset B(x_0, R_0)$.

Define a sequence of balls indexed by \mathbb{Z} centered at x and y as follows: For $i \geq 0$, set $B_i = B(x, 2^{1-i}d(x, y))$, and for $i < 0$, set $B_i = B(y, 2^{1+i}d(x, y))$. We denote the radius of the ball B_i by ρ_i ; note that $\rho_i = 2^{1-|i|}d(x, y)$. For each ball in the sequence, we can apply Proposition 4.3 with $r = 4\rho_i$.

By an application of the Poincaré inequality, the doubling condition and the Lebesgue point property of x, y for u with respect to the measure μ , we then estimate

$$\begin{aligned} |u(x) - u(y)| &\leq \sum_{i \in \mathbb{Z}} |u_{B_i} - u_{B_{i+1}}| \leq C \sum_{i \in \mathbb{Z}} \int_{B_i} |u - u_{B_i}| d\mu \\ &\leq C \sum_{i \in \mathbb{Z}} \rho_i \left(\mu(B_i)^{-1} \int_{B_i} |\nabla u|^p d\mu \right)^{1/p}. \end{aligned}$$

Therefore, invoking Proposition 4.3, applied to $4B_i$ for each i , one obtains

$$\begin{aligned}
|u(x) - u(y)| &\leq C \sum_{i \in \mathbb{Z}} \frac{\rho_i}{\mu(B_i)^{1/p}} \left[\frac{\mu(B_i)^{1/p}}{\rho_i^{1-\alpha}} + \frac{\mu(B_i)^{1/(q'p)}}{\rho_i^{\Theta/(q'p)}} \right] \\
(4.11) \quad &= C \sum_{i \in \mathbb{Z}} \left[\rho_i^\alpha + \frac{\rho_i^{1-\Theta/(q'p)}}{\mu(B_i)^{(1-1/q')/p}} \right] \\
(4.12) \quad &\leq C \sum_{i \in \mathbb{Z}} \left[\rho_i^\alpha + \left(\frac{R_0}{\mu(B(x_0, R_0))} \right)^{1/(pq)} \rho_i^{1-\Theta/(q'p)-Q/(qp)} \right] \\
&\leq C \sum_{i \in \mathbb{Z}} \rho_i^{1-\varepsilon} = Cd(x, y)^{1-\varepsilon} \sum_{i \in \mathbb{Z}} 2^{-|i|(1-\varepsilon)} \leq Cd(x, y)^{1-\varepsilon}.
\end{aligned}$$

Here inequality (4.12) follows from the lower mass bound property of μ , see (2.1). The constant C depends only on the structural constants as well as on $\mu(B(x_0, R_0))$, R_0 and $\int_{B(x_0, R_0)} |f|^q d\nu$. This completes the proof of Hölder's inequality.

Finally, to prove the Harnack inequality, we assume that $u \geq 0$ on Ω and that $f = 0$ on the relatively open set $W \subset \partial\Omega$. By Theorem 1.5(a) we have that whenever $\varphi \in N^{1,p}(\Omega) = N^{1,p}(\overline{\Omega})$ with support contained in $\Omega \cup W$, we have $\int_{\Omega \cup W} |\nabla u|^{p-2} \nabla u \cdot \nabla \varphi d\mu = 0$, and hence it follows that u is Cheeger p -harmonic in the domain $\Omega \cup W$, seen as a domain in the metric space $\overline{\Omega}$. By the strong maximum principle as in [49, Corollary 6.4], we then have that either u is identically zero in $\overline{\Omega}$ (and hence satisfies the Harnack inequality trivially), or else, $u > 0$ in Ω . In this latter case, invoking [49, Corollary 7.3], the desired Harnack inequality follows. \square

Remark 4.13. Although we were able to prove L^∞ bounds for u from the hypothesis $f \in L^{p'}(\partial\Omega, \nu)$ (see the appendix), in the argument above we need the stronger integrability condition $f \in L^q(\partial\Omega, \nu)$ with $Q - \Theta < (p - \Theta)q$. If we only have that $f \in L^{p'}(\partial\Omega)$ then the proof can be modified, so that one obtains a weaker regularity condition. Indeed, in this case, if $x, y \in \partial\Omega$ such that the maximal function $M(f^{p'})$ is finite at those points, then

$$|u(x) - u(y)| \leq Cd(x, y)^{1-\beta} \left[1 + M(f^{p'})(x)^{1/p'} + M(f^{p'})(y)^{1/p'} \right],$$

where $\beta = \max\{1 - \alpha, \Theta/p\}$, which is automatically smaller than 1 as $p > \Theta$.

5. STABILITY OF SOLUTIONS UNDER PERTURBATION OF THE NEUMANN DATA

In this section we prove two different stability results for p -Laplacian Neumann problems in the metric measure space setting, under $L^{p'}$ perturbations of the boundary data, with $p > 1$ and p' the Hölder conjugate of p . The first is Theorem 1.9. The second stability result corresponds to the formulation of the Neumann problem using only minimal upper gradients, as in [55] (i.e., the variational formulation in Theorem 1.5 part (b), with $|\nabla u|$ substituted by the minimal upper gradient,) and it is stated in Theorem 5.4.

We first begin by proving Theorem 1.9.

Proof of Theorem 1.9. Let f, g, u, v be as in the statement of the theorem. Since u and v are solutions of the Neumann problems with boundary data f and g , we have that for all $\phi \in N^{1,p}(\bar{\Omega})$,

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla \phi \, d\mu = \int_{\partial\Omega} \phi f \, d\nu \quad \text{and} \quad \int_{\Omega} |\nabla v|^{p-2} \nabla v \cdot \nabla \phi \, d\mu = \int_{\partial\Omega} \phi g \, d\nu.$$

Subtracting the second identity from the first and substituting $\phi = u - v$, yields

$$\begin{aligned} \int_{\Omega} (|\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v) \cdot (\nabla u - \nabla v) \, d\mu &= \int_{\partial\Omega} (f - g)(u - v) \, d\nu \\ &\leq \|u - v\|_{L^p(\partial\Omega)} \|f - g\|_{L^{p'}(\partial\Omega)}. \end{aligned}$$

We first consider the case $p \geq 2$. In this case, we invoke the monotonicity (4.5) to obtain

$$(5.1) \quad \|\nabla u - \nabla v\|_{L^p(\Omega)}^p \leq C \|u - v\|_{L^p(\partial\Omega)} \|f - g\|_{L^{p'}(\partial\Omega)}.$$

Now by using the boundedness of the trace operator in Theorem 2.4 and employing Lemma A.3, we obtain

$$\begin{aligned} \|u - v\|_{L^p(\partial\Omega)} &\leq \|u\|_{L^p(\partial\Omega)} + \|v\|_{L^p(\partial\Omega)} \leq C [\|u\|_{N^{1,p}(\Omega)} + \|v\|_{N^{1,p}(\Omega)}] \\ &\leq C [\|\nabla u\|_{L^p(\Omega)} + \|\nabla v\|_{L^p(\Omega)}] \\ &\leq C [\|f\|_{L^{p'}(\partial\Omega)}^{p'/p} + \|g\|_{L^{p'}(\partial\Omega)}^{p'/p}]. \end{aligned}$$

Therefore

$$\|\nabla u - \nabla v\|_{L^p(\Omega)} \leq C \left(\|f\|_{L^{p'}(\partial\Omega)}^{p'/p} + \|g\|_{L^{p'}(\partial\Omega)}^{p'/p} \right)^{1/p} \|f - g\|_{L^{p'}(\partial\Omega)}^{1/p},$$

which yields the desired inequality for the case $p \geq 2$.

On the other hand, if $1 < p < 2$, then we can proceed as in (4.7) and invoke Hölder's inequality and (4.5) to obtain

$$\begin{aligned} &\|\nabla u - \nabla v\|_{L^p(\Omega)}^p \\ &\leq \left(\int_{\Omega} (|\nabla u| + |\nabla v|)^p \, d\mu \right)^{(2-p)/2} \left(\int_{\Omega} |\nabla(u - v)|^2 (|\nabla u| + |\nabla v|)^{p-2} \, d\mu \right)^{p/2} \\ &\leq C \left(\|\nabla u\|_{L^p(\Omega)}^p + \|\nabla v\|_{L^p(\Omega)}^p \right)^{(2-p)/2} \left(\int_{\Omega} (|\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v) \cdot (\nabla u - \nabla v) \, d\mu \right)^{p/2} \\ (5.2) \quad &\leq C \left(\|\nabla u\|_{L^p(\Omega)}^p + \|\nabla v\|_{L^p(\Omega)}^p \right)^{(2-p)/2} \left(\|u - v\|_{L^p(\partial\Omega)} \|f - g\|_{L^{p'}(\partial\Omega)} \right)^{p/2}. \end{aligned}$$

Now, as in the case of $p \geq 2$, we use Theorem 2.4 together with Lemma A.3, but this time also to bound $\|\nabla u\|_{L^p(\Omega)}$ and $\|\nabla v\|_{L^p(\Omega)}$, to obtain

$$\|\nabla u - \nabla v\|_{L^p(\Omega)}^p \leq C \left(\|f\|_{L^{p'}(\partial\Omega)} + \|g\|_{L^{p'}(\partial\Omega)} \right)^{\frac{2-p}{2}p' + \frac{p'}{2}} \|f - g\|_{L^{p'}(\partial\Omega)}^{p/2}. \quad \square$$

Remark 5.3. Theorem 1.9, in combination with the Poincaré inequality (A.2), yields that if $(f_k)_k$ is a sequence of functions in $L^{p'}(\partial\Omega)$ with the condition that $\int_{\partial\Omega} f_k \, d\nu = 0$ for each k , and a function $f \in L^{p'}(\partial\Omega)$ such that $f_k \rightarrow f$

in $L^{p'}(\partial\Omega)$, and if u_k is the solution to the Neumann boundary value problem with boundary data f_k and with $\int_{\Omega} u_k d\mu = 0$, then $u_k \rightarrow u$ in $N^{1,p}(\Omega)$ with u the solution to the Neumann boundary value problem with boundary data f and with $\int_{\Omega} u d\mu = 0$.

Next, we turn to the version of the Neumann problem interpreted as a variational problem involving upper gradients. In this setting there is no Euler-Lagrange equation available and so the above argument would not work. Correspondingly, in this more general setting we obtain a weaker result, in the sense that we only can prove that convex combination of the solutions for data f_k converge to a solution for the limit data $f_k \rightarrow f$. We include this result here to illustrate some of the control we give up by not having access to the Euler-Lagrange equation provided by differential structure ∇u .

Given $u \in N^{1,p}(\Omega)$, and a ν -measurable function $f : \partial\Omega \rightarrow \mathbb{R}$ such that $\int_{\partial\Omega} f d\nu = 0$, consider the energy functional

$$I_f(u) := \int_{\Omega} g_u^p d\mu - p \int_{\partial\Omega} u f d\nu.$$

Here g_u is the minimal p -weak upper gradient of u . Let

$$N_*^{1,p}(\Omega) = \left\{ u \in N^{1,p}(\Omega) : \int_{\Omega} u d\mu = 0 \right\},$$

and define

$$I_{\min}(f) = \inf_{v \in N_*^{1,p}(\Omega)} I_f(v).$$

Theorem 5.4. *For each $k \in \mathbb{N}$, let $f_k \in L^{p'}(\partial\Omega)$ be such that $\int_{\partial\Omega} f_k d\nu = 0$, where p' is the Hölder dual $p/(p-1)$ of p . Let $u_k \in N_*^{1,p}(\Omega)$ be such that $I_{f_k}(u_k) = I_{\min}(f_k)$. Suppose that $f_k \rightarrow f$ in $L^{p'}(\partial\Omega)$. Then $\int_{\partial\Omega} f d\nu = 0$, and there is a convex combination sequence $v_k = \sum_{j=k}^{N(k)} \lambda_{j,k} u_j$ that converges in $N^{1,p}(\Omega)$ to a function $u \in N_*^{1,p}(\Omega)$ and we have that $I_f(u) = I_{\min}(f)$.*

In the above, note that for each k, j we have $0 \leq \lambda_{j,k} \leq 1$ with $\sum_{j=k}^{N(k)} \lambda_{j,k} = 1$.

Proof. Since $\int_{\partial\Omega} f_k d\nu = 0$ for all $k \in \mathbb{N}$ and $f_k \rightarrow f$ in $L^{p'}(\partial\Omega)$, it is clear that $\int_{\partial\Omega} f d\nu = 0$.

For each k , we have that

$$I_{f_k}(u_k) = I_{\min}(f_k) \leq 0,$$

since the zero function belongs to $N_*^{1,p}(\Omega)$. Thus by [55, Proposition 4.1], there exists $C > 0$ such that for all $k \in \mathbb{N}$,

$$0 \geq I_{f_k}(u_k) \geq \int_{\Omega} g_{u_k}^p d\mu - C \left(\int_{\Omega} g_{u_k}^p d\mu \right)^{1/p} \left(\int_{\partial\Omega} |f_k|^{p'} d\nu \right)^{1/p'}.$$

We then have that

$$\int_{\Omega} g_{u_k}^p d\mu \leq C \int_{\partial\Omega} |f_k|^{p'} d\nu \leq C_0,$$

for some $0 < C_0 < \infty$, since $f_k \rightarrow f$ in $L^{p'}(\partial\Omega)$. Furthermore, since $u_k \in N_*^{1,p}(\Omega)$, it follows from the Sobolev-type inequality [55, (3.6)] that there exists some $C > 0$ such that for all $k \in \mathbb{N}$,

$$\|u_k\|_{L^p(\Omega)} \leq C \|g_{u_k}\|_{L^p(\Omega)}.$$

Thus, $(u_k)_{k \in \mathbb{N}}$ is a bounded sequence in $N^{1,p}(\Omega)$, and so for each $k \in \mathbb{N}$, there is a convex combination

$$v_k := \sum_{j=k}^{N(k)} \lambda_{j,k} u_j, \quad g_k := \sum_{j=k}^{N(k)} \lambda_{j,k} g_{u_j}$$

with $0 \leq \lambda_{j,k} \leq 1$ and $\sum_{j=k}^{N(k)} \lambda_{j,k} = 1$, such that $v_k \rightarrow u$ in $L^p(\Omega)$ and $g_k \rightarrow g$ to $L^p(\Omega)$ where g is some p -weak upper gradient of u , see for instance [59, Lemma 3.6] or [46, Proposition 7.3.7]. From the boundedness of the trace $T : N^{1,p}(\Omega) \rightarrow L^p(\partial\Omega)$,

$$\begin{aligned} \left| \int_{\overline{\Omega}} v_k f \, d\nu - \int_{\partial\Omega} u f \, d\nu \right| &\leq \int_{\overline{\Omega}} |(v_k - u) f| \, d\nu \\ &\leq \|v_k - u\|_{L^p(\partial\Omega)} \|f\|_{L^{p'}(\partial\Omega)} \\ &\leq C \|v_k - u\|_{N^{1,p}(\Omega)} \|f\|_{L^{p'}(\partial\Omega)} \\ &\leq C (\|v_k - u\|_{L^p(\Omega)} + \|g_k - g\|_{L^p(\Omega)}) \|f\|_{L^{p'}(\partial\Omega)} \rightarrow 0 \end{aligned}$$

as $k \rightarrow \infty$. Thus we have that

$$\begin{aligned} I_f(u) &= \int_{\overline{\Omega}} g_u^p \, d\mu - p \int_{\partial\Omega} u f \, d\nu \leq \int_{\overline{\Omega}} g^p \, d\mu - p \int_{\partial\Omega} u f \, d\nu \\ (5.5) \quad &= \lim_{k \rightarrow \infty} \left(\int_{\overline{\Omega}} g_k^p \, d\mu - p \int_{\partial\Omega} v_k f \, d\nu \right). \end{aligned}$$

Let $\varepsilon > 0$. Then, there exists $v_0 \in N_*^{1,p}(\Omega)$ such that

$$I_f(v_0) < I_{\min}(f) + \varepsilon.$$

We note that for any $v \in N_*^{1,p}(\Omega)$,

$$|I_{f_k}(v) - I_f(v)| \leq p \int_{\partial\Omega} |v(f_k - f)| \, d\nu \leq p \|v\|_{L^p(\partial\Omega)} \|f_k - f\|_{L^{p'}(\partial\Omega)} \rightarrow 0$$

as $k \rightarrow \infty$, since $f_k \rightarrow f$ in $L^q(\partial\Omega)$ and $v \in L^p(\partial\Omega)$ by the boundedness of the trace operator. Since $I_{f_k}(u_k) = I_{\min}(f_k)$ for each k , we then have that

$$(5.6) \quad \limsup_{k \rightarrow \infty} I_{f_k}(u_k) \leq \limsup_{k \rightarrow \infty} I_{f_k}(v_0) = I_f(v_0) < I_{\min}(f) + \varepsilon.$$

This also shows that $\lim_{k \rightarrow \infty} I_{\min}(f_k) = I_{\min}(f)$.

By the triangle inequality, we have that

$$\left(\int_{\overline{\Omega}} g_k^p \, d\mu \right)^{1/p} \leq \sum_{j=k}^{N(k)} \lambda_{j,k} \left(\int_{\overline{\Omega}} g_{u_j}^p \, d\mu \right)^{1/p} = \sum_{j=k}^{N(k)} \lambda_{j,k} \left(I_{f_j}(u_j) + p \int_{\partial\Omega} u_j f_j \, d\nu \right)^{1/p}.$$

By (5.6), it follows that for sufficiently large $k \in \mathbb{N}$,

$$\left(\int_{\overline{\Omega}} g_k^p d\mu \right)^{1/p} \leq \sum_{j=k}^{N(k)} \lambda_{j,k} \left(I_{\min}(f) + \varepsilon + p \int_{\partial\Omega} u_j f_j d\nu \right)^{1/p}.$$

By Hölder's Inequality, we have that

$$\begin{aligned} \left(\int_{\overline{\Omega}} g_k^p d\mu \right)^{1/p} &\leq \left(\sum_{j=k}^{N(k)} \lambda_{j,k} \left(I_{\min}(f) + \varepsilon + p \int_{\partial\Omega} u_j f_j d\nu \right) \right)^{1/p} \left(\sum_{j=k}^{N(k)} \lambda_{j,k} \right)^{1/p'} \\ &= \left(\sum_{j=k}^{N(k)} \lambda_{j,k} \left(I_{\min}(f) + \varepsilon + p \int_{\partial\Omega} u_j f_j d\nu \right) \right)^{1/p}. \end{aligned}$$

Therefore it follows that

$$\int_{\overline{\Omega}} g_k^p d\mu \leq I_{\min}(f) + \varepsilon + p \sum_{j=k}^{N(k)} \lambda_{j,k} \int_{\partial\Omega} u_j f_j d\nu.$$

Hence,

$$\begin{aligned} \int_{\overline{\Omega}} g_k^p d\mu - p \int_{\partial\Omega} v_k f d\nu &\leq I_{\min}(f) + \varepsilon + \sum_{j=k}^{N(k)} \lambda_{j,k} p \int_{\partial\Omega} u_j f_j d\nu - p \int_{\partial\Omega} v_k f d\nu \\ &= I_{\min}(f) + \varepsilon + p \sum_{j=k}^{N(k)} \lambda_{j,k} \int_{\partial\Omega} u_j (f_j - f) d\nu \\ &\leq I_{\min}(f) + \varepsilon + p \sum_{j=k}^{N(k)} \lambda_{j,k} \|u_j\|_{L^p(\partial\Omega)} \|f_j - f\|_{L^{p'}(\partial\Omega)} \\ &\leq I_{\min}(f) + \varepsilon + C p \sum_{j=k}^{N(k)} \lambda_{j,k} \|u_j\|_{N^{1,p}(\Omega)} \|f_j - f\|_{L^{p'}(\partial\Omega)}. \end{aligned}$$

Here we again used the boundedness of the trace operator. Since the functions u_k are bounded in $N^{1,p}(\Omega)$ and $f_k \rightarrow f$ in $L^{p'}(\partial\Omega)$, it follows that for sufficiently large k ,

$$\int_{\overline{\Omega}} g_k^p d\mu - p \int_{\partial\Omega} v_k f d\nu < I_{\min}(f) + 2\varepsilon.$$

Therefore by (5.5), we have that $I_f(u) \leq I_{\min}(f) + 2\varepsilon$. By definition, $I_{\min}(f) \leq I_f(u)$. It follows that $I_{\min}(f) = I_f(u)$; that is, u is a solution to the Neumann boundary value problem with boundary data f . \square

6. CONSTRUCTING AN INDUCED NON-LOCAL FRACTIONAL LAPLACIAN FOR COMPACT DOUBLING METRIC MEASURE SPACES

6.1. The general case $1 < p < \infty$. In this section we provide a method to reproduce the strategy in [20] and define an analog of fractional p -Laplacian operators $(-\Delta_p)^\theta$ on doubling metric measure spaces for $1 < p < \infty$ and $0 < \theta < 1$. We recall that in [33] a fractional Laplacian $(-\Delta_2)^\theta$ corresponding

to a Cheeger differential structure on a complete doubling metric measure space supporting a 2-Poincaré inequality was constructed using a method different from the one employed in this paper. In Section 7 below we will show that the definition in [33], gives rise to the same operator we construct in this section.

While in the Euclidean case [20], the $(-\Delta_2)^\theta$ operator in \mathbb{R}^n arises as the Dirichlet-to-Neumann map in the upper half-space \mathbb{R}_+^{n+1} , in our more general setting the operators $(-\Delta_p)^\theta$ are defined in terms of the Dirichlet-to-Neumann map in a hyperbolic filling of (Z, d, ν) , which satisfies the hypotheses (H0), (H1), and (H2).

As outlined in Section 2.7, the hyperbolic filling construction in [14], shows that any compact doubling metric measure space (Z, d, ν) , for any fixed and $1 < p < \infty$ with $0 < \theta < 1$, arises as the boundary of a uniform domain Ω , equipped with a measure μ , such that (Ω, d, μ) is doubling and supports a 1-Poincaré inequality (and hence, a p -Poincaré inequality for each $1 < p < \infty$). The original space Z is biLipschitz equivalent to $\partial\Omega$, and the Besov space $B_{p,p}^\theta(Z)$ is the trace space of the Sobolev class $N^{1,p}(\Omega)$. Indeed, it is shown there that with the same domain Ω , for each choice of $1 < p < \infty$ and $0 < \theta < 1$, there is a choice of measure μ on Ω satisfying the above properties. Moreover, for that choice of measure μ we also have from [14, Theorem 10.3, Theorem 11.3, Theorem 12.1] that when $x \in Z = \partial\Omega$ and $0 < r < 2 \text{diam}(Z)$.

$$\nu(B(x, r)) \approx \frac{\mu(B(x, r))}{r^\Theta} \text{ with } \theta = \frac{p - \Theta}{p},$$

Hence, *every* compact doubling metric measure space (Z, d, ν) arises as the boundary of a metric measure space (Ω, d, μ) with Ω a uniform domain and (Ω, d, μ) doubling metric measure space supporting a p -Poincaré inequality, with the link between the measure μ on Ω and the doubling measure ν on Z given in terms of the codimensionality condition (1.1). Since (Ω, d, μ) satisfies properties (H0), (H1), and (H2), we can then fix a Cheeger differential structure on Ω as described in Section 2, and apply all the results in the previous sections. We will use this notation for the rest of the section.

One natural norm on the Besov class $B_{p,p}^\theta(Z)$ corresponds to a form \mathcal{E}_p given by

$$\mathcal{E}_p(u, v) = \int_Z \int_Z \frac{|u(y) - u(x)|^{p-2} (u(y) - u(x))(v(y) - v(x))}{d(x, y)^{p\theta} \nu(B(y, d(x, y)))} d\nu(x) d\nu(y).$$

Note that when $u, v \in B_{p,p}^\theta(\partial\Omega)$, we have that $\mathcal{E}_p(u, v) \in \mathbb{R}$ and that $\mathcal{E}_p(u, u) = \|u\|_{\theta,p}^p$. Moreover, $\mathcal{E}_p(u, u) = 0$ if and only if u is constant ν -a.e. in $\partial\Omega$. While \mathcal{E}_p is not a bilinear form nor is symmetric in general, it is both bilinear and symmetric when $p = 2$. However, there are other comparable forms in the Besov class, see for example [44, 50]. Given the preceding results of this paper, we have another equivalent form on $B_{p,p}^\theta(Z)$ that is more adapted to seeing this Besov space as a trace space. For $u, v \in B_{p,p}^\theta(Z)$, we

set

$$\mathcal{E}_T(u, v) := \int_{\Omega} |\nabla \widehat{u}|^{p-2} \nabla \widehat{u} \cdot \nabla \widehat{v} d\mu,$$

where $\widehat{u} \in N^{1,p}(\Omega)$ is such that the trace $T\widehat{u} = u$ and \widehat{u} is Cheeger p -harmonic in Ω (that is, \widehat{u} solves the Dirichlet problem for the Cheeger p -Laplacian on Ω with boundary data u). Note that \mathcal{E}_T is bilinear if and only if $p = 2$; otherwise, it is only linear in the second entry.

Lemma 6.1. *There exists $C \geq 1$, depending only on the structure constants, such that for each $u \in B_{p,p}^\theta(\partial\Omega)$, we have*

$$\frac{1}{C} \mathcal{E}_T(u, u) \leq \mathcal{E}_p(u, u) \leq C \mathcal{E}_T(u, u).$$

Proof. From [54, Theorem 1.1], with $T : N^{1,p}(\Omega) \rightarrow B_{p,p}^\theta(\partial\Omega)$ the trace operator and $E : B_{p,p}^\theta(\partial\Omega) \rightarrow N^{1,p}(\Omega)$ the extension operator, we have by the p -harmonicity property of \widehat{u} and by the fact that $T\widehat{u} = TEu$,

$$\mathcal{E}_T(u, u) \leq \int_{\Omega} |\nabla Eu|^p d\mu \leq C \|u\|_{\theta,p}^p = C \mathcal{E}_p(u, u)$$

and

$$\mathcal{E}_p(u, u) = \|T\widehat{u}\|_{\theta,p}^p \leq C \int_{\Omega} |\nabla \widehat{u}|^p d\mu = C \mathcal{E}_T(u, u). \quad \square$$

We are now ready to construct a Cheeger fractional p -Laplacian on Z induced by the Cheeger p -Laplacian on Ω ; recall that Z is seen as the boundary of Ω . This construction is given via the following theorem, and is analogous to the notion of weak Laplacian Δ .

Proposition 6.2. *For each $f \in L^{p'}(Z)$ with $\int_Z f d\nu = 0$ there is a function $u_f \in B_{p,p}^\theta(Z)$ such that for each $\varphi \in B_{p,p}^\theta(Z)$,*

$$\mathcal{E}_T(u_f, \varphi) = \int_Z \varphi f d\nu.$$

Moreover, there is a constant $C > 0$, which depends solely on the structural constants of Ω (or Z), such that for each $f \in L^{p'}(Z)$,

$$\mathcal{E}_p(u_f, u_f) \leq C \int_Z |f|^{p'} d\nu.$$

If $f \in L^q(Z)$ for sufficiently large q , then u_f is Hölder continuous on Z .

Proof. Given a function f as in the hypothesis of the theorem, let $u_f \in N^{1,p}(\Omega)$ be the solution to the Neumann boundary value problem for the Cheeger p -Laplacian on Ω , with Neumann boundary data f . We will denote the trace of u_f to the boundary Z also by u_f .

Note that u_f is Cheeger p -harmonic on Ω ; hence in the construction of \mathcal{E}_T , we have that $\widehat{u}_f = u_f$, and so for $\varphi \in B_{p,p}^\theta(Z)$ we have from Theorem 1.5 that

$$\mathcal{E}_T(u_f, \varphi) = \int_{\Omega} |\nabla u_f|^{p-2} \nabla u_f \cdot \nabla \widehat{\varphi} d\mu = \int_Z \varphi f d\nu.$$

Moreover, by Lemma 6.1,

$$\mathcal{E}_p(u_f, u_f) \leq C \mathcal{E}_T(u_f, u_f) = C \int_{\Omega} |\nabla u_f|^p d\mu.$$

Combining the above with Lemma A.3 yields the desired inequality $\mathcal{E}_p(u_f, u_f) \leq C \int_Z |f|^{p'} d\nu$. \square

Remark 6.3. From Theorem 1.5(c), we also know that

$$|\nabla u|^{p-2} \nabla u \cdot \nabla \eta_\epsilon d\mu \rightharpoonup -f d\nu.$$

This behavior corresponds to the behavior of functions u_f as identified in [20, page 1247] for $p = 2$ and $a = 0$ (corresponding to $(-\Delta)^{1/2}$), see also [34, page 454] for the Carnot groups setting and [33] for the setting of metric measure spaces with a doubling measure supporting a 2-Poincaré inequality. From Theorem 1.5(c), with the codimensionality between μ and ν given by the exponent Θ , we have $a = \Theta - 1$ in [20]. This justifies our definition of fractional p -Laplacian in Definition 1.10.

Proof of Theorem 1.12. We first prove Claim (i) of the theorem. We fix p, θ as in the statement of the theorem. Then, from the results of [14], we know that there is a uniform domain Ω , equipped with a doubling measure μ and supporting a 1-Poincaré inequality, such that Z is biLipschitz equivalent to $\partial\Omega$ and with ν, μ linked via the co-dimensionality condition (1.1) for $\Theta = p(1 - \theta)$, as in the discussion in Subsection 2.7. With this choice of Ω we have the construction of \mathcal{E}_T as described at the beginning of this section, and the existence of u_f now follows from Proposition 6.2.

The Hölder regularity of u_f follows from Theorem 1.6 upon noting that in gaining Hölder estimates for points $x, y \in Z$, we consider only the balls B_i centered at points in Z , and for such balls we have $\mu(B) \approx (\text{rad}(B))^{(1-\theta)p} \nu(B)$, and so the relevant lower mass bound exponent for μ here is $Q_Z + (1 - \theta)p$. Thus, if $q > \max\{1, Q_Z/\theta\}$, then the hypotheses of Theorem 1.6 is satisfied, and the Hölder regularity of u_f follows.

From Theorem 1.9 it follows that if both u_f and \widetilde{u}_f are solutions with the same data f , then $\|\widehat{\nabla u_f} - \widehat{\nabla \widetilde{u}_f}\|_{L^p(\Omega)} = 0$. Recall that for $v \in B_{p,p}^\theta(Z)$, the function \widehat{v} is the p -harmonic extension of v to Ω . As Ω supports a p -Poincaré inequality, we can conclude that $\widehat{u_f} - \widehat{\widetilde{u}_f}$ is constant on Ω and hence the trace $u_f - \widetilde{u}_f$ is constant on Z . This completes the proof of Claim (i) of the theorem.

Now we focus on proving Claim (ii) of the theorem. The $L^{p'}$ stability with respect to the boundary data f follows from Theorem 1.9. Indeed, if u_{f_1} and u_{f_2} are the solutions corresponding to the data f_1 and f_2 respectively, then their p -harmonic extensions $\widehat{u_{f_1}}$ and $\widehat{u_{f_2}}$ to Ω solve the corresponding p -Neumann problems with Neumann boundary data f_1 and f_2 . Now combining Theorem 1.9 together with the trace theorem, Theorem 2.4, yields Claim (ii).

The remainder of this proof is dedicated to Claim (iii). In order to prove the Harnack inequality, let \widehat{u} be the solution of the Dirichlet problem for the p -Laplacian in Ω , with boundary data u in Z . We first observe that as $u \geq 0$ on $Z = \partial\Omega$, by the strong maximum principle we have that either $\widehat{u} > 0$ in

Ω or else \hat{u} is identically zero in Ω (and hence in Z), see [49, Corollary 6.4]. As the zero function trivially satisfies any Harnack inequality, we only focus on the case that u is not constant in Ω . Next, we note that \hat{u} is actually p -harmonic in the open set $\Omega \cup W \subset \bar{\Omega}$. In fact, since $(-\Delta_p)^\theta u = f = 0$ in W , then for every function $\varphi \in N^{1,p}(\Omega) = N^{1,p}(\bar{\Omega})$ with support contained in $\Omega \cup W$, we have from Theorem 1.5(a) that $\int_{\Omega \cup W} |\nabla \hat{u}|^{p-2} \nabla \hat{u} \cdot \nabla \varphi \, d\mu = 0$, which tells us that \hat{u} is p -harmonic in the domain $\Omega \cup W$ (seen as a domain in the metric space $\bar{\Omega}$). The normal derivative of \hat{u} vanishes identically in W , and this allows us to use the results in [49]. Invoking [49, Corollary 7.3], one has that \hat{u} satisfies a Harnack inequality on all balls B such that $4B \subset \Omega \cup W$. In particular, for each such ball one has

$$\sup_{B \cap W} u \leq \sup_B \hat{u} \leq C \inf_B \hat{u} \leq C \inf_{B \cap W} u,$$

concluding the proof. \square

Remark 6.4. Bilinear forms such as \mathcal{E}_T and \mathcal{E}_p , for $p = 2$, correspond to a Hunt process or a jump process, see for example [9, 29, 36].

6.2. The case $p = 2$. Much of the extant literature on fractional operators deal with the linear case $p = 2$, as in [6, 7, 8, 20, 21, 22, 24, 27, 28, 29, 33, 41]. The fractional Laplacian, $(\Delta)^\theta$ on Z , as considered there was studied using spectral theory, and, in the case of [6, 7, 8, 20, 21, 22, 24, 33], was related to the behavior of the harmonic extension of the solution to a higher-dimensional domain X with the aid of the 2-Poincaré inequality on the lower-dimensional space Z . It was also shown there that the spectral construction of the fractional Laplacian operators agree with the infinitesimal generator (see for example [36]) of the non-local bilinear form \mathcal{E}_2 . In the current paper we give an intrinsic construction of a fractional Laplacian on Z by realizing it as the boundary of a John domain, and so in the case $p = 2$ we have two approaches to constructing the fractional Laplacian operators. In the case that Z itself does not support a 2-Poincaré inequality, the spectral construction as described in [20, 33] is not possible; however, the infinitesimal generator \mathcal{A} of \mathcal{E}_2 and the operator \mathcal{A}_T constructed in the current paper both exist. For completeness of discussion, we now consider the case $p = 2$ and discuss the construction of the infinitesimal generator \mathcal{A} associated with the bilinear form \mathcal{E}_2 , as considered in the above-mentioned literature. To do so, we need to consider \mathcal{E}_2 as a norm; however, it is not a norm on $B_{2,2}^\theta$, and hence we need to extend the bilinear form to the homogeneous Besov classes; see Subsection 2.4 for the relevant notions.

In the proposition below, we fix $p = 2$, and note by \mathcal{A} the infinitesimal generator associated with the symmetric bilinear form \mathcal{E}_2 on $HB_{2,2}^\theta(\partial\Omega)$. For functions $u \in B_{2,2}^\theta(\partial\Omega)$, we say that u is in the domain of \mathcal{A} if there is a function $f \in L^2(\partial\Omega)$ such that for each $v \in B_{2,2}^\theta(\partial\Omega)$ we have $\mathcal{E}_2(u, v) = -\int_{\partial\Omega} v f \, d\nu$. In this case we denote $\mathcal{A}u = f$.

Since $\partial\Omega$ is a bounded set and hence $\nu(\partial\Omega)$ is finite, it follows that constant functions are in $B_{2,2}^\theta(\partial\Omega)$. Hence, by using the choice of $v = 1$ in the defining identity of $\mathcal{A}u$ in the above paragraph, we must have $\int_{\partial\Omega} f \, d\nu = 0$ if $\mathcal{A}u = f$.

Note that if $w, u \in B_{2,2}^\theta(\partial\Omega)$ such that $u - w$ is constant on $\partial\Omega$, then u is in the domain of \mathcal{A} if and only if w is; moreover, $\mathcal{A}u = \mathcal{A}w$.

Proposition 6.5. *Let $f \in L^2(\partial\Omega)$ such that $\int_{\partial\Omega} f \, d\nu = 0$. Then there exists $w_f \in B_{2,2}^\theta(\partial\Omega)$ such that $\mathcal{A}w_f = f$.*

Proof. From [55] we know that u_f that satisfies any of the conditions set forth in Theorem 1.5 exists. It follows from the above discussion that $\mathcal{A}_T T u_f = f$ and $\widehat{T u_f} = u_f$. From the discussion above we have

$$\left| \int_{\partial\Omega} \varphi f \, d\nu \right| \leq C_0 \|\varphi\|_{\theta,2} = C_0 \mathcal{E}_2(\varphi, \varphi)^{1/2},$$

with C_0 depending on f . We know that $\|\cdot\|_{\theta,2}$ is a norm on the homogeneous space $HB_{2,2}^\theta(\partial\Omega) = B_{2,2}^\theta(\partial\Omega)/\sim$, and by above, $\varphi \mapsto \int_{\partial\Omega} \varphi f \, d\nu$ is a bounded linear map on $HB_{2,2}^\theta(\partial\Omega)$ because $\int_{\partial\Omega} \varphi f \, d\nu = \int_{\partial\Omega} (\varphi - c) f \, d\nu$ for each real number c . By Lemma 2.2 we have that $HB_{2,2}^\theta(\partial\Omega)$ is a reflexive Banach space, and so it follows from the Riesz representation theorem that there is some function (up to a constant) $w_f \in HB_{2,2}^\theta(\partial\Omega)$ such that for each $\varphi \in HB_{2,2}^\theta(\partial\Omega)$,

$$\int_{\partial\Omega} \varphi f \, d\nu = \mathcal{E}_2(w_f, \varphi). \quad \square$$

The symmetric non-local bilinear form \mathcal{E}_2 , as described above, is part of a class of symmetric non-local bilinear forms studied in [29]. The version of Poincaré inequality considered in [29, Definition 1.19] is tautological for the form \mathcal{E}_2 considered above with $\phi(r) = r^2$, and so by [29, Theorem 1.20] we have that an \mathcal{E}_2 -harmonic function is necessarily Hölder continuous on its domain of harmonicity. Here, from the discussion in [29, Section 2], a function u is \mathcal{E}_2 -harmonic in an open set $U \subset X$ if $u \in B_{2,2}^\theta(X)$ and for all $\varphi \in B_{2,2}^\theta(X)$ with compact support in U we have $\mathcal{E}_2(u, \varphi) = 0$. In particular, our construction w_f is \mathcal{E}_2 -harmonic in an open set $U \subset X$ if $f = 0$ on U ; it then follows from [29, Theorem 1.20] that w_f is Hölder continuous on U . The results in [29] do not extend to the case where f is not zero on U , and so for more general f , we do not know whether w_f is Hölder continuous, but from Theorem 1.6 above we know that u_f is indeed Hölder continuous when $f \in L^q(X) = L^q(\partial\Omega)$ for q sufficiently large.

7. RECONCILING CONSTRUCTION OF FRACTIONAL LAPLACIAN WITH [33]

In Section 6 we gave a possible construction of a fractional p -Laplacian operator $(-\Delta_2)^\theta$ on a doubling metric measure space $\partial\Omega$. In the special case $p = 2$, and with the additional hypothesis that $\partial\Omega$ also supports a 2-Poincaré inequality, an alternative construction based on spectral theory can be found in [33], which corresponds to the operator \mathcal{A} described in Subsection 6.2. The aim of this section is to reconcile these two different approaches, and show that the construction given in [33] gives rise to the same fractional operator we define in Section 6 above.

In [33] the object of study was a compact doubling metric measure space (Z, d_Z, μ) that supports a 2-Poincaré inequality, and this metric space is

naturally seen as the boundary of the unbounded domain $Z \times (0, \infty)$, where $X = Z \times [0, \infty)$ is equipped with the ℓ_2 -product metric d . Whereas, in our paper we consider the boundary of a *bounded* domain. Therefore, to show that the two approaches are not contradictory, we show that we can modify X so that it becomes a bounded doubling metric measure space supporting a 2-Poincaré inequality and that functions that are 2-harmonic in $Z \times (0, \infty)$ or X are also 2-harmonic in this modified space, and that Z is isometric to the boundary of this modified space. We consider the metric d_∞ on X given by $d_\infty((x_1, y_1), (x_2, y_2)) = \max\{d(x_1, x_2), |y_1 - y_2|\}$.

Fix $\beta > 1$, and set ρ and ω to be the following continuous functions on X :

$$\rho(x, y) := \min\{1, y^{-\beta}\}, \quad \omega(x, y) = \min\{1, y^{-2\beta}\}.$$

The metric d_ρ on X is given by

$$d_\rho((x_1, y_1), (x_2, y_2)) = \inf_{\gamma} \int_{\gamma} \rho \, ds,$$

where the infimum is over all rectifiable curves in X with end points $(x_1, y_1), (x_2, y_2)$. Since ρ is a positive continuous function and X is complete, it follows that the topology generated by d_ρ is the same as the topology generated by the original metric on X . Moreover, for $(x_0, y_0) \in X$,

$$\lim_{(x, y) \rightarrow (x_0, y_0)} \frac{d_\rho((x, y), (x_0, y_0))}{d((x, y), (x_0, y_0))} = \rho(x_0, y_0),$$

and hence (X, d_ρ) is a geodesic space. The metric d_ρ is motivated by the procedure of sphericalization as constructed in [4], see also [30, 31, 51, 52]. However, the measure also needs to be modified, not as in [30, 31, 51, 52], but in the manner of [13]. The modified measure μ_ω is given by

$$\mu_\omega(A) = \int_A \omega \, d\mu_X$$

where μ_X is the product measure on X given by $d\mu_X(x, y) = y^a \, d\mu(x) \, dy$, with $a = 1 - 2\theta$ as in [33]. From the construction, it is clear that the completion \widehat{X} of X with respect to the metric d_ρ is compact.

We denote the arc-length measure on a curve γ in (X, d_∞) by ds ; then under the deformed metric d_ρ , the arc-length measure ds_ρ is given by $ds_\rho = \rho \, ds$. It follows that if g is an upper gradient of a function u on X or $Z \times (0, \infty)$, then $\rho^{-1}g$ is an upper gradient of u on (X, d_ρ) or $(Z \times (0, \infty), d_\rho)$. This is because

$$\int_{\gamma} g_0 \, ds_\rho = \int_{\gamma} g_0 \, \rho \, ds$$

whenever g_0 is a Borel function on X or $Z \times (0, \infty)$. Observe that

$$\int_X \left(\frac{g}{\rho} \right)^2 \, d\mu_\omega = \int_X g^2 \, d\mu_X,$$

and so a family of curves Γ is of zero 2-modulus in (X, d_∞) if and only if it is of zero 2-modulus in (X, d_ρ) . Moreover, if g_u is a minimal 2-weak upper

gradient of u in (X, d_∞) and $g_{u,\rho}$ is a minimal 2-weak upper gradient of u in (X, d_ρ) , then $g_{u,\rho} = \rho^{-1} g_u$, with

$$\int_X g_{u,\rho}^2 d\mu_\omega = \int_X g_u^2 d\mu_X.$$

This means that the upper gradient energy is the same with respect to both metrics.

Note that (X, d_ρ) is a bounded metric space, for

$$\frac{1}{2} \text{diam}_\rho(X) \leq \int_0^\infty \rho(x_0, y) dy = 1 + \int_1^\infty y^{-\beta} dy = 1 + \frac{1}{\beta-1} = \frac{\beta}{\beta-1} < \infty.$$

On the other hand, (X, d_ρ) is not complete; we complete it by including the “point at infinity”. So we set $\widehat{X} = X \cup \{\infty\}$, with $d_\rho((x, y), \infty) := \int_y^\infty \rho(x, y) dy$, that is,

$$(7.1) \quad d_\rho((x, y), \infty) = \begin{cases} \frac{1}{(\beta-1)y^{\beta-1}} & \text{if } y \geq 1, \\ \frac{\beta}{\beta-1} - y & \text{if } 0 \leq y \leq 1. \end{cases}$$

We wish to consider as the domain Ω the set $\widehat{X} \setminus (Z \times \{0\})$, and note that the restriction of d_ρ to $\partial\Omega = Z \times \{0\}$ is isometric to d_Z . Observe that if $x_0 \in Z$ and $(x_1, y_1) \in \widehat{X}$ such that $d_\rho((x_0, 0), (x_1, y_1)) \leq 1$, then the ρ -geodesic connecting $(x_0, 0)$ to (x_1, y_1) must lie in the region $Z \times [0, 1]$, and so $d_\rho((x_0, 0), (x_1, y_1)) = d((x_0, 0), (x_1, y_1)) = \max\{d_X(x_1, x_0), |y_1|\}$. Hence for each $x_0 \in Z$ and $r \leq 1$, we have

$$\mu_\omega(B_\rho((x_0, 0), r)) = (1+a)^{-1} r^{1+a} \mu_Z(B(x_0, r)).$$

Thus, for radii $r < 1$, balls centered at $(x_0, 0)$ are of co-dimension $1+a$ measure with respect to μ_X . Note that then we have $\Theta = 1+a$ and $\theta = \frac{1-a}{2}$, as required in the current note.

We now need to know what the effect of the inclusion of the point at infinity has on the class of 2-weak upper gradients.

Lemma 7.2. *Let Γ be the collection of all non-constant rectifiable curves in (\widehat{X}, d_ρ) that passes through ∞ . Then $\text{Mod}_2(\Gamma) = 0$.*

Proof. It suffices to show that for each fixed $L \geq 1$, the collection Γ_L of all curves with one end point in $Z \times \{L\}$ and the other at ∞ satisfies $\text{Mod}_2(\Gamma_L) = 0$, for $\Gamma = \bigcup_{n \in \mathbb{N}} \Gamma_n$. For each $H > L$, we set $\rho_H = (\beta-1)H^{\beta-1} \chi_{Z \times [H, \infty)}$. A direct calculation shows that ρ_H is admissible for computing $\text{Mod}_2(\Gamma_L)$ as every curve in Γ_L has a subcurve in $Z \times [H, \infty)$ with one end point at $Z \times \{H\}$ and the other at ∞ . Note that as $a < 1$, we have $\frac{a+1}{2} < 1 < \beta$. Therefore

$$\text{Mod}_2(\Gamma_L) \leq \int_Z \int_H^\infty \rho_H^2 d\mu_\omega = \mu(Z) (\beta-1)^2 H^{2\beta-2} \int_H^\infty y^{a-2\beta} dy = \frac{(\beta-1)^2}{2\beta-a-1} \frac{1}{H^{1-a}}.$$

Letting $H \rightarrow \infty$ and noting that $a < 1$ yields the desired conclusion. \square

From the above lemma, it is clear that a 2-weak upper gradient of a function u on $Z \times (0, \infty)$ (or X) extends as a 2-weak upper gradient of u on Ω (or \widehat{X}). Moreover, the total Newton-Sobolev 2-capacity of $\{\infty\}$ is zero.

7.1. Doubling property of μ_ω . We want to show the existence of a constant $C \geq 1$ such that whenever $0 < r < 1/4$, $\mu_\omega(B_\rho((x_0, y_0), 2r)) \leq C\mu_\omega(B_\rho((x_0, y_0), r))$. We first need the following lemma.

Lemma 7.3. *There exists $C_\omega > 0$ such that whenever B is a ball (with respect to the metric d_ρ) in \widehat{X} with radius $R > 0$ and center (x_0, y_0) such that $\infty \notin 2B$, then for each $(x, y) \in B$ we have*

$$\frac{\omega(x_0, y_0)}{C_\omega} \leq \omega(x, y) \leq C_\omega \omega(x_0, y_0), \quad \frac{\rho(x_0, y_0)}{C_\omega} \leq \rho(x, y) \leq C_\omega \rho(x_0, y_0).$$

Moreover, for each $(x, y) \in B$ we have

$$\frac{\rho(x_0, y_0)}{C_\omega} d((x_0, y_0), (x, y)) \leq d_\rho((x_0, y_0), (x, y)) \leq C_\omega \rho(x_0, y_0) d((x_0, y_0), (x, y)),$$

and if in addition we have $y_0 \geq 1$ and $y > 1$, then

$$\begin{aligned} \frac{1}{C_\omega(\beta-1)} \left| \frac{1}{y_0^{\beta-1}} - \frac{1}{y^{\beta-1}} \right| + \frac{1}{C_\omega y_0^\beta} d_Z(x, x_0) &\leq d_\rho((x, y), (x_0, y_0)) \\ (7.4) \qquad \qquad \qquad &\leq \frac{C_\omega}{\beta-1} \left| \frac{1}{y_0^{\beta-1}} - \frac{1}{y^{\beta-1}} \right| + \frac{C_\omega}{y_0^\beta} d_Z(x, x_0) \end{aligned}$$

Proof. If $y_0 \leq 1$, then $\omega(x_0, y_0) = 1$. If for all $(x, y) \in B$ we have that $y \leq 1$, then $\omega(x, y) = 1 = \omega(x_0, y_0)$. Hence, without loss of generality, we may assume that there is some $(x, y) \in B$ with $y > 1$. As $\infty \notin 2B$, we have

$$2R \leq \int_{y_0}^{\infty} \rho dy = 1 - y_0 + \int_1^{\infty} y^{-\beta} dy = 1 - y_0 + \frac{1}{\beta-1}.$$

As $(x, y) \in B_R$,

$$R > \int_{y_0}^y \rho dy = 1 - y_0 + \frac{1}{\beta-1} (1 - y^{1-\beta}).$$

By combining the previous two estimates we get

$$\frac{1-y_0}{2} + \frac{1}{2(\beta-1)} (1 - 2y^{1-\beta}) < 0.$$

This can only be satisfied if the second term is negative. Thus we have to have

$$y < 2^{1/(\beta-1)}$$

and so $\omega(x, y) \approx 1 \approx \rho(x_0, y_0)$, satisfying the first claim of the lemma.

Now we consider the case $y_0 > 1$. As $\infty \notin 2B$, we obtain

$$2R < \int_{y_0}^{\infty} \rho dy \leq \int_{y_0}^{\infty} y^{-\beta} dy = \frac{1}{\beta-1} y_0^{1-\beta}.$$

If $y \geq 1$, then $(x, y) \in B$ implies that

$$\left| \int_{y_0}^y y^{-\beta} dy \right| = \frac{1}{\beta-1} \left| y_0^{1-\beta} - y^{1-\beta} \right| < R.$$

A combination of the above two estimates gives us that

$$\frac{1}{2} y_0^{1-\beta} < y^{1-\beta} < \frac{3}{2} y_0^{1-\beta}.$$

It follows that $y \approx y_0$, and so again $\omega(x, y) \approx \omega(x_0, y_0)$ and $\rho(x, y) \approx \rho(x_0, y_0)$. If $y \leq 1$, then as $(x_0, y_0) \in B((x, y), R)$, it follows by the discussion in the first paragraph above that we have $\omega(x, y) = 1 \approx \omega(x_0, y_0)$. This completes the proof of the first claim.

Now we let $(x, y) \in B$, and note that any d_ρ -geodesic connecting (x_0, y_0) to (x, y) lies in B ; hence by the above, we have that

$$\frac{1}{C_\omega} \rho(x_0, y_0) d((x, y), (x_0, y_0)) \leq d_\rho((x, y), (x_0, y_0)).$$

On the other hand, let γ_Z be a geodesic curve in Z with end points x, x_0 and parametrized to be from the interval $[0, |y - y_0|]$ and with constant speed, and $\gamma_{\mathbb{R}}$ be the vertical line segment $\gamma_{\mathbb{R}}(t) = t + \min\{y, y_0\}$, for $0 \leq t \leq |y_0 - y|$. Let γ be the curve given by $\gamma(t) = (\gamma_Z(t), \gamma_{\mathbb{R}}(t))$. Note by the discussion above, we have that either $y_0 \leq 1$, in which case $0 \leq y \leq 2^{1/(\beta-1)}$ and so for each t we have that $\omega(\gamma(t)) \approx \omega(x_0, y_0)$, or $y_0 > 1$, in which case $y \approx y_0$ and so again $\rho(\gamma(t)) \approx \rho(x_0, y_0)$. Hence

$$d_\rho((x, y), (x_0, y_0)) \leq \ell_\rho(\gamma) \leq C \rho(x_0, y_0) d((x, y), (x_0, y_0)).$$

Thus the second claim of the lemma is also verified.

To verify the last claim, we now suppose that $y \geq 1$ and $y_0 \geq 1$. Note that \widehat{X} is a geodesic space, and hence every geodesic (with respect to the metric d_ρ) connecting the center (x_0, y_0) of B to $(x, y) \in B$ lies entirely in B . Let $\gamma : [0, L] \rightarrow X$ be such a geodesic, arc-length parametrized with respect to the original metric d on X . Then with $\gamma(t) = (\gamma_Z(t), \gamma_{\mathbb{R}}(t))$, and as both $y_0, y \geq 1$,

$$\begin{aligned} d_\rho((x_0, y_0), (x, y)) &= \int_0^L \gamma_{\mathbb{R}}(t)^{-\beta} \max\{|\gamma'_Z|(t), |\gamma'_{\mathbb{R}}|(t)\} dt \\ &\geq \int_0^L \gamma_{\mathbb{R}}(t)^{-\beta} |\gamma'_{\mathbb{R}}|(t) dt \geq \frac{1}{\beta - 1} \left| \frac{1}{y_0^{\beta-1}} - \frac{1}{y^{\beta-1}} \right|. \end{aligned}$$

Moreover, for each $t \in [0, L]$ we have that $\gamma_{\mathbb{R}}(t) \approx y_0$ by the argument above related to the first claim of the lemma. It follows that

$$d_\rho((x_0, y_0), (x, y)) \geq \frac{1}{C_\omega y_0^\beta} \int_0^L |\gamma'_Z|(t) dt \geq \frac{1}{C_\omega y_0^\beta} d_Z(x, x_0).$$

Combining this with the above inequality yields the first of the two inequalities in (7.4). The right-hand side of (7.4) is obtained by considering the ρ -length of the curve γ_0 obtained by concatenating the curve $\gamma_{0, \mathbb{R}} : [0, |y - y_0|] \rightarrow X$ given by $\gamma_{0, \mathbb{R}}(t) = (x_0, y_0 + \frac{y - y_0}{|y - y_0|} t)$ with a geodesic $\gamma_{0, Z}(t) = (\gamma_{1, Z}(t), y)$ where $\gamma_{1, Z}$ is any geodesic in Z with end points x_0, x . \square

Lemma 7.5. *The measure μ_ω is doubling on \widehat{X} .*

In the proof below, each occurrence of $C_{\beta, a}$ denotes a possibly different constant, whose choice depends solely on the parameters β and a .

Proof. Let B_ρ be a ball of radius $r > 0$ in \widehat{X} . Since \widehat{X} is compact, it suffices to prove the doubling property for balls of radius r small enough, namely,

$$(7.6) \quad 0 < r \leq \min \left\{ 1/4, \frac{1}{(8 \operatorname{diam}(Z))^{\beta-1} (4(\beta-1))^\beta}, \frac{1}{8(\beta-1)} \right\}.$$

We consider three cases.

Case 1: $\infty \in \frac{1}{2}B_\rho$. In this case, note that $B_\rho(\infty, r/2) \subset B_\rho$ and $2B_\rho \subset B_\rho(\infty, 3r)$. Note that when $0 < R < 1/(\beta-1)$, $B_\rho(\infty, R) = Z \times [H_R, \infty)$ with

$$(7.7) \quad H_R = \left(\frac{1}{(\beta-1)R} \right)^{1/(\beta-1)}.$$

It follows that

$$(7.8) \quad \begin{aligned} \mu_\omega(B_\rho(\infty, R)) &= \mu(Z) \int_{H_R}^\infty y^{a-2\beta} dy = \frac{\mu(Z)}{2\beta-1-a} H_R^{1+a-2\beta} \\ &= \mu(Z) \frac{(\beta-1)^{\frac{2\beta-1-a}{\beta-1}}}{2\beta-1-a} R^{\frac{2\beta-1-a}{\beta-1}}. \end{aligned}$$

Therefore,

$$\begin{aligned} \mu_\omega(2B_\rho) &\leq \mu_\omega(B_\rho(\infty, 3r)) = C_{\beta,a} \mu(Z) r^{\frac{2\beta-a-1}{\beta-1}} = C_{\beta,a} \mu_\omega(B_\rho(\infty, r/2)) \\ &\leq C_{\beta,a} \mu_\omega(B_\rho), \end{aligned}$$

that is, the doubling property holds for balls falling within this case, with doubling constant that depends only on β and a .

Case 2: $\infty \in 4B_\rho \setminus \frac{1}{2}B_\rho$. Since we assume that r is small enough (see (7.6)), we have that necessarily $y_0 > 1$. Since $r/4 \leq d_\rho((x_0, y_0), \infty) < 4r$, by (7.1) it follows that

$$(7.9) \quad \frac{r}{2} \leq \frac{1}{(\beta-1)y_0^{\beta-1}} < 4r.$$

We fix a positive real number Δ (independently of r) such that

$$\Delta < \min \left\{ 1, \left(\frac{16}{15} \right)^{\beta/(\beta-1)} - 1, 1 - \left(\frac{16}{17} \right)^{1/(\beta-1)} \right\}.$$

Then by (7.6) and (7.9), we see that $X \times [y_0(1-\Delta), y_0(1+\Delta)] \subset B_\rho$. A direct computation shows then that

$$\begin{aligned} \mu_\omega(B_\rho) &\geq \mu_\omega(X \times [y_0(1-\Delta), y_0(1+\delta)]) \\ &= \frac{\mu(Z)}{(2\beta-1-a)y_0^{2\beta-1-a}} \left[(1-\Delta)^{1+a-2\beta} - (1+\Delta)^{1+a-2\beta} \right] \\ &\approx r^{(2\beta-1-a)/(\beta-1)}. \end{aligned}$$

The comparison constant in the above depends only on the choice of Δ and on $\mu(Z)$, β . As $\infty \in 4B_\rho$, we see that $4B_\rho \subset B_\rho(\infty, 8r)$. From the computations as in Case 1 above tells us that

$$\mu_\omega(2B_\rho) \leq \mu_\omega(B_\rho(\infty, 8r)) = \mu(Z) C_{\beta,a} r^{(2\beta-1-a)/(\beta-1)}.$$

Combining this estimate with the antecedent estimate gives $\mu_\omega(2B_\rho) \lesssim \mu_\omega(B_\rho)$.

Case 3: $\infty \notin 4B_\rho$. In this case, by the first part of Lemma 7.3 we know that $\mu_\omega(2B_\rho) \approx \omega(x_0, y_0)\mu(2B_\rho)$ and $\mu_\omega(B_\rho) \approx \omega(x_0, y_0)\mu(B_\rho)$. Therefore it suffices to show that $\mu(2B_\rho) \leq C\mu(B_\rho)$ for some constant C that does not depend on B_ρ . Now by the last part of the lemma, we know that for $(x, y) \in 2B_\rho$ we have that

$$d_\rho((x_0, y_0), (x, y)) \approx y_0^{\beta-1} d((x, y), (x_0, y_0)).$$

Thus $2B_\rho \subset B((x_0, y_0), 2Cry_0^\beta)$ and $B((x_0, y_0), ry_0^\beta/C) \subset B_\rho$. Now the doubling property of μ yields the desired inequality, with the doubling constant depending only on the doubling constant of μ and the constant C used here. \square

7.2. Modification of the norm on Cheeger differential structure.

The paper [33] considers D to be a (fixed choice of) Cheeger differential structure on Z . The existence of such a structure is guaranteed from the results in [25]. The corresponding differential structure ∇ on X is obtained as a tensorization of the structure D with the Euclidean differential structure on $[0, \infty)$. When the metric on X is modified as considered in this section, the norm associated with the differential structure on X is also changed. Note that the norm, as considered in Cheeger in [25], has the property that

$$|\nabla u(x, y)|_{(x, y)} \approx g_u(x, y)$$

for $\mu \times \mathcal{L}^1$ -a.e. $(x, y) \in X$; here, g_u is a 2-weak upper gradient of u with respect to the original (unmodified) metric. Since $\rho^{-1}g_u$ is a 2-weak upper gradient of u with respect to the metric d_ρ and the measure μ_ω , it follows that the norm on the inner product structure ∇ should change accordingly in order to preserve the above comparison. Hence we choose

$$|\nabla u(x, y)|_{\rho, (x, y)} = \rho(x, y)^{-1} |\nabla u(x, y)|_{(x, y)}.$$

With this modification, we have that for each Borel $A \subset \widehat{X}$,

$$\int_A \langle \nabla u(x, y), \nabla v(x, y) \rangle_{\rho, (x, y)} \mu_\omega(x, y) = \int_A \langle \nabla u(x, y), \nabla v(x, y) \rangle_{(x, y)} d(\mu \times \mathcal{L}^1)(x, y).$$

7.3. 2-Poincaré inequality with respect to d_ρ and μ_ω . It was shown in [33, Proof of Lemma 3.1] that as Z supports a 2-Poincaré inequality, the product space $X = Z \times (0, \infty)$ also supports a 2-Poincaré inequality. In this subsection we will show that the modified space \widehat{X} also supports a 2-Poincaré inequality.

Proposition 7.10. *The space $(\widehat{X}, d_\rho, \mu_\omega)$ supports a $(1, 2)$ -Poincaré inequality.*

Proof. As the metric d_ρ is geodesic, a weak Poincaré inequality implies the strong Poincaré inequality. Thus it is enough to prove that for every $u \in$

$D^{1,2}(\widehat{X})$ and every ball $B \subset \widehat{X}$, there exists a constant $c_{u,B}$ such that

$$\int_{B_\rho(x,r)} |u - c_{u,B}| d\mu_\omega \leq C_P r \left(\int_{B_\rho(x,\lambda r)} g_u^p d\mu_\omega \right)^{1/2},$$

with some constants C_P and λ that only depend on the data of the space. We consider separately two cases: The case, where the point ∞ is far away from B , and the case where it is close to B .

Case 1: $\infty \notin B_\rho((x_0, y_0), 2C_\omega^2 r_0)$, where C_ω is as in Lemma 7.3. Then, with $\tilde{B} = B((x_0, y_0), C_\omega \rho(x_0, y_0)^{-1} r_0)$, by Lemma 7.3 we have

$$B_\rho = B_\rho((x_0, y_0), r_0) \subset \tilde{B} \subset C_\omega^2 B_\rho = B_\rho((x_0, y_0), C_\omega^2 r_0).$$

Now, setting $c_{u,B} = \mu_Z(\tilde{B})^{-1} \int_{\tilde{B}} u d\mu_X$, we can estimate using Lemma 7.3 and the Poincaré inequality of (X, d_X, μ_X) that

$$\begin{aligned} \int_{B_\rho} |u - c_{u,B}| d\mu_\omega &\leq C_\omega \omega(x_0, y_0) \int_{\tilde{B}} |u - c_{u,B}| d\mu_X \\ &\leq C_\omega^2 \omega(x_0, y_0) \rho(x_0, y_0)^{-1} r_0 \mu_X(\tilde{B})^{1/2} \left(\int_{\tilde{B}} g_u^2 d\mu_X \right)^{1/2} \\ &= C_\omega^2 \omega(x_0, y_0) \rho(x_0, y_0)^{-1} r_0 \mu_X(\tilde{B})^{1/2} \left(\int_{\tilde{B}} g_{u,\rho}^2 d\mu_\omega \right)^{1/2} \\ &\leq C_\omega^2 r_0 \mu_\omega(C_\omega^2 B_\rho)^{1/2} \left(\int_{C_\omega^2 B_\rho} g_{u,\rho}^2 d\mu_\omega \right)^{1/2} \end{aligned}$$

Here the first inequality follows from the inclusion $B_\rho \subset \tilde{B}$ and from having $\omega \approx \omega(x_0, y_0)$ in \tilde{B} . The second inequality follows from the $(1, 2)$ -Poincaré inequality for (X, d, μ_X) . In the last estimate, we used the facts that for $(x, y) \in B_\rho$, $\omega(x, y)^{1/2} \approx \omega(x_0, y_0)^{1/2} = \omega(x_0, y_0) \rho(x_0, y_0)^{-1}$ in \tilde{B} and that $\tilde{B} \subset C_\omega^2 B_\rho$. Thus the Poincaré inequality is satisfied with $\lambda = C_\omega^2$ and C_P that depends on C_ω as well as the doubling constant of \widehat{X} and the constants associated with the Poincaré inequality for (X, d_X, μ_X) .

Case 2: $\infty \in B_\rho((x_0, y_0), 2C_\omega^2 r_0)$.

Now we have $B_\rho((x_0, y_0), r_0) \subset B_\rho(\infty, (2C_\omega^2 + 1)r_0) = Z \times (H_{(2C_\omega^2 + 1)r_0}, \infty]$, where $H_R = [(\beta - 1)R]^{-1/(\beta - 1)}$ as in (7.7). Therefore as we already know that μ_ω is doubling, it is enough to prove that the balls centered at ∞ satisfy the Poincaré inequality. Thus let us consider the ball $B_\rho = B_\rho(\infty, R)$ with $R < 1/(2\beta - 2)$.

For $k = 1, 2, \dots$, let $A_k = B_\rho(\infty, 2^{1-k}R) \setminus B_\rho(\infty, 2^{-k}R)$. Then from equation (7.8) above, we have $\mu_\omega(A_{k+1})/\mu_\omega(A_k) = \gamma = (\frac{1}{2})^{1+(\beta-a)/(\beta-1)} < 1$. We may assume without loss of generality that $u_{A_1} = 0$ (with respect to the measure μ_ω). Then our goal is to prove that

$$\int_{B_\rho} |u| d\mu_\omega \leq CR \left(\int_{B_\rho} g_{u,\rho}^2 d\mu_\omega \right)^{1/2} \mu_\omega(B_\rho)^{1/2}.$$

Note from the discussion around (7.7) that $A_1 = Z \times [H_R, H_{R/2}]$ with $H_R > 2$, and so for each $(x, y) \in A_1$ we have that

$$2^{-2\beta/(\beta-1)} H_R^{-2\beta} = H_{R/2}^{-2\beta} \leq \omega(x, y) \leq H_R^{-2\beta}.$$

Note that $H_{R/2} - H_R = (2^{1/(\beta-1)} - 1)H_R \geq \text{diam}(Z)$. Since (Z, d_Z, μ_Z) supports a 2-Poincaré inequality and so does the Euclidean interval $[H_R, H_{R/2}]$ (equipped with the weighted 1-dimensional Lebesgue measure $y^a dy$), we see that

(7.11)

$$\int_{A_1} |u| d\mu_X = \int_{A_1} |u - u_{A_1}| d\mu_X \leq C H_R \mu_X(A_1)^{1/2} \left(\int_{A_1} g_u^2 d\mu_X \right)^{1/2}.$$

Here we used the fact that $H_R > 2 > \text{diam}(X)$, and so the d_X -diameter of A_1 is comparable to H_R . Now given the above comparison $\omega((x, y)) \approx H_R^{-2\beta}$, we obtain

$$\begin{aligned} \int_{A_1} |u| d\mu_\omega &\leq C H_R^{1-2\beta} \left(\int_{A_1} g_{u,\rho}^2 d\mu_\omega \right)^{1/2} H_R^\beta \mu_\omega(A_1)^{1/2} \\ &= C(\beta - 1) R \left(\int_{A_1} g_{u,\rho}^2 d\mu_\omega \right)^{1/2} \mu_\omega(A_1)^{1/2}. \end{aligned}$$

Now it is enough to prove that $\int_{B_\rho \setminus A_1} |u| d\mu_\omega \leq C R \int_{B_\rho} g_{u,\rho} d\mu_\omega$, from which we can recover (1, 2)-Poincaré inequality by using Hölder's inequality.

As ω and ρ are approximately constant in each $A_k \cup A_{k+1}$ and the y -directional width with respect to the metric d_ρ of $A_k \cup A_{k+1}$ is smaller than $2^{2-k}R$, the use of the 1-dimensional Poincaré inequality on $\{x\} \times (0, \infty)$ for fixed $x \in X$ gives us

$$\begin{aligned} |u_{A_{k+1}} - u_{A_k}| &= \left| \int_{A_{k+1}} u d\mu_\omega - \int_{A_k} u d\mu_\omega \right| \approx \left| \int_{A_{k+1}} u d\mu_Z - \int_{A_k} u d\mu_X \right| \\ &\leq C H_{R_k} \int_{A_k \cup A_{k+1}} g_u d\mu_X \\ &\approx C H_{R_k} \int_{A_k \cup A_{k+1}} g_u d\mu_\omega \\ &\approx C H_{R_k}^{1-\beta} \int_{A_k \cup A_{k+1}} g_{u,\rho} d\mu_\omega \\ &\approx C 2^{-k} R \int_{A_k \cup A_{k+1}} g_{u,\rho} d\mu_\omega. \end{aligned}$$

Here u_{A_k} , $k \in \mathbb{N}$, are computed with respect to the measure μ_ω . Now we can apply a telescoping argument and (7.11) to estimate that

$$\begin{aligned}
\int_{B_\rho \setminus A_1} |u| d\mu_\omega &= \sum_{k=2}^{\infty} \int_{A_k} |u| d\mu_\omega \\
&\leq \sum_{k=2}^{\infty} \mu_\omega(A_k) \left(|u_{A_1}| + \sum_{m=1}^{k-1} |u_{A_m} - u_{A_{m+1}}| \right) \\
&\leq C \sum_{k=1}^{\infty} \sum_{m=1}^{k-1} \mu_\omega(A_k) 2^{-m} R \int_{A_m \cup A_{m+1}} g_{u,\rho} d\mu_\omega + C R \left(\int_{A_1} g_{u,\rho}^2 d\mu_\omega \right)^{1/2} \sum_{k=2}^{\infty} \mu_\omega(A_k) \\
&\leq C \sum_{k=1}^{\infty} \sum_{m=1}^{k-1} \mu_\omega(A_k) 2^{-m} R \int_{A_m \cup A_{m+1}} g_{u,\rho} d\mu_\omega + C R \left(\int_{A_1} g_{u,\rho}^2 d\mu_\omega \right)^{1/2} \mu_\omega(B_\rho \setminus A_1) \\
&\leq C \sum_{m=1}^{\infty} \sum_{k=m+1}^{\infty} 2^{-m} R \int_{A_m \cup A_{m+1}} g_{u,\rho} d\mu_\omega \frac{\mu_\omega(A_k)}{\mu_\omega(A_m)} + C R \left(\int_{A_1} g_{u,\rho}^2 d\mu_\omega \right)^{1/2} \mu_\omega(B_\rho) \\
&\leq C \sum_{m=1}^{\infty} \sum_{k=m+1}^{\infty} 2^{-m} R \int_{A_m \cup A_{m+1}} g_{u,\rho} d\mu_\omega + C R \left(\int_{A_1} g_{u,\rho}^2 d\mu_\omega \right)^{1/2} \mu_\omega(B_\rho) \\
&\leq C R \int_{B_\rho} g_{u,\rho} d\mu_\omega + C R \left(\int_{A_1} g_{u,\rho}^2 d\mu_\omega \right)^{1/2} \mu_\omega(B_\rho).
\end{aligned}$$

Hence by Hölder's inequality and (7.11),

$$\int_{B_\rho} |u| d\mu_\omega \leq C R \left(\int_{B_\rho} g_{u,\rho}^2 d\mu_\omega \right)^{1/2} \mu_\omega(B_\rho),$$

which yields the 2-Poincaré inequality. \square

7.4. John domain property. In this section we will show that \widehat{X} is a John domain. In fact, it is also a uniform domain (see for example [40]), but we do not need this stronger geometric condition in our paper.

Proposition 7.12. *The space $(\widehat{X}, d_\rho, \mu_\omega)$ is a John domain with John center ∞ and boundary $Z \times \{0\}$.*

Proof. Since for $0 < y \leq 1$ and $x \in Z$ we have that $\text{dist}_{d_\rho}((x, y), Z \times \{0\}) = \text{dist}_d((x, y), Z \times \{0\}) = y$, it follows that the boundary of \widehat{X} contains $Z \times \{0\}$. Moreover, if $x_1, x_2 \in Z$ such that $d_X(x_1, x_2) \leq 1/2$, then $d_\rho((x_1, 0), (x_2, 0)) = d((x_1, 0), (x_2, 0)) = d_X(x_1, x_2)$. We now show that there is no other boundary point for \widehat{X} . If $((x_k, y_k))_k$ is a Cauchy sequence in \widehat{X} that does not converge, then we must necessarily have that there is some $\tau > 0$ such that $y_k \leq \tau$ for each k . This follows from (7.1). That is, the sequence lies in $Z \times (0, \tau]$. From the fact that Z is compact, it follows that we must have $y_k \rightarrow 0$, that is, the sequence converges to a point in $Z \times \{0\}$. Hence $\partial \widehat{X} = Z \times \{0\}$.

Let $(x, y) \in X$; then $y > 0$. Let $\gamma : [0, \infty) \rightarrow X$ be given by $\gamma(t) = (x, y + t)$. Note that γ is arc-length parametrized with respect to the metric

d , but not with respect to the metric d_ρ . For $t > 0$, we see that

$$\ell_\rho(\gamma|_{[0,t]}) = \int_0^t \rho(y+s) ds = \begin{cases} \frac{1}{\beta-1} \left[\frac{1}{y^{\beta-1}} - \frac{1}{(y+t)^{\beta-1}} \right] & \text{if } y \geq 1, \\ 1 - y + \frac{1}{\beta-1} \left[1 - \frac{1}{(t+y)^{\beta-1}} \right] & \text{if } y < 1 \leq y+t, \\ t & \text{if } y+t < 1. \end{cases}$$

Moreover, for $t > 0$ we have that

$$\text{dist}_{d_\rho}(\gamma(t), Z \times \{0\}) = \int_0^{y+t} \rho(s) ds = \begin{cases} 1 + \frac{1}{\beta-1} \left[1 - \frac{1}{(y+t)^{\beta-1}} \right] & \text{if } y+t \geq 1, \\ y+t & \text{if } y+t \leq 1. \end{cases}$$

As $y > 0$, it follows from the above two computations that for each $t > 0$ we have $\text{dist}_{d_\rho}(\gamma(t), Z \times \{0\}) \geq \ell_\rho(\gamma|_{[0,t]})$. Hence, \widehat{X} is a John domain with John center ∞ and John constant $C_J = 1$. \square

7.5. Conclusion: the reconciliation. Now we have the tools necessary to compare the two fractional Laplace operators in the case that (Z, d_Z, μ_Z) is a compact doubling metric measure space supporting a 2-Poincaré inequality: the first considered in [33] and corresponding to the operator \mathcal{A} described in Subsection 6.2, and the second constructed in this paper and denoted $(-\Delta_2)^\theta$.

The domain $X = Z \times (0, \infty)$ was considered in [33]. Moreover, for $-1 < a < 1$ (corresponding to the relationship $\theta = (1-a)/2$), X is equipped with the weighted product measure $d\mu_X(x, y) = y^a d\mu_Z(x) dy$. With such a measure, it is clear to see that the measure $\nu = \mu_Z$ satisfies the co-dimension condition (1.1) with $\Theta = a + 1$.

It was shown in [33] that a function $u \in B_{2,2}^\theta(Z)$ satisfies $\mathcal{A}u = f$ on $Z = \partial X$ if and only if its Cheeger-harmonic extension, also denoted u , to X satisfies $\int_X g_u^2 d\mu_X < \infty$ and in addition, satisfied

$$(7.13) \quad \lim_{y \rightarrow 0^+} y^a \partial_y u = f.$$

Here \mathcal{A} on Z was defined via the spectral decomposition theorem, corresponding to the fractional Laplacian $(\Delta_2)^\theta$ in the Euclidean setting of [20]. Observe that as (Z, μ_Z) is compact, doubling, and supports a 2-Poincaré inequality, the measure μ_X is doubling and supports a 2-Poincaré inequality as well; this was shown in [33].

Now, when we transform $X = Z \times [0, \infty)$ into $(\widehat{Z}, d_\rho, \mu_\omega)$ as described at the beginning of Section 7, we obtain a doubling metric measure space supporting a 2-Poincaré inequality and the co-dimension condition as outlined in Conditions (H0), (H1), and (H2) in the current paper. Furthermore, with the transformation of the Cheeger differential structure as explained in this section, we also see that functions on X that were Cheeger 2-harmonic in $Z \times (0, \infty)$ are also Cheeger 2-harmonic in $\Omega := \widehat{X} \setminus (Z \times \{0\})$; moreover, d_ρ is locally isometric to the original metric d near $Z = \partial\Omega$, and the measure $\mu_\omega = \mu_X$ on $Z \times [0, 1)$. It follows that the condition $\lim_{y \rightarrow 0^+} y^a \partial_y u = f$ is satisfied by u if and only if u satisfies Condition (c) of our Theorem 1.5. Moreover, the trace of a function on Ω to $\partial\Omega$ is the same trace as the one for functions on $Z \times (0, \infty)$ to Z .

Thus the correspondence between the construction of [33] and ours is as follows.

- A function u on Z constructed in [33] to solve the equation $\mathcal{A}u = f$, is extended as a Cheeger 2-harmonic function, denoted \hat{u} , to $Z \times (0, \infty)$ so that the trace of \hat{u} at $\partial(Z \times (0, \infty)) = Z$ is u . It is shown there that \hat{u} also satisfies (7.13).
- It was also shown in [33] that \hat{u} has finite Dirichlet energy in $Z \times (0, \infty)$.
- The same function \hat{u} is then Cheeger 2-harmonic in the transformed domain $(Z \times (0, \infty), d_\rho, \mu_\omega)$, as discussed above. Moreover, \hat{u} has finite Dirichlet energy in this transformed domain.
- By Lemma 7.2 we know that the point ∞ is removable for Cheeger 2-harmonicity of functions with finite Dirichlet energy. Hence \hat{u} is Cheeger 2-harmonic in $\Omega = \hat{X} \setminus (Z \times \{0\})$.
- Combining the above-listed points, we see that \hat{u} satisfies Condition (c) of Theorem 1.5, and hence satisfies the equation $(-\Delta_2)^\theta u = f$ on Z , where θ and a are related by $\theta = (1 - a)/2 = 1 - \Theta/2$ (with $\Theta = a + 1$). Here, $(-\Delta_2)^\theta$ is as constructed using the bilinear form \mathcal{E}_T as described in Section 6 corresponding to $p = 2$.
- Since solutions to both problems exist and are unique up to additive constants, it follows that the two approaches give the same solution.

APPENDIX A. REMOVING BOUNDEDNESS CONDITION ON f IN [55] AND ESTABLISHING EXISTENCE AND BOUNDEDNESS OF SOLUTIONS

In this appendix we gather together results, and their proofs, that are adaptations of the results from [55] to our setting. In particular, we replace the boundedness condition on f , as required in [55], with the more natural condition $f \in L^{p'}(\partial\Omega)$, where $p' = p/(p-1)$ is the Hölder dual of p . We will first prove that for each $f \in L^{p'}(\partial\Omega)$ a solution to the Neumann boundary value problem on Ω exists with boundary data f . To this end, we use the adapted energy

$$I_f(v) := \int_{\Omega} |\nabla v|^p d\mu - p \int_{\partial\Omega} T v f d\nu$$

for $v \in N^{1,p}(\Omega)$. We set $\alpha := \inf_v I_f(v)$, with the infimum over all $v \in N^{1,p}(\Omega)$. Note that $\alpha \leq 0$. Solutions to the Neumann boundary value problem are precisely those functions u for which $I_f(u) = \alpha$; thus to consider existence of solutions, it suffices first to consider only those $v \in N^{1,p}(\Omega)$ for which $I_f(v) \leq 0$.

Lemma A.1. *Let $f \in L^{p'}(\partial\Omega)$ with $\int_{\partial\Omega} f d\nu = 0$. Then there is a constant $C > 0$, which is independent of f , such that whenever $v \in N^{1,p}(\Omega)$ such that $I_f(v) \leq 0$ and $\int_{\Omega} v d\nu = 0$, we have that*

$$\|\nabla v\|_{L^p(\Omega)}^{p-1} \leq C \|f\|_{L^{p'}(\partial\Omega)}.$$

The constant C depends only on the constant related to the Poincaré inequality on Ω , the diameter of Ω , p , and the norm-bound of the trace operator $T : N^{1,p}(\Omega) \rightarrow B_{p,p}^\theta(\partial\Omega)$. Moreover, for such v , we have that

$$\|\nabla v\|_{L^p(\Omega)}^p - C \|f\|_{L^{p'}(\partial\Omega)} \|\nabla v\|_{L^p(\Omega)} \leq I_f(v) \leq 0.$$

Proof. Since Ω is bounded and supports a p -Poincaré inequality, and since $\int_\Omega u \, d\mu = 0$, we have that

$$(A.2) \quad \int_\Omega |v|^p \, d\mu \leq C \int_\Omega |\nabla v|^p \, d\mu.$$

Let $v \in N^{1,p}(\Omega)$ such that $I_f(v) \leq 0$. Note by Theorem 2.4 and (A.2) that

$$\begin{aligned} \left| \int_{\partial\Omega} v f \, d\nu \right| &\leq \int_{\partial\Omega} |f| |v| \, d\nu \leq \|f\|_{L^{p'}(\partial\Omega)} \|Tu\|_{L^p(\partial\Omega)} \\ &\leq C \|f\|_{L^{p'}(\partial\Omega)} \|u\|_{N^{1,p}(\Omega)} \\ &\leq C \|f\|_{L^{p'}(\partial\Omega)} \|\nabla u\|_{L^p(\Omega)}. \end{aligned}$$

The claim follows from the above estimate. \square

As a corollary to the above lemma, we see that

$$0 \geq \alpha \geq \frac{C}{p} \|f\|_{L^{p'}(\partial\Omega)} - \frac{C^{1+1/p}}{p^{1/p}} \|f\|_{L^{p'}(\partial\Omega)}^{1+1/p} > -\infty.$$

From the above lemma it is clear that if $(v_k)_k$ is a sequence of functions in $N^{1,p}(\Omega)$ such that the sequence of numbers $I_f(v_k)$ converges monotonically to α , then the sequence of Cheeger derivatives ∇v_k forms a bounded sequence in $L^p(\Omega, \mathbb{R}^N)$. Thus by the assumption $\int_\Omega v_k \, d\mu = 0$, (A.2) shows that the sequence v_k itself is also bounded in $N^{1,p}(\Omega)$. Invoking Mazur's lemma yields that a convex combination of this sequence v_k converges in the reflexive Banach space self-same convex combination to the boundary $\partial\Omega$ converges to the trace of u_f in $B_{p,p}^\theta(\partial\Omega)$ because the trace operator is bounded and linear. Finally, as in the proof of Theorem 5.4, the convexity of the map $v \mapsto I_f(v)$ tells us that $I_f(u_f) \leq \alpha$, and so we must have $I_f(u_f) = \alpha$. This proves the existence of solutions to the Neumann boundary value problem. This proves the existence part of Theorem 2.5.

The rest of this appendix is devoted to showing that such solutions are bounded, thus completing the proof of Theorem 2.5.

Lemma A.3. *Let u be the solution to the Neumann boundary value problem on Ω with boundary data $f \in L^{p'}(\partial\Omega)$ such that $\int_\Omega u \, d\mu = 0$. Then*

$$\int_\Omega |u|^p \, d\mu \leq C \int_\Omega |\nabla u|^p \, d\mu \leq C^2 \int_{\partial\Omega} |f|^{p'} \, d\nu.$$

The constant C depends only on the structural constants of Ω .

The above lemma is a consequence of [55, Proposition 4.1] when the Cheeger differential structure is replaced with the upper gradients in the formulation of the Neumann boundary value problem.

Proof. For each $v \in N^{1,p}(\Omega)$ we let $I(v) = \int_{\Omega} |\nabla v|^p d\mu - p \int_{\partial\Omega} v f d\nu$. Since u is a minimizer of I as shown in Theorem 1.5, and as $I(0) = 0$, it follows that $I(u) \leq 0$, that is,

$$\int_{\Omega} |\nabla u|^p d\mu \leq p \int_{\partial\Omega} u f d\nu.$$

Hence by the trace theorem [54, Theorem 1.1], we have

$$\begin{aligned} \int_{\Omega} |\nabla u|^p d\mu &\leq p \|u\|_{L^p(\partial\Omega)} \|f\|_{L^{p'}(\partial\Omega)} \leq C (\|u\|_{L^p(\Omega)} + \|\nabla u\|_{L^p(\Omega)}) \|f\|_{L^{p'}(\partial\Omega)} \\ &\leq C \|\nabla u\|_{L^p(\Omega)} \|f\|_{L^{p'}(\partial\Omega)}, \end{aligned}$$

where we used (A.2) in the last line above. It follows that

$$\|\nabla u\|_{L^p(\Omega)}^{p-1} \leq C \|f\|_{L^{p'}(\partial\Omega)},$$

from whence the second inequality in the statement of the lemma follows. \square

The proof of boundedness of solutions to the Neumann problem as given in [55] uses the boundedness of the Neumann data f . In this appendix we show how to modify the proof of [55, Theorem 5.2] by relaxing the requirement that $f \in L^\infty(\partial\Omega)$ to $f \in L^{p'}(\partial\Omega)$. We also relax the co-dimension 1 condition to a co-dimension Θ condition, namely that there is some $0 < \Theta < p$ so that for each $x \in \partial\Omega$ and $0 < r \leq \text{diam}(\partial\Omega)$ we have

$$(A.4) \quad \nu(B(x, r)) \approx \frac{\mu(B(x, r))}{r^\Theta}.$$

Here the ball $B(x, r)$ is the ball in $\bar{\Omega}$, with $\mu(\partial\Omega) = 0$ and $\nu(\Omega) = 0$. Observe that since μ is a doubling measure, there is an exponent $s > 0$ such that for all $x \in \bar{\Omega}$ and $0 < r < R \leq \text{diam}(\Omega)$,

$$(A.5) \quad \left(\frac{r}{R}\right)^s \lesssim \frac{\mu(B(x, r))}{\mu(B(x, R))}.$$

Note that we can make s as large as we like; thus, once $\Theta < p$ is fixed, with $1 < p < \infty$, we can then choose $s > 0$ so that $p < s$. Moreover, given the assumptions on Ω , we can replace the original metric with the biLipschitz equivalent length-metric, and so we can assume that Ω is a length space (the class of upper gradients and the Cheeger differential structure are not changed by this); it follows from [54, Proposition 4.21] that given a choice of $p < \tilde{p} < p^* := \frac{p(s-\Theta)}{s-p}$, the trace operator $T : N^{1,p}(\Omega) \rightarrow L^{\tilde{p}}(\partial\Omega)$ is local in the sense that whenever $x \in \partial\Omega$ and $r > 0$, we have

$$(A.6) \quad \|Tu - u_{B(x,r)}\|_{L^{\tilde{p}}(B(x,r) \cap \partial\Omega)} \lesssim r^{(\tilde{p}^{-1} - p^{*-1})(s-\Theta)} \|\nabla u\|_{L^p(B(x,r))}.$$

Up to Remark 5.9 of [55] holds without any change even if we only assume that $f \in L^{p'}(\partial\Omega)$ where $p' = p/(p-1)$ is the Hölder dual of p . The subsequent parts of Section 5 of [55] are modified as follows.

Lemma 5.10 of [55] is modified to the following. Note that if f is bounded, then the integrability condition of f required by us, namely $f \in L^{p'}(\partial\Omega)$, trivially holds. In the lemma below, $\tilde{p} \in (p, p^*)$ is the exponent associated with the trace operator as in (A.6), and $\tilde{p}' = \tilde{p}/(\tilde{p}-1)$ is the Hölder dual

of \tilde{p} and p' the Hölder dual of p . As $\tilde{p} > p$, it follows that $p' > \tilde{p}'$, and so necessarily $f \in L^{\tilde{p}'}(\partial\Omega)$, and this property is needed instead of boundedness of f in the proof of the lemma.

We now turn to the boundedness of the solution of the Neumann problem. For ease of notation, in the following, integrals over balls B with respect to μ stand in for integrals over $B \cap \Omega$, and integrals with respect to ν stand in for integrals over $B \cap \partial\Omega$.

Lemma A.7. *Suppose that $\int_{\partial\Omega} f \, d\nu = 0$ with $f \in L^{p'}(\partial\Omega)$, and set $t = \frac{p' - \tilde{p}'}{p' - 1}$. Let u be a solution to the Neumann boundary value problem with boundary data f . For $x \in \partial\Omega$, $k, h \in \mathbb{R}$ with $k > h$, and $0 < R/2 \leq r < R \leq \text{diam}(\partial\Omega)$, set*

$$u(k, r) := \left(\int_{B(x, r)} (u - k)_+^p \, d\mu \right)^{1/p}, \quad \psi(k, r) := \int_{B(x, r)} |f|(u - k)_+ \, d\nu.$$

Then

$$(A.8) \quad \begin{aligned} u(k, r) &\leq C \left(\frac{u(h, R)}{k - h} \right)^{1-1/\kappa} \left[\frac{R}{R - r} u(h, R) + R^{1-\Theta/p} \psi(h, R)^{1/p} \right], \\ \psi(k, r) &\leq C_{f, R} \left(\frac{\psi(h, R)}{k - h} \right)^{t/\tilde{p}'} \left[\frac{R^{\aleph_1}}{R - r} u(h, R) + R^{\aleph_2} \psi(h, R)^{1/p} \right], \end{aligned}$$

where $\kappa = s/(s - p)$ and the constants \aleph_1, \aleph_2 are positive and independent of f , u , x , r , and R , while $C_{f, R} > 1$ is independent of u , x , and r .

The choice of $\kappa = s/(s - p)$ comes from the Sobolev-Poincaré inequality on $\overline{\Omega}$, namely, $(\kappa p, p)$ -Poincaré inequality; see [43]. We remind the readers that the parameter s is the lower mass bound dimension of μ from (A.5).

Proof. As in the proof of [55, Lemma 5.10], we pick a cut-off function η that is Lipschitz on $\overline{\Omega}$, with $\eta = 1$ on $B(x, r)$, $\eta = 0$ on $\overline{\Omega} \setminus B(x, (R + r)/2)$ and apply Hölder's inequality and the $(\kappa p, p)$ -Poincaré inequality to $\eta(u - k)_+$ to obtain

$$\int_{B(x, r)} (u - k)_+^p \, d\mu \lesssim \left(\frac{\mu(A(k, r))}{\mu(B(x, r))} \right)^{1-1/\kappa} R^p \int_{B(x, (R+r)/2)} |\nabla(\eta(u - k)_+)|^p \, d\mu.$$

Here $A(k, r) = \{y \in \overline{\Omega} : u(y) > k\} \cap B(x, r)$. Invoking the Leibniz rule (with $|\nabla\eta| \leq 2(R - r)^{-1} \chi_{B(x, (R+r)/2) \setminus B(x, r)}$) and the version of De Giorgi-type inequality from [55, Theorem 5.3], we now obtain

$$\begin{aligned} u(k, r)^p &\lesssim \left(\frac{\mu(A(k, r))}{\mu(B(x, r))} \right)^{1-1/\kappa} \left[\frac{R^p}{(R - r)^p} \int_{B(x, R)} (u - k)_+^p \, d\mu \right. \\ &\quad \left. + \frac{1}{\mu(B(x, R))} \int_{B(x, R)} |f|(u - k)_+ \, d\nu \right]. \end{aligned}$$

Now applying the codimensionality (A.4), we obtain

$$u(k, r)^p \lesssim \left(\frac{\mu(A(k, r))}{\mu(B(x, r))} \right)^{1-1/\kappa} \left[\frac{R^p}{(R-r)^p} u(k, R)^p + R^{p-\Theta} \int_{B(x, R)} |f|(u-k)_+ d\nu \right].$$

Now, as in [55, bottom of page 2446], we see that when $h < k$,

$$\mu(A(k, r)) \lesssim \frac{\mu(B(x, r))}{(k-h)^p} u(h, R)^p.$$

Combining the above two inequalities proves the first of the two inequalities in (A.8) (note that we have used the doubling property of μ as well as the facts that $R/2 \leq r < R$ and $u(k, R) \leq u(h, R)$, $\psi(k, R) \leq \psi(h, R)$ here).

Now we turn our attention to $\psi(k, r)$ where the bulk of the modification lies. We set $\aleph_0 := (\tilde{p}^{-1} - p^{*-1})(s - \Theta)$. By Hölder's inequality and (A.6), Hölder's inequality again, and then by the definition of $u(k, r)$,

$$\begin{aligned} \psi(k, r) &\leq \left(\int_{B(x, r)} (u-k)_+^{\tilde{p}} d\nu \right)^{\frac{1}{\tilde{p}}} \left(\int_{B(x, r)} |f\chi_{A(k, r)}|^{\tilde{p}'} d\nu \right)^{\frac{1}{\tilde{p}'}} \\ &\lesssim \nu(B(x, r))^{-\frac{1}{\tilde{p}}} \left(\int_{B(x, r)} |f\chi_{A(k, r)}|^{\tilde{p}'} d\nu \right)^{\frac{1}{\tilde{p}'}} \times \\ &\quad \left[r^{\aleph_0} \left(\int_{B(x, r)} |\nabla(u-k)_+|^p d\mu \right)^{\frac{1}{p}} + \nu(B(x, r))^{\frac{1}{\tilde{p}}} \int_{B(x, r)} (u-k)_+ d\mu \right] \\ &\lesssim \left(\int_{B(x, r)} |f\chi_{A(k, r)}|^{\tilde{p}'} d\nu \right)^{\frac{1}{\tilde{p}'}} \left[\frac{r^{\aleph_0}}{\nu(B(x, r))^{\frac{1}{\tilde{p}}}} \left(\int_{B(x, r)} |\nabla(u-k)_+|^p d\mu \right)^{\frac{1}{p}} + u(k, r) \right]. \end{aligned}$$

Combining this inequality with the De Giorgi-type inequality from [55, Theorem 5.3] yields

$$\begin{aligned} \psi(k, r) &\lesssim \left(\int_{B(x, r)} |f\chi_{A(k, r)}|^{\tilde{p}'} d\nu \right)^{1/\tilde{p}'} \left[u(k, r) + \frac{r^{\aleph_0}}{R-r} \frac{\mu(B(x, R))^{1/p}}{\nu(B(x, r))^{1/\tilde{p}}} u(k, R) \right. \\ &\quad \left. + r^{\aleph_0} \nu(B(x, R))^{1/p-1/\tilde{p}} \psi(k, R)^{1/p} \right]. \end{aligned}$$

Here we have also used the facts that nu is doubling and $R/2 \leq r < R$. Now using the co-dimensionality of ν with respect to μ , we obtain

$$\begin{aligned} \psi(k, r) &\lesssim \left(\int_{B(x, r)} |f\chi_{A(k, r)}|^{\tilde{p}'} d\nu \right)^{1/\tilde{p}'} \left[u(k, r) + \frac{R^{\aleph_0+\Theta/p}}{R-r} \nu(B(x, R))^{1/p-1/\tilde{p}} u(k, R) \right. \\ &\quad \left. + r^{\aleph_0} \nu(B(x, R))^{1/p-1/\tilde{p}} \psi(k, R)^{1/p} \right]. \end{aligned}$$

As $\tilde{p} > p$ and $\nu(\partial\Omega) < \infty$, we have
(A.9)

$$\psi(k, r) \lesssim \left(\int_{B(x, r)} |f \chi_{A(k, r)}|^{\tilde{p}'} d\nu \right)^{1/\tilde{p}'} \left[\left(1 + \frac{R^{\aleph_0 + \Theta/p}}{R - r} \right) u(k, R) + r^{\aleph_0} \psi(k, R)^{1/p} \right].$$

Note that by assumption, $f \in L^{p'}(\partial\Omega)$. As $\tilde{p} > p$, it follows that $f \in L^{\tilde{p}'}(\partial\Omega)$. We set $t = \frac{p' - \tilde{p}'}{p' - 1}$ as in the statement of the lemma. As $p < \tilde{p}$ and $p' > 1$, we have that $t > 0$. Moreover, as $\tilde{p}' > 1$, it follows that $t < 1$. Now by Hölder's inequality and the fact that $(\tilde{p}' - t)/(1 - t) = p'$,

$$\begin{aligned} \int_{B(x, r)} |f \chi_{A(k, r)}|^{\tilde{p}'} d\nu &\leq \left(\int_{B(x, r)} |f| \chi_{A(k, r)} d\nu \right)^t \left(\int_{B(x, r)} |f|^{(\tilde{p}' - t)/(1 - t)} d\nu \right)^{1 - t} \\ &= \left(\int_{B(x, r)} |f| \chi_{A(k, r)} d\nu \right)^t \left(\int_{B(x, r)} |f|^{p'} d\nu \right)^{\frac{\tilde{p}' - 1}{p' - 1}} \\ &\leq C \left(\int_{B(x, r)} |f| \chi_{A(k, r)} d\nu \right)^t \left(\int_{B(x, R)} |f|^{p'} d\nu \right)^{\frac{\tilde{p}' - 1}{p' - 1}}. \end{aligned}$$

We set

$$C_{f, R} := C \left(\int_{B(x, R)} |f|^{p'} d\nu \right)^{(\tilde{p}' - 1)/(p' - 1)}$$

Then by the above,

$$\int_{B(x, r)} |f \chi_{A(k, r)}|^{\tilde{p}'} d\nu \leq C_{f, R} \left(\int_{B(x, r)} |f| \chi_{A(k, r)} d\nu \right)^t.$$

When $h, k \in \mathbb{R}$ with $h < k$, we have

$$\int_{B(x, r)} |f| \chi_{A(k, r)} d\nu \leq \frac{1}{k - h} \int_{B(x, r)} |f| (u - h)_+ d\nu = \frac{\psi(h, r)}{k - h} \leq \frac{\psi(h, R)}{k - h}.$$

Combining this with inequality (A.9), we obtain

$$\psi(k, r) \lesssim C_{f, R}^{1/\tilde{p}'} \left(\frac{\psi(h, R)}{k - h} \right)^{t/\tilde{p}'} \left[\left(1 + \frac{R^{\aleph_0 + \Theta/p}}{R - r} \right) u(k, R) + R^{\aleph_0} \psi(k, R)^{1/p} \right].$$

As $R \leq \text{diam}(\partial\Omega) < \infty$ and $R > R - r$, we obtain the second part of (A.8) by choosing $\aleph_2 = \aleph_0$ and $\aleph_1 = \aleph_0 + \Theta/p$. \square

Now we are ready to prove the boundedness of u on $B(x, R)$ even when f is not bounded. To do so, fix $k_0 \in \mathbb{R}$. We fix $d > 0$ for now, but we will add some conditions in on d towards the end. For non-negative integers n we set

$$r_n = \frac{R}{2}(1 + 2^{-n}), \quad k_n = k_0 + d(1 - 2^{-n}).$$

Note that $0 < k_{n+1} - k_n = 2^{-(n+1)}d$ and $0 < r_n - r_{n+1} = 2^{-(n+2)}R$. Then from the above lemma, we see that

$$u(k_{n+1}, r_{n+1}) \leq C \left(\frac{u(k_n, r_n)}{2^{-(n+1)}d} \right)^{1-1/\kappa} \left[2^{n+2}u(k_n, r_n) + R^{1-\Theta/p}\psi(k_n, r_n)^{1/p} \right].$$

Setting $\alpha = 1 - 1/\kappa < 1$ and noting that $R \leq \text{diam}(\partial\Omega) < \infty$, we obtain

$$u(k_{n+1}, r_{n+1}) \leq C \frac{2^{(n+1)\alpha}}{d^\alpha} \left[2^{n+2}u(k_n, r_n)^{1+\alpha} + u(k_n, r_n)^\alpha \psi(k_n, r_n)^{1/p} \right],$$

and as $2^{n+2} \geq 1$, we obtain

$$(A.10) \quad u(k_{n+1}, r_{n+1}) \leq C \frac{2^{n(\alpha+1)}}{d^\alpha} \left[u(k_n, r_n)^{1+\alpha} + u(k_n, r_n)^\alpha \psi(k_n, r_n)^{1/p} \right].$$

Similarly, setting $\beta = t/\tilde{p}' < 1$, we have from the above lemma that

$$(A.11) \quad \psi(k_{n+1}, r_{n+1}) \leq C_{f,R} \frac{2^{n(\beta+1)}}{d^\beta} \left[u(k_n, r_n) \psi(k_n, r_n)^\beta + \psi(k_n, r_n)^{\beta+1/p} \right].$$

Lemma A.12. *There exist a choice of positive real numbers d , σ , and τ such that with k_n , r_n as defined above for each non-negative integer n , we have*

$$u(k_n, r_n) \leq 2^{-\sigma n} u(k_0, R), \quad \psi(k_n, r_n) \leq 2^{-\tau n} \psi(k_0, R).$$

Note that with d, σ , and τ as in the above lemma, we obtain that

$$0 = \lim_{n \rightarrow \infty} u(k_n, r_n) = u(k_0 + d, R/2),$$

from whence we can conclude that $u \leq k_0 + d$ on $B(x, R/2)$. Hence, to prove the boundedness of u on $B(x, R/2)$ it suffices to prove the above lemma.

Proof. If $u(k_0, R) = 0$, then $u \leq k_0$ on $B(x, R)$, and we have the boundedness. Therefore, without loss of generality, we assume that $u(k_0, R) > 0$. If $\psi(k_0, R) = 0$, then again for each n we have $\psi(k_n, r_n) = 0$, and the required inequality for ψ would be satisfied, and then we can directly focus on the part of the proof below that is relevant to $u(k_n, r_n)$. Hence we now assume also that $\psi(k_0, R) > 0$, and prove the claim via induction on n . The base case $n = 0$ holds trivially. So suppose that n is a non-negative integer such that the claims hold for n ; and we wish to then show that the claims also hold for $n + 1$.

By (A.11) and by the hypotheses holding for n ,

$$\begin{aligned} \psi(k_{n+1}, r_{n+1}) &\leq C \frac{2^{n(\beta+1)}}{d^\beta} \left[2^{-(\tau\beta+\sigma)n} u(k_0, R) \psi(k_0, R)^\beta + 2^{-\tau n(\beta+1/p)} \psi(k_0, R)^{\beta+1/p} \right] \\ &\leq \frac{C}{d^\beta} \left[2^{-n(\sigma+\tau\beta-\beta-1)} \psi(k_0, R)^\beta u(k_0, R) + 2^{-n(\tau\beta+\tau/p-\beta-1)} \psi(k_0, R)^{\beta+1/p} \right] \\ &\leq 2^{-\tau(n+1)} \psi(k_0, R) \frac{2^\tau C}{d^\beta} \left[2^{-n(\sigma+\tau\beta-\beta-1-\tau)} \frac{u(k_0, R)}{\psi(k_0, R)^{1-\beta}} \right. \\ &\quad \left. + 2^{-n(\tau\beta+\tau/p-\beta-1-\tau)} \psi(k_0, R)^{\beta+1/p-1} \right]. \end{aligned}$$

Similarly, by (A.10) and by the hypothesis,

$$u(k_{n+1}, r_{n+1}) \leq 2^{-\sigma(n+1)} u(k_0, R) \frac{C}{d^\beta} \left[2^{-n(\sigma+\sigma\alpha-\alpha-1-\sigma)} u(k_0, R)^\alpha + 2^{-n(\sigma\alpha+\tau/p-\alpha-1-\sigma)} \frac{\psi(k_0, R)^{1/p}}{u(k_0, R)^{1-\alpha}} \right].$$

We choose σ and τ such that

$$\begin{aligned} \sigma + \sigma\alpha - \alpha - 1 - \sigma &\geq 0, & \sigma\alpha + \frac{\tau}{p} - \alpha - 1 - \sigma &\geq 0, \\ \sigma + \tau\beta - \beta - 1 - \tau &\geq 0, & \tau\beta + \frac{\tau}{p} - \beta - 1 - \tau &\geq 0. \end{aligned}$$

Such a choice is possible because $\beta = t/\tilde{p}' < 1 < p' = p/(p-1)$, see the conditions listed at the top of [55, page 2450]. Subsequently, we choose d large enough so that

$$d^\beta \geq C \left[u(k_0, R)^\alpha + \frac{u(k_0, R)}{\psi(k_0, R)^{1-\beta}} + \psi(k_0, R)^{\beta+1/p-1} \frac{\psi(k_0, R)^{1/p}}{u(k_0, R)^{1-\alpha}} \right]$$

to complete the proof of the lemma. \square

REFERENCES

- [1] H. Aikawa, N. Shanmugalingam: *Carleson-type estimates for p -harmonic functions and the conformal Martin boundary of John domains in metric measure spaces*. Michigan Math. J. **53** (2005), no. 1, 165–188.
- [2] L. Ambrosio, N. Gigli, G. Savaré: *Bakry–Émery curvature-dimension condition and Riemannian Ricci curvature bounds*. Ann. Probab. **43** (2015), no. 1, 339–404.
- [3] L. Ambrosio, N. Gigli, G. Savaré: *Metric measure spaces with Riemannian Ricci curvature bounded from below*. Duke Math. J. **163** (2014), no. 7, 1405–1490.
- [4] Z. Balogh, S. Buckley: *Sphericalization and flattening*. Conform. Geom. Dyn. **9** (2005), 76–101.
- [5] A. Banerjee, P. Garain, J. Kinnunen: *Some local properties of subsolutions and supersolutions for a doubly nonlinear nonlocal parabolic p -Laplace equation*. preprint, <https://arxiv.org/abs/2010.05727> (2020).
- [6] A. Banerjee, N. Garofalo: *Monotonicity of generalized frequencies and the strong unique continuation property for fractional parabolic equations*. Adv. Math. **336** (2018), 149–241.
- [7] A. Banerjee, N. Garofalo, I. Munive, D.-M. Nhieu: *The Harnack inequality for a class of nonlocal parabolic equations*. Commun. Contemp. Math. **23** (2021), no. 6, Paper No. 2050050, 23 pp.
- [8] V. Banica, M. González, M. Sáez: *Some constructions for the fractional Laplacian on noncompact manifolds*. Rev. Mat. Iberoam. **31** (2015), no. 2, 681–712.
- [9] M. Barlow, R. Bass, Z.-Q. Chen, M. Kassmann: *Non-local Dirichlet forms and symmetric jump processes*. Trans. Amer. Math. Soc. **361** (2009), no. 4, 1963–1999.
- [10] F. Baudoin, Q. Lang, Y. Sire: *Powers of generators on Dirichlet spaces and applications to Harnack principles*. preprint available at <https://arxiv.org/abs/2010.01036> (2020).
- [11] P. Bérard, D. L. Webb: *One can't hear orientability of surfaces*. Math. Z. **300** (2022), 139–160.
- [12] A. Björn, J. Björn: *Nonlinear potential theory on metric spaces*. EMS Tracts in Mathematics, **17**, European Mathematical Society (EMS), Zürich, 2011. xii+403 pp.
- [13] A. Björn, J. Björn, X. Li: *Sphericalization and p -harmonic functions on unbounded domains in Ahlfors regular spaces*. J. Math. Anal. Appl. **474** (2019), no. 2, 852–875.

- [14] A. Björn, J. Björn, N. Shanmugalingam: *Extension and trace results for doubling metric measure spaces and their hyperbolic fillings*. J. Math. Pures Appl. (9) **159** (2022), 196–249.
- [15] J. Björn, N. Shanmugalingam: *Poincaré inequalities, uniform domains and extension properties for Newton–Sobolev functions in metric spaces*. J. Math. Anal. Appl. **332** (2007) 190–208.
- [16] M. Bonk, E. Saksman: *Sobolev spaces and hyperbolic fillings*. J. Reine Angew. Math **737** (2018), 161–187.
- [17] M. Bonk, E. Saksman, T. Soto: *Triebel–Lizorkin spaces on metric spaces via hyperbolic fillings*. Indiana Univ. Math. J. **67** (2018), 1625–1663.
- [18] M. Bonk, O. Schramm: *Embeddings of Gromov hyperbolic spaces*. Geom. Funct. Anal. **10** (2000), 266–306.
- [19] M. Bourdon, H. Pajot: *Cohomologie l_p et espaces de Besov*. J. Reine Angew. Math. **558** (2003), 85–108.
- [20] L. Caffarelli, L. Silvestre: *An extension problem related to the fractional Laplacian*. Comm. Partial Differential Equations **32** (2007), no. 7–9, 1245–1260.
- [21] L. Caffarelli, M. Soria-Carro: *On a family of fully nonlinear integro-differential operators: From fractional Laplacian to nonlocal Monge–Ampère*. Preprint (2021) <https://arxiv.org/pdf/2111.12781.pdf>
- [22] L. Caffarelli, P. Stinga: *Fractional elliptic equations, Caccioppoli estimates and regularity*. Ann. I. H. Poincaré: Analyse Nonlinéaire **33** (2016) 767–807.
- [23] L. Capogna, R. Gibara, R. Korte, N. Shanmugalingam, *Fractional p -Laplacians via Neumann problems in unbounded metric measure spaces*, preprint (2024).
- [24] A. Chang, M. González: *Fractional Laplacian in conformal geometry*. Adv. Math. **226** (2011), no. 2, 1410–1432.
- [25] J. Cheeger: *Differentiability of Lipschitz functions on metric measure spaces*. Geom. Funct. Anal. **9** (1999), no. 3, 428–517.
- [26] W. Chen, C. Li: *Maximum principles for the fractional p -Laplacian and symmetry of solutions*. Adv. Math. **335** (2018), 735–758.
- [27] Z.-Q. Chen, T. Kumagai: *Heat kernel estimates for stable-like processes on d -sets*. Stochastic Process. Appl. **108** (2003), no. 1, 27–62.
- [28] Z.-Q. Chen, T. Kumagai: *Heat kernel estimates for jump processes of mixed types on metric measure spaces*. Probab. Theory Related Fields **140** (2008), no. 1–2, 277–317.
- [29] Z.-Q. Chen, T. Kumagai, J. Wang: *Stability of parabolic Harnack inequalities for symmetric non-local Dirichlet forms*. J. Eur. Math. Soc. **22** (2020), no. 11, 3747–3803.
- [30] E. Durand-Cartagena, X. Li: *Preservation of p -Poincaré inequality for large p under sphericalization and flattening*. Illinois J. Math. **59** (2015), no. 4, 1043–1069.
- [31] E. Durand-Cartagena, X. Li: *Preservation of bounded geometry under sphericalization and flattening: quasiconvexity and ∞ -Poincaré inequality*. Ann. Acad. Sci. Fenn. Math. **42** (2017), no. 1, 303–324.
- [32] E. Durand-Cartagena, N. Shanmugalingam: *An elementary proof of Cheeger’s theorem on reflexivity of Newton–Sobolev spaces of functions in metric measure spaces*. J. Anal. **21** (2013), 73–83.
- [33] S. Eriksson-Bique, G. Giovannardi, R. Korte, N. Shanmugalingam, G. Speight: *Regularity of solutions to the fractional Cheeger–Laplacian on Domains in metric spaces of bounded geometry*. J. Differential Equations **306** (2022), 590–632.
- [34] F. Ferrari, B. Franchi: *Harnack inequality for fractional sub-Laplacians in Carnot groups*. Math. Z. **279** (2015), no. 1–2, 435–458.
- [35] B. Franchi, P. Hajlasz, P. Koskela: *Definitions of Sobolev classes on metric spaces*. Ann. Inst. Fourier (Grenoble) **49** (1999), no. 6, 1903–1924.
- [36] M. Fukushima, Y. Oshima, M. Takeda: *Dirichlet forms and symmetric Markov processes*. De Gruyter Studies in Mathematics, **19** Walter de Gruyter & Co., Berlin (1994) x+392 pp. ISBN: 3-11-011626-X.
- [37] N. Garofalo: *Fractional thoughts*. New developments in the analysis of nonlocal operators, AMS Contemp. Math. **723** (2019), 1–135.

- [38] M. Giaquinta: *Multiple integrals in the calculus of variations and nonlinear elliptic systems*. Annals of Mathematics Studies, 105. Princeton University Press, Princeton, NJ, 1983. vii+297 pp.
- [39] M. Giaquinta: *Introduction to regularity theory for nonlinear elliptic systems*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 1993. viii+131 pp.
- [40] R. Gibara, N. Shanmugalingam: *Transformation of geometry of uniform domains under weights that depend on distance to the boundary*. manuscript, in preparation.
- [41] N. Gigli: *On the differential structure of metric measure spaces and applications*. Mem. Amer. Math. Soc. **236** (2015), no. 1113, vi+91 pp.
- [42] N. Gigli, A. Mondino: *A PDE approach to nonlinear potential theory in metric measure spaces*. J. Math. Pures Appl. (9) **100** (2013), no. 4, 505–534.
- [43] P. Hajlasz, P. Koskela: *Sobolev met Poincaré*. Mem. Amer. Math. Soc. **145** (2000), no. 688, x+101 pp.
- [44] Y. Han, D. Müller, D. Yang: *A theory of Besov and Triebel-Lizorkin spaces on metric measure spaces modeled on Carnot-Carathéodory spaces*. Abstr. Appl. Anal. 2008, Art. ID 893409, 250 pp.
- [45] J. Heinonen, P. Koskela: *Quasiconformal maps in metric spaces with controlled geometry*. Acta Math. **181** (1998), no. 1, 1–61.
- [46] J. Heinonen, P. Koskela, N. Shanmugalingam, J. Tyson: *Sobolev spaces on metric measure spaces: an approach based on upper gradients*. New Mathematical Monographs **27**, Cambridge University Press, Cambridge, U., 2015. xi+429 pp.
- [47] T. Kilpeläinen, J. Malý: *Degenerate elliptic equations with measure data and nonlinear potentials*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **19** (1992), no. 4, 591–613.
- [48] J. Kinnunen, V. Latvala: *Lebesgue points for Sobolev functions on metric spaces*. Rev. Mat. Iberoamericana **18** (2002), 685–700.
- [49] J. Kinnunen, N. Shanmugalingam: *Regularity of quasi-minimizers on metric spaces*. Manuscripta Math. **105** (2001), no. 3, 401–423.
- [50] P. Koskela, K. Rajala, N. Shanmugalingam: *Interpolation properties of Besov spaces defined on metric spaces*. Math. Nachr. **283** (2010), no. 2, 215–231.
- [51] X. Li: *Preservation of bounded geometry under transformations of metric spaces*. Thesis (Ph.D.)—University of Cincinnati. 2015. 140 pp.
- [52] X. Li, N. Shanmugalingam: *Preservation of bounded geometry under sphericalization and flattening*. Indiana Univ. Math. J. **64** (2015), no. 5, 1303–1341.
- [53] T. Mäkeläinen: *Removable sets for Hölder continuous p -harmonic functions on metric measure spaces*. Ann. Acad. Scient. Fenn. Mat. **33**, (2008), 605–624.
- [54] L. Malý: *Trace and extension theorems for Sobolev-type functions in metric spaces*. preprint, <https://arxiv.org/abs/1704.06344> (2017).
- [55] L. Malý, N. Shanmugalingam: *Neumann problem for p -Laplace equation in metric spaces using a variational approach: existence, boundedness, and boundary regularity*. J. Diff. Eq. **265**, No. 6 (2018), 2431–2460.
- [56] P. Mikkonen: *On the Wolff potential and quasilinear elliptic equations involving measures*. Ann. Acad. Sci. Fenn. Math. Diss. **104** (1996), 71 pp.
- [57] T. Ono, Hölder continuity of solutions to quasilinear elliptic equations with measure data, in *Potential theory in Matsue*, 327–338, Adv. Stud. Pure Math., 44, Math. Soc. Japan, Tokyo.
- [58] J. M. Rakotoson, W. P. Ziemer: *Local behavior of solutions of quasilinear elliptic equations with general structure*. Transactions Amer. Math. Soc. **319**, No. 2 (1990), 747–764.
- [59] N. Shanmugalingam: *Harmonic functions on metric spaces*. Illinois J. Math. **45** No. 3 (2001), 1021–1050.
- [60] E. M. Stein: *Harmonic Analysis: Real-variable methods, orthogonality, and oscillatory integrals*. Princeton University Press, Princeton, NJ., 1993.
- [61] P.R. Stinga, P.L. Torrea: *Extension problem and Harnack’s inequality for some fractional operators*. Comm. Partial Differential Equations **35** (2010), no. 11, 2092–2122.
- [62] X. Wang, G. Du: *Properties of solutions to fractional p -subLaplace equations on the Heisenberg group*. Boundary Value Problems (2020), 2020:128.

- [63] X. Zhong: *On nonhomogeneous quasilinear elliptic equations*. Dissertation, University of Jyväskylä, Ann. Acad. Sci. Fenn. Math. Diss. **117** (1998), 46 pp.

DEPARTMENT OF MATHEMATICS AND STATISTICS, SMITH COLLEGE, NORTHAMPTON, MA 01063, U.S.A.

Email address: lcapogna@smith.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, P.O. Box 210025, UNIVERSITY OF CINCINNATI, CINCINNATI, OH 45221-0025, U.S.A.

Email address: klinejp@mail.uc.edu

AALTO UNIVERSITY, DEPARTMENT OF MATHEMATICS AND SYSTEMS ANALYSIS, P.O. Box 11100, FI-00076 AALTO, FINLAND

Email address: riikka.korte@aalto.fi

DEPARTMENT OF MATHEMATICAL SCIENCES, P.O. Box 210025, UNIVERSITY OF CINCINNATI, CINCINNATI, OH 45221-0025, U.S.A.

Email address: shanmun@uc.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, KENYON COLLEGE, 201 N. COLLEGE DR., GAMBIER, OH 43022, U.S.A.

Email address: snipesm@kenyon.edu