

MABUCHI'S SOLITON METRIC AND RELATIVE D-STABILITY

TOMOYUKI HISAMOTO

ABSTRACT. For Fano manifolds T. Mabuchi introduced a generalization of the Kähler-Einstein metric, which is characterized as the critical point of the Ricci-Calabi functional. We show that a Fano manifold admits Mabuchi's metric if and only if it is uniformly relatively D-stable. The idea of the proof includes some equivariant generalization of the recent developed variational approach to the Kähler-Einstein problem.

CONTENTS

1. Introduction	1
2. Mabuchi soliton and modified D-energy	4
3. Relative uniform D-stability	19
4. Variational approach and proof of the main theorem	29
References	33

1. INTRODUCTION

Let X be a Fano manifold. We denote the canonical line bundle by K_X and its dual by $-K_X$. In a central problem of complex geometry we are guided to look for a standard Kähler metric in the first Chern class $c_1(X) = c_1(-K_X)$. The fundamental result established in [CDS15], [Tia15] states that there exists a Kähler-Einstein metric if and only if X is K-polystable. Not all the Fano manifold satisfy the stability; for example one-point blow up of \mathbb{P}^2 is never Kähler-Einstein. On the other hand, for an *arbitrary* Fano manifold X we may consider a canonical geometric flow which should optimally destabilize X . The self-similar solution of the flow coincides with T. Mabuchi's generalization of Kähler-Einstein metric. The purpose of this paper is to clarify which Fano manifold admits such a metric.

For the definition, let us denote a Kähler metric by ω and the normalized Ricci potential function by ρ which is the unique function satisfies

$$\text{Ric } \omega - \omega = dd^c \rho, \quad \int_X (e^\rho - 1) \omega^n = 0. \quad (1.1)$$

We also write $\omega = dd^c \varphi$ locally so as to identify the metric with a collection of smooth functions φ patching together to define the fiber metric of $-K_X$. Our standard metric

Date: February 18, 2022.

first introduced by [M01] is the critical point of the Ricci-Calabi functional

$$R(\omega) = R(\varphi) := \frac{1}{V} \int_X (e^\rho - 1)^2 \omega^n. \quad (1.2)$$

Here the volume $V = \int_X \omega^n$ is independent of ω . The straightforward variational computation shows that the metric ω is a critical point iff $e^\rho - 1$ is the Hamilton function for some one-parameter subgroup $\eta: \mathbb{G}_m \rightarrow \text{Aut}(X, -K_X)$. It is also clear from the definition that the condition gives the Ricci-analogue of the extremal Kähler metric defined in terms of the classical Calabi functional. These two metrics are not the same while the above η is generated by the extremal vector field. There as well exists the infinite-dimensional GIT picture [D15] so that the Ricci-Calabi functional can be seen as the square norm of a certain moment map. The role of the Kemp-Ness functional in GIT is then played by the famous D-energy

$$D(\varphi) = -\log \frac{1}{V} \int_X e^{-\varphi} - \frac{1}{(n+1)V} \sum_{i=0}^n \int_X (\varphi - \varphi_0) \omega^i \wedge \omega_0^{n-i}.$$

The definition of the D-energy first appeared in [BM86] and was written down to this form by [D88]. The gradient flow

$$\frac{\partial}{\partial t} \varphi = 1 - e^\rho$$

was initially studied in our previous work [CHT22]. In the preprints [H19] and [X21] it was shown that the flow indeed minimizes $R(\varphi)$ and is naturally related with the optimal degeneration of the Fano manifold. From now on we call the pair of the critical point of $R(\varphi)$ and the one-parameter subgroup *Mabuchi soliton*, since it is characterized as the self-similar solution of the flow.

Our main result claims that the existence of Mabuchi soliton is equivalent to certain algebraic stability condition. It extends the result of [Y17], [N19] for the toric case to general Fano manifolds. Our approach precisely follows [BBJ21] where they give a new variational proof of [CDS15] for a Fano manifold with finite automorphism group. We in particular extended the argument to the general automorphism (with $\eta = 0$) case.

Theorem A. A Fano manifold X admits a Mabuchi soliton if and only if it is uniformly relatively D-stable, with respect to the equivariant test configurations.

If the extremal vector field is zero i.e. $\eta = 0$, the above statement essentially says that Kähler-Einstein metric if X is D-stable with respect to equivariant test configurations. This result was proved by [DS16] using so-called smooth continuity method. One advantage of the variational approach is that it can be comparatively easily applied to singular X (or more generally singular pair: X with a divisor) and for us, more than anything to the Mabuchi soliton case. The variational proof requires “uniform” version of the stability and to obtain the result we develop the equivariant formulation which was suggested in our previous work [H18]. One direction deriving stability is based on Theorem 3.20 which was proved in [H18]. In the equivariant setting, to obtain the metric one needs special discussion particularly in Lemma 3.19 and Lemma 4.3, to control the test configurations twisted by one-parameter subgroups. Also we need to be careful

for the equivariant formalism to confirm that the minimizer of the energy is a weak solution. This is concentrated in Theorem 2.13. The first version of our preprint had a flaw in the final step of the proof and this was pointed out and solved by [Li22]. This corresponds to the discussion around Lemma 4.3 and 4.4. It is not much, but we here serve another simple proof of Lemma 4.3.

The required stability condition is introduced in Definition 3.21. The concept of D-stability originates from [B16]. As the K-stability introduced by [D02] naturally arises from the Calabi functional and the K-energy, D-stability arises from the above Ricci-Calabi functional and the D-energy. Uniformity of the stability was introduced in our previous work [BHJ15] and [Der16a] independently. In regard to the torus containing the soliton vector field we may also formulate the relative version of the D-stability, especially based on [H16a] [H16b], and [H18]. Putting these together we formulate the *uniform relative stability* which reflects the coercivity of the modified D-energy.

One remarkable point is that we may restrict ourselves to test configurations equivariant for the *whole* automorphism group, in showing the existence of the metric. In our argument this fact owes to Theorem 2.13, where we used the fact that the Futaki invariant is trivial outside of the center.

Unlike K-stability, D-stability works only for Fano manifolds, however, as Theorem A and its proof show, the treatment is much easier. Existence of extremal Kähler metric is still open problem, even for the anti-canonical polarizations. A simple argument shows that the Mabuchi soliton assures the extremal Kähler metric. A new circumstance in the relative setting is that the two metrics are in fact not equivalent. The first counterexample is raised in the latest version of [NSY17].

Compared with Kähler-Ricci soliton, Mabuchi soliton has in some sense more algebraic nature. For example the soliton vector field is periodic and actually generates η . On the other hand, the gradient flow is not so flexible as the Kähler-Ricci flow. In addition toric examples in [NSY17] is in contrast to the result of [WZ04].

Along the variational approach we may naturally understand the uniqueness of Mabuchi soliton.

Theorem B ([M03], Theorem C). Let (ω_0, η_0) and (ω_1, η_1) be smooth Mabuchi solitons. Then there exists an automorphism $f \in \text{Aut}^0(X)$ in the identity component such that $f^*\omega_1 = \omega_0$, $f^*\eta_0 = \eta_1$.

Our argument also gives a new proof of the Matsushima-type theorem in [M03], [N19]. Namely, if a Fano manifold admits the Mabuchi soliton, the identity component of the group of automorphism preserving the extremal vector field is reductive. These uniqueness and the reductivity are indeed key materials for the derivation of the coercivity from existence of the metric.

Acknowledgment. The author express his gratitude to Professor C. Li, for his kind communication especially pointing out the flaws in our first version of the preprint. As we explained in the introduction, the first point is around Lemma 4.3 and 4.4. The second point is that the definition of the modified K-energy M_g was incorrect in the first version. See Remark 2.23 for the detail. The author wishes to thank Professor

R. Berman, S. Boucksom, and M. Jonsson for very fruitful discussions. Especially for the equivariant formulation he learned a lot from the three professors, as in our previous paper [H18]. We are also grateful to E. Inoue, S. Saito, R. Takahashi, and anonymous referee for helpful comments. This research was supported by JSPS KAKENHI Grant Number 15H06262 and 17K14185.

2. MABUCHI SOLITON AND MODIFIED D-ENERGY

2.1. Notation. Throughout the paper X denotes an n -dimensional Fano manifold and a Kähler metric ω is taken in the first Chern class $c_1(X)$. We adopt the additive notation writing the anti-canonical bundle as $-K_X$ and the fiber metric as φ . While we do not fix a specific covering $\{U_\alpha\}_\alpha$ of local coordinate patches U_α with index α , the symbol φ is interpreted to a function φ_α on each $U_\alpha \subset X$. In a local frame any section s of $-K_X$ is identified with a function s_α and it is evaluated by the multiplication of $e^{-\varphi_\alpha}$ to $|s_\alpha|^2$. On the intersection $U_\alpha \cap U_\beta$ for two indices α, β and coordinates z_α^i, z_β^j $1 \leq i, j \leq n$ the transition function is written to $g_{\alpha\beta} = \det \left[\frac{\partial z_\alpha^i}{\partial z_\beta^j} \right]_{ij}$ and the compatibility $\varphi_\alpha = \varphi_\beta + \log |g_{\alpha\beta}|^2$ holds. If we put $d^c = \frac{\partial - \bar{\partial}}{4\pi\sqrt{-1}}$, it follows that the Chern curvature $\omega_\varphi = dd^c\varphi$ is globally well-defined. We set $\mathcal{H}(X, -K_X)$ as the collection of smooth fiber metric φ on $-K_X$ such that ω_φ is positive. By dd^c -Lemma, any metric ω in $c_1(X)$ equals to ω_φ for some $\varphi \in \mathcal{H}(X, -K_X)$ which is unique up to addition of a constant. For this reason $\mathcal{H}(X, -K_X)$ is called *the space of Kähler metrics*. We essentially need φ instead of $\omega = \omega_\varphi$ in order to consider the action of the Hamilton diffeomorphism group.

2.2. Ricci curvature formulation. We briefly review some energy formulations to the Kähler-Einstein problem, which make use of the Ricci potential. There has been another (and probably major) scalar curvature formulation which works for a general polarized manifold. In terms of the scalar curvature one may introduce the Calabi functional and notion of K-stability observing the behavior of the K-energy along the degeneration of the manifold. See the milestone works [C82], [C85], [F83], [M86], [T97], and [D02]. D-energy which we will explain is as well classical but the determination of the corresponding D-stability [B16] and the momentum map picture [D15] were rather recent.

Let us start from defining two probability measures associated to a Kähler metric $\omega = \omega_\varphi$, or equivalently $\varphi \in \mathcal{H}(X, -K_X)$. One is the Monge-Ampère measure $V^{-1}\omega_\varphi^n$. The other one which we call the *canonical measure* is special for the Fano case and defined to be

$$\mu_\varphi := \frac{e^{-\varphi}}{\int_X e^{-\varphi}}, \quad (2.1)$$

where $e^{-\varphi}$ denotes the global volume form described as $e^{-\varphi_\alpha} \bigwedge_{i=1}^n dz_\alpha^i \wedge d\bar{z}_\alpha^i$ on a coordinate patch U_α . Note that the metric is Kähler-Einstein iff it satisfies the Monge-Ampère equation $V^{-1}\omega_\varphi^n = \mu_\varphi$. Therefore we are lead to focus on the difference of these two measures. In fact it precisely gives the infinite-dimensional moment map. Namely, once we regard a fixed Kähler metric ω as a symplectic form and instead collect all

the complex structures J compatible with ω , one may attach to each J the difference measure

$$J \mapsto \mu_\varphi - V^{-1}\omega^n. \quad (2.2)$$

The group of Hamilton diffeomorphisms naturally acts on the complex structures. The Lie algebra of this group is naturally identified with smooth function space $C^\infty(X; \mathbb{R})$ with Poisson bracket and hence the above defines a map to the dual Lie algebra. It indeed satisfies the moment map condition. More precisely, we should impose to J the compatibility condition with the fiber metric φ , but see [D15] for the detail explanation.

The square norm of the moment map is written down to

$$R(\varphi) = \frac{1}{V} \int_X (e^\rho - 1)^2 \omega^n, \quad (2.3)$$

which we call the Ricci-Calabi functional. Our interest is the critical point of the Ricci-Calabi functional which gives a generalization of the Kähler-Einstein metric. The first variation of $R: \mathcal{H}(X, -K_X) \rightarrow \mathbb{R}$ is given as follows. See also [N19] for calculating the second variation.

Proposition 2.1 ([CHT22], Proposition 2.3). *Set the twisted Laplacian on functions $f \in C^\infty(X, \mathbb{C})$ by*

$$L_\rho f = \Delta_\omega f + (\bar{\partial}\rho, \bar{\partial}f)_\omega. \quad (2.4)$$

Then the first variation of the Ricci-Calabi functional is given as

$$\delta R(\varphi) = -\frac{2}{V} \int_X \delta\varphi \left(L_\rho \tilde{f} + \tilde{f} \right) d\mu_\varphi, \quad (2.5)$$

where

$$\tilde{f} = (e^\rho - 1) - \frac{1}{V} \int_X (e^\rho - 1) d\mu_\varphi.$$

As a consequence, φ is the critical point of the Ricci-Calabi functional iff $e^\rho - 1$ is a Hamilton function. One can check this by a simple application of the Bochner-Kodaira formula. Since X is Fano any holomorphic vector field defines a function h unique up to addition of a constant such that

$$\sqrt{-1}\bar{\partial}h = i_v\omega. \quad (2.6)$$

We call h a Hamilton function.

Definition 2.2. *A Kähler metric $\omega \in c_1(X)$ is called a Mabuchi soliton if $e^\rho - 1$ is a Hamilton function for some holomorphic vector field.*

The vector field is zero iff $\rho = 0$ and in this case Mabuchi soliton is nothing but Kähler-Einstein.

Going back to the moment map picture, we also have the canonical energy functional $D: \mathcal{H}(X, -K_X) \rightarrow \mathbb{R}$ with the outer derivative $(dD)_\varphi = \mu_\varphi - V^{-1}\omega^n$ at φ . We call it D-energy. It is in fact separated into two terms $D = L - E$ and each term is specifically defined as

$$L(\varphi) := -\log \frac{1}{V} \int_X e^{-\varphi}, \quad E(\varphi) := \frac{1}{(n+1)V} \sum_{i=0}^n \int_X (\varphi - \varphi_0) \omega^i \wedge \omega_0^{n-i}. \quad (2.7)$$

We here take a reference φ_0 and $\omega_0 = dd^c\varphi_0$. Note that the difference $\varphi - \varphi_0$ defines a global function while φ does not. One can easily compute to check the differential

$$(dL)_\varphi = \mu_\varphi, \quad (dE)_\varphi = V^{-1}\omega^n. \quad (2.8)$$

The definition of the Monge-Ampère energy E chose φ_0 but it is characterized by (2.8), up to addition of a constant.

2.3. Space of finite energy metrics. A fundamental property of the D-energy is that it is convex along any geodesic in the space of Kähler metrics. Since the difference $\varphi - \psi$ of any two $\varphi, \psi \in \mathcal{H}(X, -K_X)$ defines a global function, tangent space at any point of $\mathcal{H}(X, -K_X)$ is identified with $C^\infty(X; \mathbb{R})$. Mabuchi's inner product [M87] for any tangents $u, v \in C^\infty(X; \mathbb{R})$ at φ is

$$\langle u, v \rangle = \frac{1}{V} \int_X uv\omega^n. \quad (2.9)$$

Any curve φ^t ($t \in [a, b]$) in $\mathcal{H}(X, -K_X)$ defines a function $\Phi(\tau, x) := \varphi^{-\log|\tau|}(x)$ of complex variables $e^{-b} \leq |\tau| \leq e^{-a}$ and $x \in X$. It is well-known (from [S92]) that the geodesity for (2.9) is equivalent to the degenerate Monge-Ampère equation

$$(dd_{\tau,x}^c \Phi)^{n+1} = 0. \quad (2.10)$$

The left-hand side at the same time describes the Monge-Ampère energy by the fiber integration formula

$$dd_\tau^c E(\varphi) = \int_X (dd_{\tau,x}^c \Phi)^{n+1}. \quad (2.11)$$

It follows that E is affine along any geodesics. In fact for given smooth endpoints the bounded *weak* geodesic Φ connecting them uniquely exists, but it is not C^2 in general.

Variational approach even requires the appropriate completion of the space of smooth metrics. These facts strongly motivate to consider a singular fiber metric φ which is only locally integrable and satisfies $dd^c\varphi \geq 0$ in the sense of current. We denote the collection of all such singular φ by $\text{PSH}(X, -K_X)$. It equivalent to say that in a coordinate patch U_α , φ_α is pluri-subharmonic (psh for short) function. For the bounded psh function the wedge product of the current $\omega_\varphi^n = (dd^c\varphi)^n$ is safely defined thanks to the celebrated work of [BT76]. In particular we may define the Monge-Ampère energy E for locally bounded φ . For a smooth boundary data we have the bounded solution of (2.10). From the recent result [CTW17], the solution is actually of $C^{1,1}$.

The Monge-Ampère operator $\varphi \mapsto V^{-1}\omega_\varphi^n$ can not be continuously extended to $\text{PSH}(X, -K_X)$. Following [BEGZ10] and [BBGZ13], one can however take the reference φ_0 smooth and bounded approximation $\varphi^{(j)} := \max\{\varphi, \varphi_0 - j\}$ of $\varphi \in \text{PSH}(X, -K_X)$, to define the *non-pluripolar Monge-Ampère measure*

$$\text{MA}(\varphi) := \lim_{j \rightarrow \infty} 1_{\{\varphi > \varphi_0 - j\}} V^{-1}\omega_{\varphi^{(j)}}. \quad (2.12)$$

By the construction $\text{MA}(\varphi)$ drops the mass of the unbounded locus so it is no longer a probability measure. It can be further shown that $\text{MA}(\varphi)$ is local in pluri-fine topology

and has no mass on any pluripolar set. In a similar idea taking bounded ψ such that $\psi \geq \varphi$ locally we define the Monge-Ampère energy as

$$E(\varphi) := \inf_{\psi} E(\psi) \in \mathbb{R} \cup \{-\infty\}. \quad (2.13)$$

The extended Monge-Ampère energy is upper-semicontinuous in the L^1 -topology of $\text{PSH}(X, -K_X)$. Moreover, the level set $\{E \geq C\}$ is compact in this weak topology, as a non-linear analogue of the Banach-Alaoglu theorem.

Let us now consider $p \geq 1$ and the L^p -Finsler distance d_p of $\mathcal{H}(X, -K_X)$, defined by the norm of tangents

$$\|u\|_p := \left[\frac{1}{V} \int_X |u|^p \omega_\varphi^n \right]^{\frac{1}{p}}. \quad (2.14)$$

As we shall see, $p = 1$ plays the special role in the variational approach.

Theorem 2.3 ([D15], Theorem 2. See also [BBJ21], Theorem 1.7). *Take a smooth non-increasing sequence of $\varphi_j \in \text{PSH}(X, -K_X)$ converges to φ . Endow the space of metrics with finite Monge-Ampère energy*

$$\mathcal{E}^1(X, -K_X) := \left\{ \varphi \in \text{PSH}(X, -K_X) : E(\varphi) > -\infty \right\} \quad (2.15)$$

with the distance

$$d_1(\varphi, \psi) := \lim_{j \rightarrow \infty} d_1(\varphi_j, \psi_j). \quad (2.16)$$

It then gives the coarsest refinement of the L^1 -topology so that E is continuous. The Monge-Ampère energy is affine along every geodesic on $(\mathcal{E}^1(X, -K_X), d_1)$. Moreover, $(\mathcal{E}^1(X, -K_X), d_1)$ realizes the completion of $(\mathcal{H}(X, -K_X), d_1)$.

We usually refer to the L^1 -topology as the “weak” topology and the d_1 -topology as the “strong” topology. The space $\mathcal{E}^1(X, -K_X)$ is contained in the *finite energy class*

$$\mathcal{E}(X, -K_X) := \left\{ \varphi \in \text{PSH}(X, -K_X) : \int_X \text{MA}(\varphi) = 1 \right\}. \quad (2.17)$$

Restricted to $\mathcal{E}(X, -K_X)$ the non-pluripolar Monge-Ampère operator is continuous along any monotone sequence. The determination of the domain of Monge-Ampère operator owes to the pioneering work [C98]. The compact setting is treated in [GZ07], [BEGZ10]. See also the comprehensive textbook [GZ17].

As it was shown in [D17a], geodesics connecting two points are not unique in \mathcal{E}^1 , however, for any $\varphi^0, \varphi^1 \in \mathcal{E}^1(X, -K_X)$ there exists the canonical one if we impose the plurisubharmonicity to the corresponding function Φ on the product space.

Theorem 2.4 ([BBJ21], Theorem 1.7). *For any $\varphi^0, \varphi^1 \in \mathcal{E}^1(X, -K_X)$ the solution of (2.10) has the unique solution $\varphi^t \in \mathcal{E}^1(X, -K_X)$ and provides a geodesic for d_1 . This special type of geodesic is called psh geodesic.*

The convexity of the D-energy functional along such a weak geodesic is established by the fundamental work [B09], [BP08], and [B15]. Since it is not scale free, i.e. $E(\varphi + c) = E(\varphi) + c$ for constants c , it is convenient to introduce the Aubin’s J-functional:

$$J(\varphi) = L_0(\varphi) - E(\varphi) := \sup_X (\varphi - \varphi_0) - E(\varphi). \quad (2.18)$$

It follows that $J(\varphi) - d_1(\varphi, \varphi_0)$ is uniformly bounded. As we have the uniform estimate

$$\sup_X(\varphi - \varphi_0) \leq \frac{1}{V} \int_X (\varphi - \varphi_0) \omega_0^n + C \quad (2.19)$$

for $\varphi \in \mathcal{H}(X, -K_X)$, sometimes $V^{-1} \int_X (\varphi - \varphi_0) \omega_0^n - E(\varphi)$ is adopted for the definition of J . Moreover one may introduce the symmetric I-functional

$$I(\varphi, \psi) := \int_X (\varphi - \psi)(\text{MA}(\psi) - \text{MA}(\varphi)) \quad (2.20)$$

which satisfies the pseudo-triangle inequality:

$$c_n I(\varphi_1, \varphi_2) \leq I(\varphi_1, \varphi_2) + I(\varphi_2, \varphi_3) \quad (2.21)$$

for any $\varphi_1, \varphi_2, \varphi_3 \in \mathcal{E}^1(X, \omega_0)$. The constant c_n only depends on the dimension of X . The induced topology of \mathcal{E}^1 is equivalent to d_1 -topology. See [BBEGZ19], section 1.

We say that the D-energy is coercive if

$$D(\varphi) \geq \varepsilon J(\varphi) - C$$

for any smooth φ . From the weak compactness of the level set $\{E \geq -C\}$ the coercivity guarantees the existence of a minimizer.

Remark 2.5. *If one considers coercivity for d_2 there is no example of Fano manifolds satisfy the condition. This is confirmed by [BHJ15], Proposition 8.5 for the K-energy. We may check the same for the D-energy, using Definition 3.2.*

2.4. Modified D-energy. Exploiting the inner product (2.9) we may also modify the D-energy such that the critical point gives the Mabuchi soliton.

It is consistent to consider the group of bundle automorphism $\text{Aut}(X, -K_X)$, indeed any $g \in \text{Aut}(X, -K_X)$ pulls-back $\varphi \in \text{PSH}(X, -K_X)$ to $g^*\varphi$. More precisely, for any $x \in X$ a vector $v \in (-K_X)_x$ is evaluated as

$$|v|^2 e^{-(g^*\varphi)(x)} = |g \cdot v|^2 e^{-\varphi(gx)}. \quad (2.22)$$

Note that the local frame identifying the function $\varphi(gx)$ with the pull-backed fiber metric depends on g . Indeed we see from (2.22) that the function $\varphi(gx)$ is unbounded in g . Since the line bundle is anti-canonical, any automorphism of X can be lifted to $-K_X$, hence the group splits into $\text{Aut}(X, -K_X) = \text{Aut}(X) \times \mathbb{G}_m$. We denote the identity component by $\text{Aut}^0(X, -K_X)$. In particular, constant multiplication on each fiber defines the identical one-parameter subgroup which we denote by $\kappa: \mathbb{G}_m \rightarrow \text{Aut}^0(X, -K_X)$.

Our first step is to specify the soliton vector field of the Mabuchi soliton. For the purpose we fix an algebraic subtorus $T \subset \text{Aut}(X, -K_X)$. The compact part $S = \text{Hom}(\mathbb{S}^1, T)$ is canonically defined. Henceforth we take a compact subgroup $K \subset \text{Aut}^0(X, -K_X)$ which contains S and commutes with T . It happens $K = S$. In any case we define the space of K -invariant metrics

$$\mathcal{H}(X, -K_X)^K := \left\{ \varphi \in \mathcal{H}(X, -K_X) : g^*\varphi = \varphi \text{ for any } g \in K \right\}. \quad (2.23)$$

From the assumption that K commutes with the torus, T acts on $\mathcal{H}(X, -K_X)^K$. Tangents are identified with smooth K -invariant functions. We take K -invariant functions in 2.14 to define the distance d_1 . Similarly the space of finite energy K -invariant metrics $\mathcal{E}^1(X, -K_X)^K$ can be defined and has the same property as Theorem 2.3. Note that in the definition

$$d_1(\varphi_0, \varphi_1) = \inf_{\varphi_t} \int_0^1 \frac{1}{V} \int_X |\dot{\varphi}_s| \omega_{\varphi_t}^n ds$$

a path φ_t , connecting φ_0 to φ_1 , is taken as K -invariant. The unique psh geodesic is however K -invariant if $\varphi_0, \varphi_1 \in \mathcal{H}(X, -K_X)^K$. It follows that d_1 for $\mathcal{H}(X, -K_X)^K$ equals to the previous one for $\mathcal{H}(X, -K_X)$.

Let us denote by $N := \text{Hom}(\mathbb{G}_m, T)$ the lattice of all one-parameter subgroups $\mu: \mathbb{G}_m \rightarrow T$. The dual lattice $M := \text{Hom}(T, \mathbb{G}_m)$ is identified with the set of characters. Observe that the vector space $N_{\mathbb{R}} := N \otimes \mathbb{R}$ is identified with the Lie algebra \mathfrak{s} of S . From the basic symplectic geometry S defines the moment polytope $P \subset M_{\mathbb{R}}$ as the image of the moment map

$$m_\varphi: X \rightarrow M_{\mathbb{R}}. \quad (2.24)$$

Actually for any smooth K -invariant ω_φ and $\mu \in N_{\mathbb{R}}$ we have the unique map satisfying

$$\langle \mu, m_\varphi(x) \rangle = \left. \frac{d}{dt} \right|_{t=0} \varphi(\mu(e^t)x). \quad (2.25)$$

It is easy to show that m_φ is independent of the metric. Once $\mu \in N_{\mathbb{R}}$ is fixed $h_\mu := \langle \mu, m_\varphi(x) \rangle$ gives the (unnormalized) Hamilton function. Notice that when $\mu \in N$ is generated by a vector field $v \in \mathfrak{t}$ we have the relation (2.6). The S -invariance of ω guarantees that h_μ is real. For the identical one-parameter subgroup we observe $h_\kappa \equiv 1$.

In this convention following [FM95] we introduce the inner product

$$\langle \mu, \nu \rangle := \int_X h_\mu h_\nu \omega^n \quad (2.26)$$

for $\mu, \nu \in N_{\mathbb{R}}$. Of course h_μ depends on the choice of metric ω but as we will see in the next section the above inner product is determined only by μ, ν . The Hamilton function can be regarded as the tangent vector of the associated (smooth) geodesic ray

$$\varphi^t = \mu(e^{-t})^* \varphi^0 \quad (2.27)$$

for a given initial $\varphi^0 \in \mathcal{H}(X, -K_X)^K$. Note again that φ^t is K -invariant for each t since we assumed K commutes with T .

The slope of D-energy along this ray is independent of $t \in [0, \infty)$ and explicitly computed as

$$F(\mu) := \frac{1}{V} \int_X h_\mu (e^\rho - 1) \omega^n. \quad (2.28)$$

It is precisely the classical Futaki invariant [F83] for the vector field generating μ . Notice that using the scalar curvature S_{ω_φ} and its average \hat{S} the Futaki invariant can be also written as

$$F(\mu) = -\frac{1}{V} \int_X h_\mu (S_{\omega_\varphi} - \hat{S}) \omega^n \quad (2.29)$$

and the description leads us to the definition of the K-energy. Therefore, D-energy is rather natural in view of the original description in [F83]. The extremal vector field naturally arises from the optimization of $F(\mu)$ normalized by $\|\mu\| = \langle \mu, \mu \rangle^{\frac{1}{2}}$. Actually a simple variational computation

$$\delta \left(\frac{F(\mu)}{\|\mu\|} \right) = \frac{1}{\|\mu\|} \left(F(\delta\mu) - \frac{\langle \delta\mu, \mu \rangle}{\langle \mu, \mu \rangle} F(\mu) \right)$$

suggest us to introduce the extremal one-parameter subgroup $\eta \in N_{\mathbb{R}}$, which satisfies

$$F(\mu) - \langle \mu, \eta \rangle = 0 \quad (2.30)$$

for any $\mu \in N_{\mathbb{R}}$. The extremal η in particular satisfies $F(\eta) - \langle \eta, \eta \rangle = 0$ hence it is a critical point. On the other hand, if we take a critical point η with $F(\eta) > 0$ (regarding the symmetry $F(-\mu) = -F(\mu)$), we may assume $F(\eta) = \|\eta\|^2$ because the positive constant multiplication $\eta \mapsto c\eta$ preserves $F(\mu)/\|\mu\|$. Then the first variation formula implies that η is extremal. Since (2.30) is a system of linear equations, one can easily see that η is uniquely characterized by the above relation. It is also easy to check $\eta \in N_{\mathbb{Q}}$ and automatically

$$\langle \kappa, \eta \rangle = \int_X h_\eta \omega^n = 0. \quad (2.31)$$

On the other hand the Mabuchi soliton should minimize $R(\varphi)$. In fact if there exists a Mabuchi soliton ω_φ with $e^\rho - 1 = h_\mu$ for some $\mu \in N_{\mathbb{R}}$ we have $\mu = \eta$ and

$$R(\varphi) = \frac{F(\eta)}{\|\eta\|}.$$

That is, both the optimizers φ and η attain the same value. In general the lower bound of the Ricci-Calabi functional is attained by the normalized non-Archimedean D-energies which we introduce in the next section. See the recent work [X21] and [H19] for this topic. The following is a consequence of Theorem 2.9 in the next subsection.

Proposition 2.6. *There exists the modified Monge-Ampère energy $E_\eta: \mathcal{H}(X, -K_X)^K \rightarrow \mathbb{R}$ satisfying*

$$(dE_\eta)_\varphi = (1 + h_\eta)V^{-1}\omega_\varphi^n$$

at each point $\varphi \in \mathcal{H}(X, -K_X)^K$. Moreover, E_η is geodesically affine.

We define modified D-energy as $D_\eta := L - E_\eta$. It follows from the proposition that a smooth metric ω is Mabuchi soliton iff it is a critical point of the modified D-energy.

2.5. Modified Monge-Ampère measure. For the variational approach it is necessary to handle with $E_\eta(\varphi)$ for singular φ . In this part following [BWN14] we discuss basic properties of the modified Monge-Ampère measure. In [BWN14] the case $K = S$ is considered but the same argument works for general K which contains S and commutes with T . Let continuously $m_\varphi: X \rightarrow P$ be the moment map.

Definition 2.7. *Let $\varphi \in \mathcal{H}(X, -K_X)^K$. For a non-negative continuous function $g: P \rightarrow \mathbb{R}$ define the modified Monge-Ampère measure*

$$\text{MA}_g(\varphi) := g(m_\varphi(x)) \text{MA}(\varphi).$$

The definition further extends to general $\varphi \in \text{PSH}(X, -K_X)^K$ so that the measure $\text{MA}_g(\varphi)$ is local in plurifine topology and non-pluripolar.

Theorem 2.8 ([BWN14], Theorem 2.7). *The Duistermaat-Heckman measure*

$$\text{DH}_T := (m_\varphi)_* \text{MA}(\varphi)$$

is independent of smooth φ and defines a positive measure on $M_{\mathbb{R}}$. For any $\varphi \in \text{PSH}(X, -K_X)^K$ we have

$$\int_X \text{MA}_g(\varphi) \leq \int_P g \text{DH}_T.$$

The equality holds if $\varphi \in \mathcal{E}(X, -K_X)^K$, namely when $\text{MA}(\varphi)$ is a probability measure.

Let \langle, \rangle be the canonical pairing of the lattices N and M . We are interested in the case

$$g(x) := 1 + \langle \eta, x \rangle - \int_P \langle \eta, x \rangle \text{DH}_T. \quad (2.32)$$

Note that g of this form is not necessarily non-negative. At least when $g \geq 0$ and φ smooth we observe $\text{MA}_g(\varphi) = (1 + h_\eta) \text{MA}(\varphi)$ and

$$(\inf_P g) \text{MA}(\varphi) \leq \text{MA}_g(\varphi) \leq (\sup_P g) \text{MA}(\varphi). \quad (2.33)$$

Notice that $1 + h_\eta > 0$ holds if X admits a Mabuchi soliton. As in the next section we shall see that the condition $1 + h_\eta > 0$ is numerical, from now on we assume that the above g is positive. Then the equation of Mabuchi soliton may be interpreted into the Monge-Ampère type equation

$$\text{MA}_g(\varphi) = \mu_\varphi. \quad (2.34)$$

We call $\varphi \in \text{PSH}(X, -K_X)^K$ satisfying this condition a *weak Mabuchi soliton*.

On the other hand, if we choose

$$g(x) = \frac{e^{\langle \mu, x \rangle}}{\int_P e^{\langle \mu, x \rangle} \text{DH}_T} \quad (2.35)$$

with certain μ , equation (2.34) gives the weak Kähler-Ricci soliton. In this case g is always positive but $\mu \notin N_{\mathbb{Q}}$.

Theorem 2.9 ([BWN14], Lemma 2.14, Proposition 2.15). *We have the canonical energy $E_g: \mathcal{H}(X, -K_X)^K \rightarrow \mathbb{R}$ such that $(dE_g)_\varphi = \text{MA}_g(\varphi)$. For general $\varphi \in \text{PSH}(X, -K_X)^K$ we have*

$$E_g(\varphi) := \inf_{\psi \geq \varphi} E_g(\psi),$$

where ψ runs through bounded ones, or $\mathcal{H}(X, -K_X)^K$. The functional E_g is monotone, upper-semicontinuous in L^1 -topology, and continuous for any non-increasing sequence in $\text{PSH}(X, -K_X)^K$.

We would write $E_\eta := E_g$ in the case (2.32). The description of E_g is easily specified so we briefly sketch it. For the path $\varphi_t = (1-t)\varphi + t\varphi_0$ the required E_g is computed as

$$\begin{aligned} E_g(\varphi) &= \int_0^1 \frac{d}{dt} E_g(\varphi_t) dt = \frac{1}{V} \int_0^1 dt \int_X (\varphi - \varphi_0) g(m_{\varphi_t}) \omega_{\varphi_t}^n \\ &= \frac{1}{V} \sum_{i=0}^n \binom{n}{i} \int_0^1 t^i (1-t)^{n-i} dt \int_X (\varphi - \varphi_0) g(m_{\varphi_t}) \omega_\varphi^i \wedge \omega_0^{n-i}. \end{aligned}$$

Note $m_{\varphi_t} = tm_\varphi + (1-t)m_{\varphi_0}$ and that the last integrand is just a variant of modified Monge-Ampère measure. Therefore we may exploit Definition 2.7 to derive the required property of E_g . If $\inf_P g$ is positive it follows

$$(\sup_P g)E(\varphi) \leq E_g(\varphi) \leq (\inf_P g)E(\varphi) \quad (2.36)$$

provided $\sup_X(\varphi - \varphi_0) = 0$. It implies that $E_g(\varphi) > -\infty$ if φ has finite Monge-Ampère energy. At any case we define the g -modified J-energy by

$$J_g(\varphi) := L_0(\varphi) - E_g(\varphi). \quad (2.37)$$

Let again $J_\eta := J_g$ in the case (2.32). This is after all equivalent to the ordinal J -functional.

Lemma 2.10. *When $g > 0$, we have*

$$(\inf_P g)J_g(\varphi) \leq J(\varphi) \leq (\sup_P g)J_g(\varphi)$$

for all $\varphi \in \mathcal{H}(X, -K_X)^K$.

Proof. Since $J_g(\varphi+c) = J_g(\varphi)$ for any constant $c \in \mathbb{R}$, we may assume $\sup_X(\varphi - \varphi_0) = 0$. The claim is then a consequence of (2.36). \square

For a given probability measure μ one can consider the Monge-Ampère type equation $\text{MA}_g(\varphi) = \mu$. It was also shown in [BWN14] Theorem 2.18 that there exists the unique solution $\varphi \in \mathcal{E}^1(X, -K_X)^K$ iff the Legendre dual of the Monge-Ampère energy

$$E_g^*(\mu) := \sup_{\varphi \in \mathcal{E}^1(X, -K_X)} \left[E_g(\varphi) - \int_X (\varphi - \varphi_0) d\mu \right] \in \mathbb{R} \cup \{\infty\} \quad (2.38)$$

is finite. Moreover, the above supremum is attained by the solution. We denote the dual of E_η by E_η^* .

What we will study is the g -modified D-energy $D_g(\varphi) := L(\varphi) - E_g(\varphi)$ and the equation (2.34). The convexity of L -functional follows from the main result of [B15], while a direct computation shows the fiber integration formula:

$$dd_\tau^c E_g(\varphi^t) = \frac{1}{(n+1)V} \int_X g(m_\Phi) (dd_{x,\tau}^c \Phi(x, \tau))^{n+1}, \quad (2.39)$$

which generalizes (2.11). We observe that E_g is affine along any psh geodesic. Moreover, if $g > 0$, the affineness of $E_g(\varphi^t)$ and $E(\varphi^t)$ are equivalent for any ray φ^t on $\mathcal{E}(X, \omega_0)$. In particular the affineness of $E_g(\varphi^t)$ implies that φ^t is a psh geodesic. As a consequence we obtain the convexity of the modified D -energy.

Theorem 2.11 ([B15], [BWN14]). *The g -modified D -energy $D_g(\varphi) := L(\varphi) - E_g(\varphi)$ is convex along any weak geodesic φ^t ($t \in [a, b]$) which is a bounded solution of (2.10).*

We will prepare the following lemma which can be seen as a variant of Theorem 2.11.

Lemma 2.12. *Let $D_\eta: \mathcal{E}^1(X, \omega_0)^K \rightarrow \mathbb{R}$ be the modified D -energy. For each $\varphi \in \mathcal{E}^1(X, -K_X)^K$, the map $\mathbf{d}_\eta: K_{\mathbb{C}} \rightarrow \mathbb{C}$ defined by $\mathbf{d}_\eta(g) := D_\eta(g^*\varphi)$ is pluriharmonic. In particular if D_η is bounded from below, then \mathbf{d}_η is constant on the center.*

Proof. This is similar to [H18], Theorem 1.6 and Remark 2.6 which are for the case $\eta = 0$. The statement simply interprets geodesically affineness of D_η into the complex variables.

The log part is obviously pluriharmonic. We show that $\mathbf{e}_\eta(g) := E_\eta(g^*\varphi)$ is pluriharmonic. Let us take an arbitrary holomorphic map $g: \Delta \rightarrow \text{Aut}^0(X, L)$ which sends $z \in \Delta$ in the one dimensional disk to the automorphism $g(z)$. The formula 2.39 is translated to

$$dd^c E_\eta(\varphi_{g(z)}) = \frac{1}{(n+1)V} \int_X (1 + h_{\varphi, \eta})(dd_{z,x}^c \varphi_{g(z)}(x))^{n+1}.$$

For the holomorphic map $F: \Delta \times X \rightarrow X$ by $F(z, x) := g(z) \cdot x$ we have

$$(dd_{z,x}^c \varphi_{g(z)}(x))^{n+1} = (dd_{z,x}^c F^* \varphi)^{n+1} = F^*(dd_x^c \varphi)^{n+1} = 0.$$

It implies that \mathbf{e} is pluriharmonic. □

Using the above lemma let us show that the weak minimizer of the energy is actually the Mabuchi soliton. This forms one of the critical step in our proof of Theorem A.

Theorem 2.13. *Assume that T contains the center of the complexified Lie group $K_{\mathbb{C}}$. If there exists a minimizer of modified D -energy $D_\eta: \mathcal{E}^1(X, \omega_0)^K \rightarrow \mathbb{R}$ it defines a weak solution of (2.34).*

Proof. For a function v we define the point-wise upper envelope

$$Pv := \sup \left\{ \psi \in \text{PSH}(X, -K_X)^K, \psi \leq v \right\}.$$

The proof is due to the highly non-trivial derivation formula ([BWN14], Proposition 2.16):

$$\left. \frac{d}{dt} \right|_{t=0} E_g(P(\varphi + tu)) = \int_X u \text{MA}_g(\varphi)$$

for $\varphi \in \mathcal{E}^1(X, -K_X)^K$, $u \in C^0(X; \mathbb{R})^K$. This was first established in [BB10] for $g = 1$, $K = \{\text{id}\}$ case. If φ is a minimizer of D_η , we observe

$$\begin{aligned} f(t) &:= L(\varphi + tu) - E_\eta(P(\varphi + tu)) \\ &\geq L(P(\varphi + tu)) - E_\eta(P(\varphi + tu)) \\ &\geq L(\varphi) - E_\eta(\varphi) = f(0). \end{aligned}$$

The derivation formula yields $f'(0) = 0$ and hence

$$\int_X u \text{MA}(\varphi) = \int_X u \mu_\varphi. \tag{2.40}$$

for every $u \in C^0(X; \mathbb{R})^K$.

We should show that the same holds for any $u \in C^0(X; \mathbb{R})$. By Lemma 2.12, \mathbf{d}_η is constant on the center. We observe that for any one-parameter subgroup $\mu \in N$ the slope of $\mathbf{d}(\mu(e^{-t}))$ is equivalent to the classical Futaki character. Since the character is defined on the reductive Lie algebra $\mathfrak{k}_\mathbb{C}$ which can be written as the direct sum of the center and the derived algebra, the slopes are nontrivial only on the center. Therefore \mathbf{d}_η is actually constant on whole $K_\mathbb{C}$. Thus the measure $\mu := (dD)_\varphi$ is K -invariant. It then follows that for any smooth function v and $g \in K$

$$\int_X v \mu = \int_X g_*(v \mu) = \int_X ((g^{-1})^* v) g_*(\mu) = \int_X ((g^{-1})^* v) \mu.$$

Integrating against the Haar measure we have

$$\int_X v \mu = \int_X u \mu = 0$$

so that $\mu = 0$ as desired.

Conversely, if $\varphi \in \mathcal{E}^1(X, -K_X)^K$ is a weak solution, convexity of D_g implies that φ is a minimizer. \square

Remark 2.14. *More generally for arbitrary smooth $(1, 1)$ form α one can consider the energy $E_\alpha: \mathcal{H} \rightarrow \mathbb{R}$ with the property*

$$(dE_\alpha)_\varphi = V^{-1} \alpha \wedge (dd^c \varphi)^{n-1}.$$

The g -modified energy $E_{g,\alpha}$ is similarly defined and extended to $\mathcal{E}^1(X, -K_X)$. The particular case $\alpha = \text{Ric}(\omega_0)$ for some fixed metric is denoted by $E_{g,\text{Ric}}$. This is called Ricci energy and plays a role in the study of K -energy. See section 2.7.

In the next subsection a refinement of the latter argument will show the uniqueness of the weak Mabuchi soliton.

Let us discuss about the coercivity of the modified D-energy. From now on we rather start from the extremal one-parameter subgroup η . Let

$$\text{Aut}(X, \eta) := \left\{ g \in \text{Aut}(X, -K_X) : \eta(\tau)g = g\eta(\tau) \text{ for all } \tau \in \mathbb{G}_m. \right\} \quad (2.41)$$

The identity component is denoted by $\text{Aut}^0(X, \eta)$. We take afresh $T = C(\text{Aut}^0(X, \eta))$ as the center of the automorphisms commuting with η . Moreover, we entirely consider a maximal compact subgroup K containing S . It clearly commutes with the center. We set

$$J_T(\varphi) := \inf_{\sigma \in T} J(\sigma^* \varphi). \quad (2.42)$$

Definition 2.15. *Let $T = C(\text{Aut}^0(X, \eta))$ and K be a maximal compact subgroup of $\text{Aut}^0(X, \eta)$, which contains the compact part of T . We say that the modified D-energy is coercive if there exists a positive constants ε, C such that*

$$D_\eta(\varphi) \geq \varepsilon J_T(\varphi) - C$$

holds for every invariant metric $\varphi \in \mathcal{H}(X, -K_X)^K$.

By the standard argument we may obtain the weak minimizer from the coercivity. Actually for a minimizing sequence φ_j , we have $\sigma_j \in T$ by the coercivity such that $\sigma_j^* \varphi_j$ is contained in the sublevel set $\{J_\eta \leq C\}$. Since $\{E \geq -C\}$ is weakly compact, we obtain a weakly convergent subsequence $\sigma_j^* \varphi_j \rightarrow \varphi$ in $\mathcal{E}^1(X, -K_X)^K$. From Lemma 2.12 the map $\sigma \mapsto D_\eta(\sigma^* \varphi_j)$ is constant. That is, D_η must be T -invariant. Especially $D_\eta(\sigma_j^* \varphi_j) = D_\eta(\varphi_j)$. The lower-semicontinuity concludes that φ is a minimizer of D_η . Theorem 2.13 concludes that the obtained minimizer is a weak Mabuchi soliton. It is indeed a smooth Mabuchi soliton, by Theorem 2.17. Let us define Mabuchi's numerical constant as

$$m_X = \inf_X(1 + h_\eta), \quad (2.43)$$

where η is the extremal one-parameter subgroup.

Our goal in this subsection is:

Theorem 2.16 ([LZ17]). *A Fano manifold X admits a Mabuchi soliton if and only if $m_X > 0$, $\text{Aut}^0(X, \eta)$ is reductive, and the modified D -energy is coercive.*

We have already explained that the coercivity implies the existence of the metric. The converse direction is based on [DR17]. For the Mabuchi soliton [LZ17] already obtained a result adopting a different definition of the coercivity. Since the difference of the formulation is subtle point for the equivariant setting, we briefly sketch the proof in our framework.

Proof. Let φ be the Mabuchi soliton. Trivially $K \subset \text{Aut}(M, \varphi)$ so the maximality implies $K = \text{Aut}(M, \varphi)$. Let $G = \text{Aut}^0(X, \eta)$. By Corollary 2.21 $G = K_{\mathbb{C}}$ is reductive. We consider the normalizer and the centralizer

$$\begin{aligned} N_K(G) &:= \{g \in G : gkg^{-1} \subset K\}, \\ C_K(G) &:= \{g \in G : gkg^{-1} = k \text{ for every } k \in K\}. \end{aligned}$$

We first observe $C_K(G) = C(G)$. Indeed any $t \in C_K(G)$ we have the map $\tau : G \rightarrow G$ defined by $\tau(g) = tgt^{-1}$ and this is identical for $g \in K$. By Corollary 2.21 it implies that τ is identical on G .

From the general theory of Lie groups we know that $N_K(G)/KC_K(G)$ is finite. Let us show $N_K(G) = KC(G)$ in our situation. Since $N_K(G)/KC_K(G)$ is finite, we may write $N_K(G) = K'C_K(G) = K'C(G)$ for some maximal compact subgroup K' . By construction $K \subset K'$ so the maximality of K implies $N_K(G) = KC(G)$.

We now check that $T = C(G)$ acts transitively on the smooth Mabuchi solitons. By Theorem 2.20, for two Mabuchi solitons φ and φ' we have $f \in \text{Aut}^0(X, \eta)$ such that $f^* \varphi = \varphi'$. Since $K = \text{Aut}(M, \varphi) = \text{Aut}(M, \varphi')$ as we have already observed, it follows $f^{-1}Kf \subset K$. Namely, $f \in N_K(G) = KC(G)$.

The above transitivity of T and the regularity of weak minimizers (Theorem 2.17), we may apply [DR17], Theorem 3.4 (with $\mathcal{R} = \mathcal{H}(X, -K_X)^K$, $G = C(\text{Aut}^0(X, \eta))$ there) so that have constants ε, C and

$$D_\eta(\varphi) \geq \varepsilon \inf_{\sigma \in T} J(\sigma^* \varphi) - C \quad (2.44)$$

for every $\varphi \in \mathcal{H}(X, -K_X)^K$. □

2.6. Uniqueness of Mabuchi soliton. We shall first check the regularity. In [LZ17] the corresponding step is carried out by the continuity method assuming the coercivity. We here introduce a direct argument.

Theorem 2.17. *Assume $m_X = \inf_X(1+h_\eta)$ is strictly positive. Then the weak Mabuchi soliton of (2.34) is actually smooth.*

Proof. Since φ has finite Monge-Ampère energy it has zero Lelong number (see [GZ17], Exercise 10.7). By the uniform version of Skoda's integrability theorem ([GZ17], Theorem 8.11), μ_φ has L^p -density for any $p > 1$. Noting (2.33) and applying the viscosity theory: [EGZ11], Theorem C to $\text{MA}_g(\varphi) = \mu_\varphi$, we deduce that φ is continuous. We may further show φ is C^∞ essentially using Yau's C^2 -estimate. For example one can apply the idea of [ST09] Theorem 1 to the present setting. See [ST19] for the detail exposition. \square

For the uniqueness the fact $\varphi \in L^\infty$ is important, because we need the following.

Proposition 2.18 ([B15], Theorem 1.2). *Let φ^t be a weak geodesic which is uniformly bounded in the sense that $|\varphi^t - \varphi_0| \leq C$. If the convex function $L(\varphi^t)$ is affine, there exists a $f_t \in \text{Aut}(X, -K_X)$ such that*

$$f_t^* \omega_{\varphi^t} = \omega_{\varphi^0}.$$

Moreover $f_t = \exp(-t \text{Re } v)$ for some holomorphic vector v lifted to $-K_X$ such that $\text{Im } v$ preserves ω_{φ^t} .

Remark 2.19. *By [B16], Proposition 3.3, we may further conclude $f_t^* \varphi^t = \varphi^0$.*

For $\mu \in N_{\mathbb{R}}$ we denote by $\text{Aut}(X, \mu)$ the group of bundle automorphisms preserving μ . Set $\text{Aut}(X, \varphi)$ for a fiber metric φ in a similar manner.

Theorem 2.20. *Let (ω_0, η_0) and (ω_1, η_1) be smooth Mabuchi solitons. Then there exists some $f \in \text{Aut}^0(X, -K_X)$ such that*

$$f^* \omega_1 = \omega_0, \quad f^* \eta_1 = \eta_0.$$

If $\eta_0 = \eta_1$ we have $f \in \text{Aut}^0(X, \eta_1)$ and one can further take f generated by the imaginary part of $\text{Aut}(X, \varphi_1)_{\mathbb{C}}$.

Proof. First we consider the case $\eta = \eta_0 = \eta_1$ contained in the torus T . Take potentials φ^0, φ^1 of ω_0, ω_1 and bounded geodesic φ^t ($t \in [0, 1]$). Since φ^0, φ^1 are minimizers the convex function $D_\eta(\varphi^t)$ should be affine. In particular $L(\varphi^t)$ is affine. We may apply Theorem 2.18 so that $f_t^* \varphi^t = \varphi^0$. Observe that $\varphi^t = (f_t^{-1})^* \varphi^0$ is a weak Mabuchi soliton, since it is a minimizer of D_η . If we take the extremal vector field v generating η and set $w := (f_t)_* v - v$, it follows $L_w \omega_0 = 0$ and hence $dd^c h_w = 0$. That is, f_t preserves η .

When $\eta_0 \neq \eta_1$ noting that the maximal tori are conjugate to each other we may take some f so that $\eta_1 = f^* \eta_0$, by the uniqueness of the extremal vector field. \square

The uniqueness argument is closely related to the reductivity result.

Corollary 2.21. *If a Fano manifold X admits a Mabuchi soliton (ω_φ, η) we have*

$$\text{Aut}^0(X, \eta) = \text{Aut}(X, \varphi)_\mathbb{C}.$$

That is, $\text{Aut}^0(X, \eta)$ is a complexification of the compact Lie group $\text{Aut}(X, \varphi)$.

Proof. From $h_\eta = e^\rho - 1$ we know $\text{Aut}(X, \varphi) \subset \text{Aut}^0(X, \eta)$. If we take $g \in \text{Aut}^0(X, \eta)$, $g^*\varphi$ is Mabuchi soliton hence some $f \in \text{Aut}(X, \varphi)_\mathbb{C}$ satisfies $f^*\varphi = g^*\varphi$. It follows $g = (g \circ f^{-1}) \circ f \in \text{Aut}^0(X, \varphi)_\mathbb{C}$. \square

Mabuchi first showed Theorem 2.20 using the inverse-continuity method of [BM85]. In [M03] Corollary 2.21 is also proved by the twisted Laplacian calculus similarly to [M57]. As it was shown in [N19], one can also derive Corollary 2.21 directly from the second variation of the Ricci-Calabi functional. The present proofs are based on the idea of [B15] for the Kähler-Einstein metric. A virtue of this idea more directly links reductivity to the uniqueness.

2.7. Thermodynamical formalism and modified K-energy. In this final part of the section, following the thermodynamical formalism of [B13] and its modified version in [BWN14], we introduce the modified K-energy in terms of D-energy.

Recall for two probability measures μ, ν the relative entropy is defined to be

$$H(\mu|\nu) = \int_X \log \left[\frac{d\mu}{d\nu} \right] d\mu. \quad (2.45)$$

Its relation with D-energy is based on the Legendre transformation formula:

$$H(\mu|\mu_0) = \sup_{f \in C^0(X; \mathbb{R})} \left[\int_X f d\mu - \log \int_X e^f d\mu_0 \right]. \quad (2.46)$$

Definition 2.22. *Fix a reference $\varphi_0 \in \mathcal{H}(X, -K_X)^K$ and $\mu_0 := \mu_{\varphi_0}$. Let $g: P \rightarrow \mathbb{R}$ be a positive continuous function on the moment polytope. For μ with finite $E^*(\mu)$ we define the free energy*

$$F(\mu) := H(\mu|\mu_0) - E^*(\mu). \quad (2.47)$$

For $\varphi \in \mathcal{E}^1(X, -K_X)^K$ we define the K-energy as $M(\varphi) := F(\text{MA}(\varphi))$ and the modified K-energy as

$$M_g(\varphi) := F(\text{MA}_g(\varphi)) = H(\text{MA}_g(\varphi)|\mu_0) - E_g(\varphi) + \int_X (\varphi - \varphi_0) \text{MA}_g(\varphi). \quad (2.48)$$

In [BWN14] the Kähler-Ricci soliton case (2.35) was discussed. In this case M_g is equivalent to the energy introduced by [TZ02] for smooth metrics. The treatment is valid for arbitrary g including Mabuchi soliton case. From the definition it follows

$$\begin{aligned} F(\mu) &= H(\mu|\mu_0) - E^*(\mu) \\ &= \sup_f \left[\int_X f d\mu - \log \int_X e^f d\mu_0 \right] - \sup_\varphi \left[E(\varphi) - \int_X (\varphi - \varphi_0) d\mu \right] \end{aligned}$$

and the second supremum is attained by the weak solution of $\text{MA}(\varphi) = \mu$. Therefore we obtain $M_1 = M$, which is analogues to the Chen-Tian formula ([C00]). In particular the $g = 1$ case gives the original definition of K-energy.

Remark 2.23. (1) *In the first version of our preprint we adopted different (and fail to work) definition of M_g . This is mainly because the convexity property of $F(\text{MA}_g(\varphi))$ was unclear. This was confirmed by [HL20]. We give a brief exposition in the below.*

- (2) *There are already several functionals called modified K-energy in the literatures, which are defined mainly to characterize the extremal Kähler metric. For example the functional $M' := M - E_\eta$ gives one such candidate. An extremal metric might not be a Mabuchi soliton unless it is Kähler-Einstein. Since $M \geq D$ clearly implies $M' \geq D_\eta$, if D_η is coercive so does M' . It follows that if X admits a Mabuchi soliton it also has an extremal Kähler metric. The converse does not holds. For example, $X = \mathbb{P}(\mathcal{O}_{\mathbb{P}^2} \oplus \mathcal{O}_{\mathbb{P}^2}(2))$ admits an extremal Kähler metric but not Mabuchi soliton. See [NSY17] for the detail.*

First of all, we have the following convexity property of M_g .

Theorem 2.24 (The special case of [HL20], Section 4). *Assume $g > 0$. The modified K-energy M_g is convex along any psh geodesic in $\mathcal{E}^1(X, -K_X)^K$.*

Proof. The result for $g = 1$ was first proved by [BB17], assuming that the psh geodesic has bounded Laplacian. It was extended to arbitrary psh geodesic in \mathcal{E}^1 by [BDL17]. [HL20] treated much more general situation using parallel but more complicated computations. We here give a brief account for the present case. First we assume φ^t is smooth and rewrite (2.48) in the form:

$$M_g(\varphi) = \int_X \log \left[\frac{\omega_\varphi^n}{e^{-(\varphi-\varphi_0)} d\mu_0} \right] \text{MA}_g(\varphi) + \int_X \log [g(m_\varphi)] \text{MA}_g(\varphi) - E_g(\varphi). \quad (2.49)$$

The second term in the right-hand side is constant in t . We have already observed that E_g is geodesically affine. For the first term we compute the fiber integral along the projection $(z, \tau) \mapsto \tau$. Since m_φ is a moment map there exists a (not necessarily holomorphic but vertical) vector field V_g such that

$$\sqrt{-1} \bar{\partial}_{z,\tau} g(m_\varphi) = i_{V_g} dd_{z,\tau}^c \Phi \quad (2.50)$$

holds. Consequently

$$\begin{aligned} dd_\tau^c \int_X \log \left[\frac{(dd_z^c \varphi)^n}{e^{-(\varphi-\varphi_0)} d\mu_0} \right] g(m_\varphi) (dd_{z,\tau}^c \Phi)^n &= \int_X g(m_\varphi) dd^c \log \left[\frac{(dd_z^c \varphi)^n}{e^{-(\varphi-\varphi_0)} d\mu_0} \right] \wedge (dd_{z,\tau}^c \Phi)^n \\ &= \int_X g(m_\varphi) dd^c \log \left[\frac{(dd_z^c \varphi)^n}{(dd_z^c \varphi_0)^n} \right] \wedge (dd_{z,\tau}^c \Phi)^n. \end{aligned}$$

In the second equality we used the fact that $E_{g,\text{Ric}}$ is geodesically affine. (See Remark 2.14.) A standard approximation technique extends the result to singular psh geodesics and the last line in the above is nonnegative from the local Bergman kernel approximation argument of [BB17], Theorem 3.2. \square

The present choice of M_g shares the same minimizer with D .

Theorem 2.25 ([BWN14], Proposition 3.2). *We have $M_g \geq D_g$ on $\mathcal{E}^1(X, -K_X)^K$ and a metric φ attains the equality iff it is a weak Mabuchi soliton. The modified K-energy M_g is lower bounded iff D_g is. In this case the infimums of the both functionals coincide.*

Proof. For the reader's convenience we give the proof. The one-side inequality in (2.46) is a simple consequence of Jensen's inequality and actually holds for lower-semicontinuous function of the form $f = -(\varphi - \varphi_0)$. It immediately shows $M_g \geq D_g$.

On the other hand, the supremum is attained by the solution f of

$$\frac{e^f d\mu_0}{\int_X e^f d\mu_0} = d\mu.$$

Consequently, $M_g(\varphi) = D_g(\varphi)$ iff φ is a weak Mabuchi soliton.

It remains to show $\inf_\varphi M_g = \inf_\varphi D_g \in \mathbb{R} \cup \{-\infty\}$. Set $m := \inf_\varphi M_g$. By the properness result [BBEGZ19], Theorem 2.18, we have

$$H(\mu|\mu_0) \geq \alpha E^*(\mu) - C \tag{2.51}$$

for any α smaller than Tian's α -invariant. In particular, $H(\mu|\mu_0) < \infty$ implies that μ has finite energy so that some $\varphi \in \mathcal{E}^1(X, -K_X)^K$ solves $\text{MA}_g(\varphi) = \mu$. Substitution to (2.48) yields

$$H(\mu|\mu_0) \geq m + E_g(\varphi) - \int_X (\varphi - \varphi_0) \text{MA}_g(\varphi).$$

Since the infimum of the inversion formula

$$L(\varphi) = \inf_\mu \left[H(\mu|\mu_0) + \int_X (\varphi - \varphi_0) d\mu \right]$$

is attained by $\mu = \mu_\varphi$, it follows $D_g \geq m$. \square

Totally in the same manner we observe that the coercivity $M_g(\varphi) \geq \varepsilon \inf_{\sigma \in T} J_g(\sigma^* \varphi) - C$ holds on $\mathcal{E}^1(X, -K_X)^K$ iff $D_g \geq \varepsilon \inf_{\sigma \in T} J_g(\sigma^* \varphi) - C$ on $\mathcal{E}^1(X, -K_X)^K$.

3. RELATIVE UNIFORM D-STABILITY

Bearing in mind of the last section, we introduce the algebraic (non-Archimedean) counterpart of modified energies and define the appropriate notion of stability which should characterize the existence of Mabuchi soliton. We first recall the notion of D-stability introduced by [B16]. The terminology here is due to our previous work [BHJ15], [BHJ19].

3.1. Uniform D-stability. We first require that a test configuration $\pi: (\mathcal{X}, \mathcal{L}) \rightarrow \mathbb{A}^1$ of a polarized manifold (X, L) is a family of polarized schemes defined over the affine line \mathbb{A}^1 . In fact we may and should allow \mathcal{L} to be only relatively semiample and \mathbb{Q} -Cartier divisor. Further, \mathcal{X} is normal variety and endowed with a lifted \mathbb{G}_m -action $\lambda: \mathbb{G}_m \rightarrow \text{Aut}(\mathcal{X}, \mathcal{L})$ such that the projection π is equivariant for λ and the standard \mathbb{G}_m -action to \mathbb{A}^1 . The datum includes the isomorphism

$$\pi^{-1}(\mathbb{A}^1 \setminus \{0\}) \simeq X \times (\mathbb{A}^1 \setminus \{0\})$$

which sends the line bundle \mathcal{L} equivariantly to $L_{\mathbb{A}^1} = p_1^* L$. Although we are concerned with the case $L = -K_X$, \mathcal{L} is still not equivalent to $-K_{\mathcal{X}/\mathbb{P}^1}$. Since we assumed \mathcal{X} is normal, $K_{\mathcal{X}}$ is at least well-defined as a Weil divisor, however, it is even not the line bundle in general. Note that some literatures consider the family over the projective

line \mathbb{P}^1 . This is equivalent to our setting because one can always obtain the unique compactified family $(\bar{\mathcal{X}}, \bar{\mathcal{L}}) \rightarrow \mathbb{P}^1$ which is trivial around $\infty \in \mathbb{P}^1$.

Example 3.1. *Every one-parameter subgroup $\mu \in N$ defines a product family $X_{\mathbb{A}^1} = X \times \mathbb{A}^1$ endowed with the non-trivial action: $\lambda(\sigma)(x, \tau) := (\mu(\sigma)x, \sigma\tau)$ for $\sigma \in \mathbb{G}_m$. We call it a product configuration generated by μ . Therefore, test configuration can be seen as a far generalization of one-parameter subgroup. Note that the compactified family is no longer a product space. For example, the product space $\mathbb{P}^1 \times \mathbb{A}^1$ endowed with a non-trivial \mathbb{G}_m -action is compactified to the Hirzebruch surface $\mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(d))$.*

After the compactification $(\bar{\mathcal{X}}, \bar{\mathcal{L}})$ we may take the intersection number e.g. $\bar{\mathcal{L}}^{n+1}$. Since we assumed \mathcal{X} is normal, $K_{\mathcal{X}}$ is at least well-defined as a Weil divisor and $K_{\bar{\mathcal{X}}}\bar{\mathcal{L}}^n$ essentially gives the famous Donaldson-Futaki invariant, or equivalently, non-Archimedean K-energy $M^{\text{NA}}(\mathcal{X}, \mathcal{L})$ introduced in [BHJ15]. This is a natural generalization of (2.29) to arbitrary test configurations.

Let us define the non-Archimedean D-energy. Recall that for given divisors \mathcal{B}, \mathcal{D} the log-canonical threshold $\text{lct}_{(\bar{\mathcal{X}}, \mathcal{B})}(\mathcal{D})$ is defined to be the supremum of $c \in \mathbb{R}$ such that the log pair $(\bar{\mathcal{X}}, \mathcal{B} + c\mathcal{D})$ has at worst log canonical singularities. Choosing the boundary divisor \mathcal{B} linearly equivalent to $-K_{\bar{\mathcal{X}}/\mathbb{P}^1} - \bar{\mathcal{L}}$, the quantity reflects the positivity of the canonical divisor. Notice that in this choice the log-canonical divisor $K_{\bar{\mathcal{X}}} + \mathcal{B} \sim_{\mathbb{Q}} -\bar{\mathcal{L}} + \pi^*K_{\mathbb{P}^1}$ is \mathbb{Q} -Cartier so that the log discrepancies and log canonical singularities are well defined for any $(\mathcal{X}, \mathcal{L})$.

Definition 3.2. *For a test configuration $\pi: (\mathcal{X}, \mathcal{L}) \rightarrow \mathbb{A}^1$ we define*

$$L^{\text{NA}}(\mathcal{X}, \mathcal{L}) := \text{lct}_{(\bar{\mathcal{X}}, \mathcal{B})}(\mathcal{X}_0) - 1, \quad E^{\text{NA}}(\mathcal{X}, \mathcal{L}) := \frac{\bar{\mathcal{L}}^{n+1}}{(n+1)V},$$

where \mathcal{X}_0 is the scheme theoretic central fiber and the boundary divisor is chosen $\mathcal{B} \sim_{\mathbb{Q}} -K_{\bar{\mathcal{X}}/\mathbb{P}^1} - \bar{\mathcal{L}}$. We say a Fano manifold X is D-semistable if the non-Archimedean D-energy

$$D^{\text{NA}}(\mathcal{X}, \mathcal{L}) := L^{\text{NA}}(\mathcal{X}, \mathcal{L}) - E^{\text{NA}}(\mathcal{X}, \mathcal{L})$$

is semipositive for all test configurations.

If $(\mathcal{X}, \mathcal{L})$ is the product test configuration generated by $\mu \in N$, we write $D^{\text{NA}}(\mathcal{X}, \mathcal{L})$ as $D^{\text{NA}}(\mu)$. It is known to be equivalent to the Futaki character (2.28), i.e.

$$D^{\text{NA}}(\mu) = F(\mu) \tag{3.1}$$

holds for every $\mu \in N$. One can also define D-stability and D-polystability of a Fano manifold. See [B16], [BHJ15], and [F16] for the detail treatment. The uniform version is more important for us. We say that a test configuration $(\mathcal{X}', \mathcal{L}')$ is a pull-back of $(\mathcal{X}, \mathcal{L})$ if a birational equivariant morphism $f: \mathcal{X}' \rightarrow \mathcal{X}$ yields $\mathcal{L}' = f^*\mathcal{L}$. Two test configurations are called *equivalent* if they admit a common pull-back. It is easy to see that the above invariants have the same values for the equivalent test configurations. More substantially, by [BHJ15], any test configurations can be seen as a fiber metric of the Berkovich analytification $(\mathcal{X}^{\text{NA}}, \mathcal{L}^{\text{NA}})$, evaluating each valuation on the central fiber. The equivalence of test configurations are precisely of corresponding non-Archimedean

fiber metrics. The above L^{NA} and E^{NA} are actually functionals defined on these non-Archimedean fiber metrics. From this reason, taking a pull-back we may assume a domination $\rho: (\mathcal{X}, \mathcal{L}) \rightarrow X_{\mathbb{A}^1}$ to the product family endowed with a possibly non-trivial action. By the projection formula the following definition is actually independent of ρ .

Definition 3.3. *We define*

$$L_0^{\text{NA}}(\mathcal{X}, \mathcal{L}) := V^{-1}(\rho^* L_{\mathbb{A}^1}) \bar{\mathcal{L}}^n.$$

and the non-Archimedean counterpart of Aubin's J -functional as $J^{\text{NA}}(\mathcal{X}, \mathcal{L}) := L_0^{\text{NA}}(\mathcal{X}, \mathcal{L}) - E^{\text{NA}}(\mathcal{X}, \mathcal{L})$. A Fano manifold is called uniformly D -stable if there exists a constant $\varepsilon > 0$ such that

$$D^{\text{NA}}(\mathcal{X}, \mathcal{L}) \geq \varepsilon J^{\text{NA}}(\mathcal{X}, \mathcal{L})$$

holds for all test configurations.

Let us illustrate a key relation between the functionals E, J, D and their non-Archimedean version. It explains that test configuration gives the algebraic formulation of the geodesic ray on \mathcal{H} . In the sequel we denote the fiber of $\tau \in \mathbb{A}^1$ by \mathcal{X}_τ and the restricted line bundle by \mathcal{L}_τ . As well, for the unit disk Δ we set $\mathcal{X}_\Delta := \pi^{-1}(\Delta)$ and $\mathcal{L}_\Delta := \mathcal{L}|_{\mathcal{X}_\Delta}$. For the punctured disk $\Delta^* = \Delta \setminus \{0\}$ we have the isomorphism $\mathcal{X}_{\Delta^*} \simeq X \times \Delta^*$ so that identify a point of \mathcal{X}_{Δ^*} with (x, τ) . Let Φ be a smooth fiber metric of \mathcal{L}_Δ , having the semipositive curvature. It defines the ray

$$\varphi^t(x) = \Phi(\lambda(e^{-t})(x, 1)) \quad (3.2)$$

so that φ^t for each $t \in [0, \infty)$ defines a fiber metric of L , having the semipositive curvature. This type of ray is said to be compatible with the test configuration. Any two metrics defines the same asymptotic because the difference of the associated rays is bounded uniformly in t . The following type of results is predicted in the origination of K-stability and proved for arbitrary test configurations in [B16], [BHJ19].

Theorem 3.4. *Let $F: \mathcal{H} \rightarrow \mathbb{R}$ be a functional either E, J , or D . For a test configuration and a ray φ^t compatible with $(\mathcal{X}, \mathcal{L})$ we have*

$$F^{\text{NA}}(\mathcal{X}, \mathcal{L}) = \lim_{t \rightarrow \infty} \frac{F(\varphi^t)}{t}.$$

The above formula is indeed true for non-smooth but bounded Φ for which the semipositivity $dd^c \Phi \geq 0$ holds in the sense of current. In particular the same result holds for the associated psh geodesic ray φ^t which is characterized by the degenerate Monge-Ampère equation

$$(dd_{\tau,x}^c \Phi)^{n+1} = 0 \quad (3.3)$$

on \mathcal{X}_Δ . Given a smooth boundary value φ^0 , the bounded solution uniquely exists. For example [B16] gives the solution in terms of the Peron-Bremermann type envelope. After [CTW18] it is known to have the best-possible $C^{1,1}$ -regularity. We have already emphasized that the consideration of weak geodesic is necessary for the variational approach.

Regarding Theorem 3.4, as in [DL20] we introduce for psh geodesic ray φ^t the *radial energy*

$$\hat{F}(\Phi) := \lim_{t \rightarrow \infty} \frac{F(\varphi^t)}{t}, \quad (3.4)$$

which is well-defined by the convexity property.

As a consequence of Theorem 3.4, the coercivity of the D-energy implies that X is uniformly D-stable. The heart of [BBJ21] is showing the converse direction. It is known that the uniform stability implies that the automorphism group is finite (see [BHJ19] for the analytic discussion and the purely algebraic proof [BX19]). What we are going to discuss suggests the one of the treatment for general automorphism groups.

3.2. Associated concave function and Duistermaat-Heckman measure. We continuously fix an extremal one-parameter subgroup η and a torus $T \subset \text{Aut}^0(X, \eta)$. As in the previous subsections we denote the lattice of one-parameter subgroups by N and the dual by M . Let $P \subset M_{\mathbb{R}}$ be the moment polytope of the maximal torus and

$$m_\varphi: X \rightarrow P$$

the moment map. Recall that the Duistermaat-Heckman measure is the push-forward

$$\text{DH}_T := (m_\varphi)_*(V^{-1}\omega^n) \quad (3.5)$$

which is also independent of the metric. Let us first give an algebraic definition of DH_T . Any $\mu \in N_{\mathbb{R}}$ is identified with the affine function $G_\mu(x) := \langle \mu, x \rangle$ on $M_{\mathbb{R}}$ hence we may integrate by DH_T . Let $k \in \mathbb{N}$ and μ_1, \dots, μ_{N_k} be the weight of the \mathbb{G}_m -action on $H^0(X, kL)$, induced by μ . For any $p \geq 1$, the equivariant Riemann-Roch formula implies

$$\int_P G_\mu^p(x) \text{DH}_T = \lim_{k \rightarrow \infty} \frac{1}{k^p N_k} \sum \mu_i^p. \quad (3.6)$$

If set $\text{DH}_\mu := (h_\mu)_*(V^{-1}\omega^n) = (G_\mu)_*\text{DH}_T$, by the Hausdorff moment theorem we obtain the convergence of the measures on \mathbb{R} :

$$\text{DH}_\mu = \lim_{k \rightarrow \infty} \frac{1}{N_k} \sum \delta_{\frac{\mu_i}{k}}. \quad (3.7)$$

A simple argument checks that P is the closed convex hull of the set

$$\left\{ \frac{\chi}{k} \in M_{\mathbb{Q}} : \chi \in M, s_\chi \in H^0(X, kL) \text{ with } \sigma \cdot s_\chi = \chi(\sigma)s_\chi \right\} \quad (3.8)$$

and that

$$\text{DH}_T = \lim_{k \rightarrow \infty} \frac{1}{N_k} \sum \delta_{\frac{\chi}{k}}, \quad (3.9)$$

where $\chi \in M$ runs for all $s_\chi \in H^0(X, kL)$. As a consequence

Proposition 3.5. *For every $\mu \in N_{\mathbb{Q}}$ we have*

$$m_X := \inf_P G_{\kappa+\mu} = \lim_{k \rightarrow \infty} \min \frac{k + \mu_i}{k} = \inf_X (1 + h_\mu).$$

In particular $\inf_X (1 + h_\mu)$ is independent of the metric.

More generally, a test configuration defines a concave function $G_{(\mathcal{X}, \mathcal{L})}$ on P . For the purpose it is convenient to describe the test configuration in terms of the filtration.

Definition 3.6. *Let $(\mathcal{X}, \mathcal{L})$ be a test configuration. Given $s \in H^0(X, L)$, we have a rational section*

$$\bar{s}(x, \tau) = \lambda(\tau) \cdot s(\lambda(\tau^{-1})(x, \tau))$$

of \mathcal{L} . Considering how extent \bar{s} is holomorphic we obtain a filtration of the section ring, which fully recovers the test configuration. For each $\lambda \in \mathbb{R}$ we set

$$F^\lambda H^0(X, kL) := \{s \in H^0(X, kL) : \tau^{-\lceil \lambda \rceil} \bar{s} \in H^0(\mathcal{X}, k\mathcal{L})\}. \quad (3.10)$$

We may easily show that the filtration is monotone, left-continuous, and multiplicative in both k and λ . By [BHJ15] lemma 2.14, λ -weightspace of the induced action to $H^0(\mathcal{X}_0, k\mathcal{L}_0)$ is given by

$$H^0(\mathcal{X}_0, k\mathcal{L}_0)_\lambda \simeq F^\lambda H^0(X, kL) / F^{\lambda+1} H^0(X, kL). \quad (3.11)$$

The identification also follows from Proposition 3.13 below. A non-trivial fact proved in [PS07] is the linearly boundedness. Namely there exists a constant $C > 0$ such that

$$F^{kt} H^0(X, kL) = \{0\} \text{ (resp. } H^0(X, kL)\text{)} \quad (3.12)$$

for any $t > C$ (resp. $t < -C$) and $k \geq 1$. It is equivalent to say: $|\lambda| \leq Ck$ for the induced \mathbb{G}_m -action.

Imitating [WN12], we construct a concave function from the filtration.

Definition 3.7. *For each $t \in \mathbb{R}$ we define P^t as the closed convex hull of the set*

$$\left\{ \frac{\chi}{k} \in M_{\mathbb{Q}} : \chi \in M, s_\chi \in F^{kt} H^0(X, kL) \text{ with } \sigma \cdot s_\chi = \chi(\sigma) s_\chi \right\}.$$

The associated concave function is then defined as

$$G_{(\mathcal{X}, \mathcal{L})}(x) := \sup\{t \in \mathbb{R} : x \in P^t\}.$$

It is easy to check $G_{(\mathcal{X}, \mathcal{L})} = G_\mu$ when $(\mathcal{X}, \mathcal{L})$ is the product configuration generated by $\mu \in N$. Indeed from the definitions we compute

$$\begin{aligned} \tau^{-kt} \bar{s}_\chi(x, \tau) &= \tau^{-kt} \mu(\tau) \cdot s_\chi(\mu(\tau^{-1})x) \\ &= \tau^{-kt} (\mu(\tau) \cdot s_\chi)(x) \\ &= \tau^{-kt + \langle \mu, \chi \rangle} s_\chi(x). \end{aligned}$$

In terms of the associated weak geodesic ray (3.3) we may extend (3.7) to T -equivariant test configurations. Notice that the weak geodesic ray φ^t has $C^{1,1}$ -regularity and the right-derivative

$$\dot{\varphi}^0 := \inf_{t>0} \frac{\varphi^t - \varphi^0}{t} \quad (3.13)$$

pointwisely defined is hence a bounded function. This fact reflects the linearly boundedness (3.12).

Theorem 3.8. *Let $(\mathcal{X}, \mathcal{L})$ be a T -equivariant test configuration. For each $k \in \mathbb{N}$ $\lambda_1, \dots, \lambda_{N_k}$ denote the weights of the induced \mathbb{G}_m -action to $H^0(\mathcal{X}_0, k\mathcal{L}_0)$. The push-forward*

$$\mathrm{DH}_{(\mathcal{X}, \mathcal{L})} := \dot{\varphi}_*^0(V^{-1}\omega^n)$$

defines a probability measure on \mathbb{R} , which is independent of the metric. Moreover, it is equivalent to

$$(G_{(\mathcal{X}, \mathcal{L})})_* \mathrm{DH}_T = \lim_{k \rightarrow \infty} \frac{1}{N_k} \sum \delta_{\frac{\lambda_i}{k}}.$$

Proof. The identity $\mathrm{DH}_{(\mathcal{X}, \mathcal{L})} = \lim_{k \rightarrow \infty} \frac{1}{N_k} \sum \delta_{\frac{\lambda_i}{k}}$ is shown in [H16a]. For any $p \geq 1$ we have

$$\int_{\mathbb{R}} t^p (G_{(\mathcal{X}, \mathcal{L})})_* \sum \delta_{\frac{\lambda_i}{k}} = \int_P G_{(\mathcal{X}, \mathcal{L})}^p \sum \delta_{\frac{\lambda_i}{k}} = \sum G_{(\mathcal{X}, \mathcal{L})}^p \left(\frac{\chi}{k} \right),$$

where the summation is for all $s_\chi \in H^0(X, kL)$. In view of the Hausdorff moment theorem it remains to show

$$\sum G_{(\mathcal{X}, \mathcal{L})}^p \left(\frac{\chi}{k} \right) = \sum \left(\frac{\lambda_i}{k} \right)^p.$$

We fix χ and by the linearly boundedness (3.12) take the largest t such that $s_\chi \in F^{\lceil kt \rceil} H^0(X, kL)$ but $s_\chi \notin F^{\lceil kt \rceil + 1} H^0(X, kL)$. From (3.11) such kt one-to-one corresponds to λ_i so we complete the proof. \square

Remark 3.9. *In [WN12], the associated concave function on the Okounkov body is in fact defined for any possibly non-equivariant test configuration. If X is toric polarized manifold the Okounkov body and the associated concave function is equivalent to the present construction. In our setting $G_{(\mathcal{X}, \mathcal{L})}$ is to be a piecewise-linear function.*

We conclude this subsection describing the invariants $E^{\mathrm{NA}}, J^{\mathrm{NA}}$ in terms of the \mathbb{G}_m -action.

Proposition 3.10 ([BHJ15] Proposition 7.8, Theorem 5.16). *For any test configuration, the non-Archimedean Monge-Ampère energy satisfies*

$$E^{\mathrm{NA}}(\mathcal{X}, \mathcal{L}) = \int_{\mathbb{R}} t \mathrm{DH}_{(\mathcal{X}, \mathcal{L})} = \lim_{k \rightarrow \infty} \frac{1}{N_k} \sum_{i=1}^{N_k} \frac{\lambda_i}{k}.$$

The functional L_0 satisfies

$$L_0^{\mathrm{NA}}(\mathcal{X}, \mathcal{L}) = \sup \mathrm{supp} \mathrm{DH}_{(\mathcal{X}, \mathcal{L})} = \lim_{k \rightarrow \infty} \max_i \frac{\lambda_i}{k}.$$

Moreover, $\max_i \frac{\lambda_i}{k}$ is stable in k . More precisely, it is enough to take a sufficiently divisible k so that $k\mathcal{L}$ is globally generated. If there exists a domination $\rho: \mathcal{X} \rightarrow X_{\mathbb{A}^1}$, let E_0 be the strict transform of $X \times \{0\}$ and $D := \mathcal{L} - \rho^ L_{\mathbb{A}^1}$ be the unique \mathbb{Q} -divisor supported on \mathcal{X}_0 . We then have*

$$L_0^{\mathrm{NA}}(\mathcal{X}, \mathcal{L}) = \mathrm{ord}_{E_0} D.$$

3.3. Relative setting. Let us discuss the relative stability. From now on we fix η , $G = \text{Aut}(X, \eta)$, $T = C(G)$, and $K \subset G$ a maximal compact subgroup containing S . The identical one-parameter subgroup to T is denoted by $\kappa \in N$. Test configurations are assumed to be G -equivariant, as the metrics were K -invariant.

Definition 3.11. *Let G be a reductive algebraic group. A test configuration $(\mathcal{X}, \mathcal{L})$ endowed with $\mathbb{G}_m \times G$ -action is G -equivariant if it is compatible with the equipped \mathbb{G}_m -action on $(\mathcal{X}, \mathcal{L})$ and the G -action on $(X, L) = (\mathcal{X}_1, \mathcal{L}_1)$.*

Notice that we imposed the commutativity with G , on the \mathbb{G}_m -action. In particular G acts on the central fiber \mathcal{X}_0 . The G -equivariance is not too much restrictive, as one can see [DS16], Theorem 1, in the Kähler-Einstein case. See also Example 3.23 below.

Example 3.12. *Consider a G -invariant ideal $I \subset \mathcal{O}_X$. Let $\rho: \mathcal{X} \rightarrow X_{\mathbb{A}^1}$ be the normalization of the blow-up along the ideal $\mathcal{J} := I\mathcal{O}_{X_{\mathbb{A}^1}} + \tau\mathcal{O}_{X_{\mathbb{A}^1}}$ and E be the exceptional divisor. We take $\varepsilon > 0$ and set $\mathcal{L} := \rho^*L_{\mathbb{A}^1} - \varepsilon E$. This typical test configuration called deformation to the normal cone is intensively studied in [RT07]. Indeed one can show that \mathcal{L} is ample for every sufficiently small ε . If for example the support V of \mathcal{J} is smooth we may write $E = \mathbb{P}(N_{V/X} \oplus \mathcal{O}_X)$ as the normal cone. The induced \mathbb{G}_m -action is trivial on the normal bundle $N_{V/X}$ and the is the simple multiplication on \mathcal{O}_X . Since \mathcal{J} is G -invariant, $(\mathcal{X}, \mathcal{L})$ inherits the G -action so that ρ is equivariant. In this construction we observe that the two actions actually commute to each other.*

Let us consider the case $G = \text{Aut}(X, -K_X)$. For example, \mathbb{P}^2 does not have any G -invariant ideal. If X is the one point blow-up of \mathbb{P}^2 any G -invariant ideal is supported on the exceptional divisor. We may check that the deformation to the normal cone prevent X to be D -semistable.

The starting point here is to take the inner product of such a test configuration with arbitrary one-parameter subgroups, extending the definition of [FM95]. The equipped $\lambda: \mathbb{G}_m \rightarrow \text{Aut}(\mathcal{X}, \mathcal{L})$ induces the action to $H^0(\mathcal{X}_0, k\mathcal{L}_0)$ for every $k \geq 1$. Since \mathcal{X} is normal and is a family over the curve, it is flat. It follows $H^0(\mathcal{X}_0, k\mathcal{L}_0) \simeq H^0(X, kL)$ for any sufficiently large k . In fact we may have a G -equivariant trivialization of the vector bundle $\pi_*(k\mathcal{L})$ over \mathbb{A}^1 .

Proposition 3.13. *The G -equivariant algebraic vector bundle $E = \pi_*(k\mathcal{L})$ on the affine line \mathbb{A}^1 is G -equivariantly isomorphic to $E_0 \times \mathbb{A}^1$.*

Proof. For the case $G = \{\text{id}\}$ we refer [BHJ15] Proposition 1.3. Taking M into the account the same argument works. Indeed from the commutativity of the first component \mathbb{G}_m with the second G , G -action does not effect. Let M_G be the lattice of weights and

$$H^0(\mathbb{A}^1, E) = \bigoplus_{(\lambda, \chi) \in \mathbb{Z} \oplus M_G} H^0(\mathbb{A}^1, E)_{(\lambda, \chi)} \quad (3.14)$$

be the decomposition to the irreducible representations. Set $V := E_1 = H^0(X, kL)$, $V_\chi := H^0(X, kL)_\chi$ and $F^\lambda V_\chi$ as the image of $H^0(\mathbb{A}^1, E)_{(\lambda, \chi)}$ under the restriction map $H^0(\mathbb{A}^1, E) \rightarrow E_1$. Definition 3.11 implies $F^\lambda V_\chi \subset V_\chi$. Since τ has weight -1 with respect to the \mathbb{G}_m -action on the base \mathbb{C} , multiplication by τ induces $F^{\lambda+1} V_\chi \subset F^\lambda V_\chi$. Since $F^\lambda V_\chi = V_\chi$ for $\lambda \ll 0$ and $V = \bigoplus_\chi V_\chi$, the above map sending $\sum \tau^{-\lambda} v_\lambda$ to $\sum v_\lambda$

is surjective. On the other hand, if $\sum \tau^{-\lambda} v_\lambda$ lies in the kernel, $w_\lambda := -\sum_{\lambda' \geq \lambda} v_{\lambda'}$ in $F^\lambda V := \bigoplus_{\chi} F^\lambda V_\chi$ vanishes for $\lambda \ll 0$. Since $v_\lambda = w_{\lambda+1} - w_\lambda$, it means that $\sum \tau^{-\lambda} v_\lambda$ is in $(\tau - 1)H^0(\mathbb{A}^1, E)$. Thus we have $H^0(\mathbb{A}^1, E)/(\tau - 1)H^0(\mathbb{A}^1, E) \simeq V$ and the equivariant isomorphism

$$E|_{\mathbb{A}^1 \setminus \{0\}} \simeq V \times \mathbb{A}^1 \setminus \{0\}.$$

Similarly, by sending $\sum \tau^{-\lambda} v_\lambda$ to v_λ modulo $F^{\lambda+1}V$ we may show

$$E_0 \simeq \bigoplus_{\lambda \in \mathbb{Z}} F^\lambda V / F^{\lambda+1} V.$$

It follows the equivariant isomorphism

$$H^0(X, E) \simeq \bigoplus_{\lambda \in \mathbb{Z}} \tau^{-\lambda} F^\lambda V.$$

By choosing a basis compatible the filtration and $F^\lambda V = \bigoplus_{\chi} F^\lambda V_\chi$, we obtain a required equivariant trivialization. \square

Now consider $G = \text{Aut}^0(X, \eta)$. Because the test configuration is assumed to be G -equivariant, given $\mu \in N$ we may simultaneously diagonalize the two actions on $H^0(X, kL)$ and $H^0(\mathcal{X}_0, k\mathcal{L}_0)$ so that each weights λ_i and μ_i are assigned for the common vectors under the equivariant trivialization. In the sequel we may take any such λ_i and μ_i .

Definition 3.14 ([H16b]). *Let $(\mathcal{X}, \mathcal{L})$ be a T -equivariant test configuration. For any one-parameter subgroup $\mu \in N$ we have the limit*

$$\langle (\mathcal{X}, \mathcal{L}), \mu \rangle := \lim_{k \rightarrow \infty} \frac{1}{k^2 N_k} \sum_{i=1}^{N_k} \lambda_i \mu_i.$$

When $(\mathcal{X}, \mathcal{L})$ is a product configuration generated by some $\lambda \in N$, we denote $\langle (\mathcal{X}, \mathcal{L}), \mu \rangle =: \langle \lambda, \mu \rangle$.

For the identical one-parameter subgroup $\kappa \in N$ we observe $\mu_i = k$ and hence Proposition 3.10 shows

$$\langle (\mathcal{X}, \mathcal{L}), \kappa \rangle = \lim_{k \rightarrow \infty} \frac{1}{k N_k} \sum_{i=1}^{N_k} \lambda_i.$$

It is easy to check that homogeneity naturally extends the above inner product to $\mu \in N_{\mathbb{Q}}$. We may further extend the definition to $\mu \in N_{\mathbb{R}}$ by the following description.

Theorem 3.15 ([H16b]). *Let $(\mathcal{X}, \mathcal{L})$ be a T -equivariant test configuration and $\mu \in N_{\mathbb{Q}}$. For the associated weak geodesic ray φ^t and the Hamilton function h_μ we have*

$$\langle (\mathcal{X}, \mathcal{L}), \mu \rangle = \frac{1}{V} \int_X \dot{\varphi}^0 h_\mu \omega^n.$$

In view of the above result, it is also natural to define

$$\langle \Phi, \mu \rangle := \frac{1}{V} \int_X \dot{\varphi}^0 h_\mu \omega^n \tag{3.15}$$

for arbitrary psh geodesic Φ with $C^{1,1}$ -regularity. When Φ is associated with a test configuration the right-hand side gives the slope of the affine function $E_g(\varphi^t)$.

Corollary 3.16. *Let $(\mathcal{X}, \mathcal{L})$ be a test configuration and φ^t the associated weak geodesic ray. For any $\mu \in N_{\mathbb{Q}}$ putting $g(x) := 1 + \langle x, \mu \rangle - \int_P \langle x, \mu \rangle \text{DH}_T$ (compatible with (2.32)) we have*

$$\langle (\mathcal{X}, \mathcal{L}), \mu \rangle = \lim_{t \rightarrow \infty} \frac{E_g(\varphi^t)}{t}.$$

Now we choose the extremal one-parameter subgroup $\eta \in N_{\mathbb{Q}}$ to define the non-Archimedean counterpart of E_{η} .

Definition 3.17. *The non-Archimedean counterpart of the modified Monge-Ampère energy E_{η} is defined to be*

$$E_{\eta}^{\text{NA}}(\mathcal{X}, \mathcal{L}) := \langle (\mathcal{X}, \mathcal{L}), \kappa + \eta \rangle.$$

We introduce the modified non-Archimedean energies as $D_{\eta}^{\text{NA}}(\mathcal{X}, \mathcal{L}) := L^{\text{NA}}(\mathcal{X}, \mathcal{L}) - E_{\eta}^{\text{NA}}(\mathcal{X}, \mathcal{L})$ and $J_{\eta}^{\text{NA}}(\mathcal{X}, \mathcal{L}) := L_0^{\text{NA}}(\mathcal{X}, \mathcal{L}) - E_{\eta}^{\text{NA}}(\mathcal{X}, \mathcal{L})$.

Note that J_{η}^{NA} is not necessarily non-negative, just as J_{η} was. We shall see that $T \subset \text{Aut}(X, \eta)$ is enough to examine the positivity of J_{η}^{NA} .

Proposition 3.18. *If $m_X > 0$, then $J_{\eta}^{\text{NA}}(\mathcal{X}, \mathcal{L}) \geq 0$ and the equality holds iff the T -equivariant $(\mathcal{X}, \mathcal{L})$ is the trivial test configuration.*

Proof. It is immediate from Theorem 3.15 and 3.10 that

$$J_{\eta}^{\text{NA}}(\mathcal{X}, \mathcal{L}) = \sup_X \varphi^0 - \frac{1}{V} \int_X \varphi^0 (1 + h_{\eta}) \omega^n. \quad (3.16)$$

Since we may rescale the \mathbb{G}_m -action to have $\sup_X \varphi^0 = 0$, from the formula $m_X = \inf_X (1 + h_{\eta}) > 0$ implies $J_{\eta}^{\text{NA}}(\mathcal{X}, \mathcal{L}) > 0$, otherwise φ^0 is identically zero. By [BHJ15] Theorem A, $\varphi^0 \equiv 0$ implies that $(\mathcal{X}, \mathcal{L})$ is trivial. That is, the product configuration with the trivial action. \square

In terms of the associated concave function, we may write

$$J_{\eta}^{\text{NA}}(\mathcal{X}, \mathcal{L}) = \max_P G_{(\mathcal{X}, \mathcal{L})} - \frac{1}{V} \int_P G_{(\mathcal{X}, \mathcal{L})} G_{\kappa + \eta} \text{DH}_T. \quad (3.17)$$

In our definition of stability we assume $m_X > 0$. By Proposition 3.5, this additional assumption is very much easier to check than the positivity of D_{η}^{NA} for all test configurations.

Let us return to a general $\mu \in N$ and take a G -equivariant trivialization so that the weights λ_i and μ_i are assigned for the common vectors. We endow a new \mathbb{G}_m -action with the space $(\mathcal{X}, \mathcal{L})$ such that the weights are given by $\lambda_i + \mu_i$. Since $T = C(G)$, it indeed gives a G -equivariant test configuration which we will denote by $(\mathcal{X}_{\mu}, \mathcal{L}_{\mu})$. If Φ is the weak geodesic ray associated with $(\mathcal{X}, \mathcal{L})$, it is easy to see that

$$\varphi_{\mu}^t(x) := \Phi(\lambda(e^{-t})\mu(e^{-t})(x, 1)) \quad (3.18)$$

gives the geodesic ray associated with $(\mathcal{X}_{\mu}, \mathcal{L}_{\mu})$. The homogeneity naturally extends the definition to arbitrary $\mu \in N_{\mathbb{Q}}$. From Theorem 3.15, we may further observe that

$J^{\text{NA}}(\mathcal{X}_\mu, \mathcal{L}_\mu)$ is continuous in $\mu \in N_{\mathbb{Q}}$. The next rationality lemma is a key to show the existence of the metric in Theorem A.

Lemma 3.19 (rationality of the minimizer). *The functional $\mu \mapsto J_\eta^{\text{NA}}(\mathcal{X}_\mu, \mathcal{L}_\mu)$ is rationally piecewise-linear convex function in $N_{\mathbb{R}}$. It is moreover strictly convex in $N_{\mathbb{R}}/\mathbb{R}$. Especially the infimum*

$$J_T^{\text{NA}}(\mathcal{X}, \mathcal{L}) := \inf_{\mu \in N_{\mathbb{R}}} J^{\text{NA}}(\mathcal{X}_\mu, \mathcal{L}_\mu) \quad (3.19)$$

is attained by a rational μ .

Proof. The result was observed in [H18]. Indeed by Proposition 3.10 we see that

$$J_\eta^{\text{NA}}(\mathcal{X}_\mu, \mathcal{L}_\mu) = \max_i \frac{\lambda_i + \mu_i}{k} - \frac{1}{N_k} \sum_{i=1}^{N_k} (\lambda_i + \mu_i)(\kappa + \eta_i).$$

Thanks to Proposition 3.10 the first term is independent of k , as soon as $k\mathcal{L}$ is globally generated. Note that the condition is independent of μ . The second term is affine in μ . Therefore, as the function in μ , it is the maximum for finite number of affine functions. The function is obviously non-negative and proper in $N_{\mathbb{R}}/\mathbb{R}$. \square

The notation J_T and J_T^{NA} are consistent. We indeed have the slope formula which is the main ingredient in deriving the stability in Theorem A.

Theorem 3.20 ([H18], Theorem B). *Let $(\mathcal{X}, \mathcal{L})$ be a T -equivariant test configuration and φ^t be the associated weak geodesic ray. We have*

$$J_T^{\text{NA}}(\mathcal{X}, \mathcal{L}) = \lim_{t \rightarrow \infty} \frac{J_T(\varphi^t)}{t}.$$

Notice that $\sigma \in T$ attaining the infimum of $J_T(\varphi^t)$ depends on t . It is at least technically crucial to fix one torus in obtaining this sort of slope formulas. See [H18], Remark 1.8. Based on the results, we now arrive at the definition of the desired stability.

Definition 3.21. *A Fano manifold X is uniformly relatively D -stable if $m_X > 0$, $G = \text{Aut}(X, \eta)$ is reductive, and there exists a constant $\varepsilon > 0$ such that*

$$D_\eta^{\text{NA}}(\mathcal{X}, \mathcal{L}) \geq \varepsilon J_T^{\text{NA}}(\mathcal{X}_\mu, \mathcal{L}_\mu)$$

holds for any G -equivariant test configuration. We say that X is relatively D -semistable if $D_\eta^{\text{NA}}(\mathcal{X}, \mathcal{L}) \geq 0$ for any G -equivariant test configuration.

Remark 3.22. *In a recent preprint [Y22] it was proved that The obstruction $m_X > 0$ about the automorphism group automatically follows from the condition $D_\eta^{\text{NA}}(\mathcal{X}, \mathcal{L}) \geq \varepsilon J_T^{\text{NA}}(\mathcal{X}_\mu, \mathcal{L}_\mu)$.*

Our formulation endows the test configurations with large symmetry G . This is considerably effective in checking the stability of specific Fano manifolds.

Example 3.23. *As we have observed, there is no G -invariant ideal when $X = \mathbb{P}^2$. It simply implies that \mathbb{P}^2 is uniformly relatively D -stable. When X is the one point blow-up of \mathbb{P}^2 , we have the deformation to the normal cone $(\mathcal{X}, \mathcal{L})$ for the exceptional divisor. We may check that $(\mathcal{X}, \mathcal{L})$ dominates the product test configuration generated by η . It*

means that $D_\eta^{\text{NA}}(\mathcal{X}, \mathcal{L}) = J_T^{\text{NA}}(\mathcal{X}, \mathcal{L}) = 0$. Indeed X admits a Mabuchi soliton, and hence it is uniformly relatively D -stable, by the following general result. See [Y17] for investigation of the general toric Fano manifolds.

In our framework explained so far, existence of the metric naturally implies the stability.

Theorem 3.24. *If a Fano manifold admits a Mabuchi soliton, then it is uniformly relatively D -stable.*

Proof. By Theorem 2.16 we have the coercivity. As a consequence of Theorem 3.4 and Theorem 3.20, the coercivity implies the stability. \square

4. VARIATIONAL APPROACH AND PROOF OF THE MAIN THEOREM

Standing on the preparation of the last two sections we give a proof of Theorem A. After we organized the formulation, the argument is now a simple extension of the variational approach [BBJ21], to the relative and equivariant setting.

4.1. Convergence of weak geodesics. Existence of the metric implies the stability, by Theorem 3.24. Let us assume that a Fano manifold is uniformly relatively D -stable, in the sense of Definition 3.21. Since J and J_η are equivalent, one may use J_η in place of J . In view of the thermodynamical formalism, we have already observed in section 2.7 that the coercivity properties of the modified D and K -energy (see Definition 2.22) are equivalent. We shall suppose that the modified K -energy is not coercive and lead the contradiction.

The first step is to construct a weak geodesic ray in which direction the modified K -energy (and therefore D -energy) is not coercive. If the coercivity of Definition 2.15 fails, we have a sequence $\varphi_j \in \mathcal{H}(X, -K_X)^K$ and $\varepsilon_j \rightarrow 0$ ($j = 1, 2, \dots$) so that

$$M_\eta(\varphi_j) \leq \varepsilon_j J(\sigma^* \varphi_j) - j \quad (4.1)$$

holds for *any* $\sigma \in T$. Since both sides are preserved by the constant rescaling $\varphi \mapsto \varphi + c$ we may take

$$\sup_X(\varphi_j - \varphi_0) = 0. \quad (4.2)$$

We may moreover assume

$$T_j := -E(\varphi_j) \rightarrow \infty, \quad (4.3)$$

otherwise the uniform version of Skoda's integrability and the weak-compactness of the level set $\{\varphi \in \mathcal{E}^1(X, -K_X)^K : E(\varphi) \geq -C\}$ imply

$$D_\eta(\varphi_j) \geq -\log C - E_\eta(\varphi_j) \geq -\log C'.$$

Here we used again the comparison (2.36) of E and E_η . Then (4.1) yields $J(\varphi_j) \rightarrow \infty$, which contradicts to the assumption $E(\varphi_j) \geq -C$ with (4.2).

Let us take a weak geodesic φ_j^t ($0 \leq t \leq -E(\varphi_j)$) which joins φ_0 to φ_j . For the convergence of φ_j^t we need to control the relative entropy.

Theorem 4.1 ([BBEGZ19], Theorem 2.17). *The sublevel set*

$$\left\{ \varphi \in \mathcal{E}^1(X, -K_X)^K : H(\text{MA}(\varphi)|\mu_0) \leq C, \sup_X(\varphi - \varphi_0) = 0 \right\}$$

is compact in the d_1 -topology.

Since $g > 0$ it is sufficient to control $H(\text{MA}_g(\varphi)|\mu_0)$. In view of the formula (2.48), the entropy bound is reduced to control M_g . Indeed we may control the last two terms in (2.48) by E , observing (2.36) and the elementary estimate

$$(n+1)E(\varphi) \leq \int_X (\varphi - \varphi_0) \text{MA}(\varphi) \leq E(\varphi). \quad (4.4)$$

The convexity of the modified K-energy now implies

$$M_\eta(\varphi_j^t) \leq \frac{t}{-E(\varphi_j)} M_\eta(\varphi_j) \leq \frac{t}{-E(\varphi_j)} (\varepsilon_j J(\sigma^* \varphi_j) - j). \quad (4.5)$$

In particular for $\sigma = \text{id}$ we obtain the bound of $M_\eta(\varphi_j^t)$. It follows from Theorem 4.1 that for each fixed T , φ_j^t ($0 \leq t \leq T$) is contained in a compact subset with respect to the strong topology. The geodesicity as well implies

$$d_1(\varphi_j^t, \varphi_j^s) = d_1(\varphi_j^1, \varphi_0) |t - s| \leq C(J(\varphi_j^1) + 1) |t - s|$$

for any $t, s \geq 0$. By Ascoli's theorem, passing through a subsequence if necessary, we conclude that φ_j^t strongly converges to φ^t . It is immediate from $E(\varphi_j^t) = -t$ that $E(\varphi^t) = -t$.

4.2. Demailly type approximation. The second step is to approximate φ^t constructed in the above by a sequence of test configurations. It is the non-Archimedean analogue of Demailly's approximation theorem for plurisubharmonic functions. Given φ^t , the relation (3.2) gives the singular K -invariant metric Φ on $L_{\mathbb{A}^1}$, defined over $X_{\mathbb{A}^1 \setminus \{0\}} = \mathbb{C}^* \times X$. Since $\sup_X(\varphi^t - \varphi_0) = 0$, the plurisubharmonicity uniquely extends Φ to \mathbb{A}^1 . Now for a sufficiently large $m \in \mathbb{N}$ we take the multiplier ideal sheaf $\mathcal{J}(m\Phi)$ and the normalized blow-up $\rho_m: \mathcal{X}^{(m)} \rightarrow \mathbb{A}^1$, endowed with the exceptional divisor E_m and the line bundle

$$\mathcal{L}^{(m)} := \rho_m^* L_{\mathbb{A}^1} - \frac{1}{m+m_0} E_m. \quad (4.6)$$

We may show that $\mathcal{L}^{(m)}$ is relatively semiample line bundle. See [BBJ21], Lemma 5.6 for the proof. We may check that the test configuration $(\mathcal{X}^{(m)}, \mathcal{L}^{(m)})$ inherits the equivariant G -action, since $\mathcal{J}(m\Phi)$ is G -invariant. Note that the central fiber \mathcal{X}_0 is the union of the strict transform E_0 of $X \times \{0\}$ and the exceptional divisor E_m . The \mathbb{G}_m -action of $(\mathcal{X}^{(m)}, \mathcal{L}^{(m)})$ is trivial on E_0 so that it commutes with the G -action.

Theorem 4.2 ([BBJ21], Theorem 5.4 and 6.4 for the $T = \{\text{id}\}$ case). *For the above test configurations constructed from φ^t , we have*

$$E_\eta^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)}) \geq \lim_{t \rightarrow \infty} \frac{E_\eta(\varphi^t)}{t},$$

$$\lim_{m \rightarrow \infty} L^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)}) = \lim_{t \rightarrow \infty} \frac{L(\varphi^t)}{t}.$$

We need the modified E_η in the above, however, the proof is the same as [BBJ21]. Indeed, using Demailly's approximation theorem locally, we have the estimate

$$\Phi_m \geq \Phi - C_{m,r} \quad (4.7)$$

on the shrunken area $\mathbb{B}(0, r) \times X$. The constant $C_{m,r}$ is necessarily independent of t . Since the modified Monge-Ampère energy is monotone, we apply Corollary 3.16 to obtain

$$\begin{aligned} E_\eta^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)}) &= \lim_{t \rightarrow \infty} \frac{E_\eta(\varphi_m^t)}{t} \\ &\geq \lim_{t \rightarrow \infty} \frac{E_\eta(\varphi^t - C_{m,r})}{t} = \lim_{t \rightarrow \infty} \frac{E_\eta(\varphi^t)}{t}. \end{aligned}$$

The key point in the above is the Ohsawa-Takegoshi L^2 -extension theorem [OT87] used in Demailly's approximation.

Let us next consider the upper bound of $D(\mathcal{X}_m, \mathcal{L}_m)$. With the convexity of the D-energy, the assumption (4.1) for $\sigma = \text{id}$ immediately implies

$$\lim_{t \rightarrow \infty} \frac{D_\eta(\varphi^t)}{t} \leq 0. \quad (4.8)$$

It then follows

$$\begin{aligned} \lim_{m \rightarrow \infty} L^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)}) &= \lim_{t \rightarrow \infty} \frac{L(\varphi^t)}{t} \\ &= \lim_{t \rightarrow \infty} \frac{D_\eta(\varphi^t) + E_\eta(\varphi^t)}{t} \leq \hat{E}_\eta(\Phi) = \langle \Phi, \eta \rangle. \end{aligned} \quad (4.9)$$

This is satisfactory for our purpose.

We aim for the lower bound estimate of L^{NA} to get the contradiction. From now on we follow the strategy of [Li22] modifying the original idea of [BBJ21]. Let us take some $\mu_m \in \mathbb{N}_\mathbb{Q}$ by Lemma 3.19 so that

$$J^{\text{NA}}(\mathcal{X}_{\mu_m}^{(m)}, \mathcal{L}_{\mu_m}^{(m)}) = J_T^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)}) := \inf_{\mu \in N_\mathbb{R}} J^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)})$$

holds. To subtract the convergent subsequence of μ_m , let us serve another simple proof of the boundedness lemma in [Li22].

Lemma 4.3 (boundedness of one-parameter subgroups, [Li22], discussion in the section 5.4). *The achievements μ_m is bounded in the vector space $N_\mathbb{R}$. Specifically, taking a norm of $N_\mathbb{R}$ we have a constant independent of m such that*

$$|\mu_m| \leq C.$$

Proof. It is sufficient to bound $J^{\text{NA}}(-\mu_m)$ or equivalent I-functional

$$\lim_{t \rightarrow \infty} \frac{I(\mu_m(e^t)^* \psi^0, \psi^0)}{t}$$

for some fixed ψ^0 . See (2.20) for the definition. We may write $\mu_m(e^t)^* \psi^0 = \psi_{(-\mu_m)}^0$ if regard ψ^0 as the constant ray. Let ψ^t be the associated ray with $(\mathcal{X}^{(m)}, \mathcal{L}^{(m)})$. Our

trick using the pseudo-triangle inequality (2.21) is

$$\begin{aligned} c_n I(\psi_{(-\mu_m)}^0, \psi^0) &\leq I(\psi_{(-\mu_m)}^0, \psi^t) + I(\psi^t, \psi^0) \\ &= I(\psi^0, \psi_{\mu_m}^t) + I(\psi^t, \psi^0). \end{aligned}$$

We may control the first term by $J_T^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)})$ and the second term by $J^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)})$. These two are bounded because

$$J^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)}) = -E^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)}) \leq \frac{-E(\varphi^t)}{t} = 1$$

and obviously $J_T^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)}) \leq J^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)})$. Notice that the similar argument was also applied to the proof of the slope formula. \square

By the above lemma, taking a subsequence if necessary we may assume μ_m converges to some $\mu \in N_{\mathbb{R}}$. The next lemma crucial in the proof shows that the twisted ray φ_{μ}^t is non-trivial. This is the point we use the assumption (4.1) for arbitrary $\sigma \in T$.

Lemma 4.4 (non-triviality of the twisted geodesic ray, [Li22], Corollary 5.3). *The radial J -energy of the twisted ray φ_{μ}^t is strictly positive. Namely*

$$\hat{J}_{\eta}(\Phi_{\mu}) > 0.$$

Proof. Here we follow the discussion of [Li22]. In view of (2.10), it is sufficient to prove $\hat{J}(\Phi_{\mu}) > 0$. We observe from (2.48) that

$$M(\varphi_j) = M(\sigma^* \varphi_j) \geq C - nJ(\sigma^* \varphi_j)$$

holds. The assumption (4.1) then yields

$$J(\sigma^* \varphi_j) \geq \frac{j + C}{n + \varepsilon_j} \rightarrow +\infty.$$

It implies $J(\varphi_{\mu}^t) \rightarrow +\infty$. Since J has linear growth along geodesics, we obtain $\hat{J}(\Phi_{\mu}) > 0$. \square

The above two lemmas furnish the proof of the main theorem. Let us decompose the L-functional as

$$L^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)}) = D_{\eta}^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)}) + E_{\eta}^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)}).$$

By the uniform stability (with in mind of (2.10) again) there exists some $\varepsilon' > 0$ such that the right-hand side is not less than

$$\begin{aligned} &\varepsilon' J_{\eta}^{\text{NA}}(\mathcal{X}_{\mu_m}^{(m)}, \mathcal{L}_{\mu_m}^{(m)}) + E_{\eta}^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)}) \\ &= \varepsilon' L_0^{\text{NA}}(\mathcal{X}_{\mu_m}^{(m)}, \mathcal{L}_{\mu_m}^{(m)}) + (1 - \varepsilon') E_{\eta}^{\text{NA}}(\mathcal{X}_{\mu_m}^{(m)}, \mathcal{L}_{\mu_m}^{(m)}) - \langle \mu_m, \kappa + \eta \rangle. \end{aligned}$$

As we may assume $1 - \varepsilon' > 0$, the second term is now controlled by Theorem 4.2. Since the functional $L_0(\varphi)$ is monotone in φ , we may conclude

$$\begin{aligned} \lim_{m \rightarrow \infty} L^{\text{NA}}(\mathcal{X}^{(m)}, \mathcal{L}^{(m)}) &\geq \varepsilon' \hat{L}_0(\Phi_{\mu}) + (1 - \varepsilon') \hat{E}_{\eta}(\Phi_{\mu}) - \langle \mu, \kappa + \eta \rangle \\ &= \varepsilon' \hat{J}_{\eta}(\Phi_{\mu}) + \hat{E}_{\eta}(\Phi). \end{aligned}$$

By Lemma 4.4 this is strictly greater than $\langle \Phi, \eta \rangle$ and it contradicts to 4.9.

REFERENCES

- [BM85] S. Bando, and T. Mabuchi: *Uniqueness of Kähler-Einstein metrics modulo connected group actions*. algebraic geometry, Sendai, 1985, 11–40, Adv. Stud. Pure Math., 10, North Holland, Amsterdam, 1987. 17
- [BM86] S. Bando, and T. Mabuchi: *On some integral invariants on complex manifolds, I*. Proc. Japan Acad. Ser. A Math. Sci., **62** (1986), no. 5, 197–200. 2
- [BT76] E. Bedford and A. Taylor: *The Dirichlet problem for a complex Monge-Ampère equation*. Invent. Math. **37** (1976), no. 1, 1–44. 6
- [B13] R. J. Berman: *A thermodynamical formalism for Monge-Ampère equations, Moser-Trudinger inequalities and Kähler-Einstein metrics*. Adv. Math. **248** (2013), 1254–1297. 17
- [B16] R. J. Berman: *K-polystability of Q-Fano varieties admitting Kahler-Einstein metrics*. Invent. Math. **203** (2016), no. 3, 973–1025. 3, 4, 16, 19, 20, 21
- [BB17] R. J. Berman and B. Berndtsson: *Convexity of the K-energy on the space of Kähler metrics and uniqueness of extremal metrics*. J. Amer. Math. Soc. **30** (2017), no. 4, 1165–1196. 18
- [BB10] R. J. Berman and S. Boucksom: *Growth of balls of holomorphic sections and energy at equilibrium*. Invent. Math. **181** (2010), no. 2, 337–394. 13
- [BBGZ13] R. J. Berman, S. Boucksom, V. Guedj, and A. Zeriahi: *A variational approach to complex Monge-Ampère equations*. Publ. Math. Inst. Hautes Études Sci. **117** (2013), 179–245. 6
- [BBEGZ19] R. J. Berman, S. Boucksom, P. Eyssidieux, V. Guedj, and A. Zeriahi: *Kähler-Einstein metrics and the Kähler-Ricci flow on log Fano varieties*. J. Reine Angew. Math. **751** (2019), 27–89. 8, 19, 30
- [BBJ21] R. J. Berman, S. Boucksom, and M. Jonsson: *A variational approach to the Yau-Tian-Donaldson conjecture*. J. Amer. Math. Soc. **34** (2021), no. 3, 605–652. 2, 7, 22, 29, 30, 31
- [BDL17] R. J. Berman, T. Darvas, and C. H. Lu: *Convexity of the extended K-energy and the large time behavior of the weak Calabi flow*. Geom. Topol. **21** (2017), no. 5, 2945–2988. 18
- [BWN14] R. Berman and D. Witt Nyström: *Complex optimal transport and the pluripotential theory of Kähler-Ricci solitons*. arXiv:1401.8264. 10, 11, 12, 13, 17, 18
- [B15] B. Berndtsson: *A Brunn-Minkowski type inequality for Fano manifolds and some uniqueness theorems in Kähler geometry*. Invent. Math. **200** (2015), no. 1, 149–200. 7, 12, 13, 16, 17
- [BP08] B. Berndtsson and M. Paun: *Bergman kernels and the pseudoeffectivity of relative canonical bundles*. Duke Math. J. **145** (2008), no. 2, 341–378. 7
- [B09] B. Berndtsson: *Curvature of vector bundles associated to holomorphic fibrations*. Ann. of Math. (2) **169** (2009), no. 2, 531–560. 7
- [BX19] H. Blum and C. Xu: *Uniqueness of K-polystable degenerations of Fano varieties*. Ann. of Math. (2) **190** (2019), no. 2, 609–656. 22
- [B18] S. Boucksom: *Variational and non-Archimedean aspects of the Yau-Tian-Donaldson conjecture*. arXiv:1805.03289.
- [BEGZ10] S. Boucksom, P. Eyssidieux, V. Guedj, and A. Zeriahi: *Monge-Ampère equations in big cohomology classes*. Acta Math. **205** (2010), no. 2, 199–262. 6, 7
- [BHJ15] S. Boucksom T. Hisamoto and M. Jonsson: *Uniform K-stability, Duistermaat-Heckman measures and singularities of pairs*. Ann. Inst. Fourier (Grenoble) **67** no. 2 (2017), 743–841. 3, 8, 19, 20, 23, 24, 25, 27
- [BHJ19] S. Boucksom T. Hisamoto and M. Jonsson: *Uniform K-stability and asymptotics of energy functionals in Kähler geometry*. J. Eur. Math. Soc. (JEMS) **21** (2019), no. 9, 2905–2944. 19, 21, 22

- [BJ18] S. Boucksom and M. Jonsson: *A non-Archimedean approach to K-stability*. [arXiv:1805.11160](https://arxiv.org/abs/1805.11160).
- [C82] E. Calabi: *Extremal Kähler metrics*. Seminar on Differential Geometry, pp. 259–290, Ann. of Math. Stud., **102**, Princeton Univ. Press, Princeton, N.J., 1982. 4
- [C85] E. Calabi: *Extremal Kähler metrics II*. Differential geometry and complex analysis, 95–114, Springer, Berlin, 1985. 4
- [C98] U. Cegrell: *Pluricomplex energy*. Acta Math. **180** (1998), no. 2, 187–217. 7
- [C00] X. X. Chen: *The space of Kähler metrics*. J. Differential Geom. **56** (2000), 189–234.
- [C00] X. X. Chen: *On the lower bound of the Mabuchi energy and its application*. Int. Math. Res. Not. **12** (2000), 607–623. 17
- [CDS15] X.X. Chen, S. K. Donaldson and S. Sun: *Kähler-Einstein metrics on Fano manifolds, I, II, III*. J. Amer. Math. Soc. **28** (2015), 183–278. 1, 2
- [CT08] X. Chen and Y. Tang: *Test configuration and geodesic rays*. Astérisque No. **321** (2008), 139–167.
- [CTW17] J. Chu, V. Tossatti and B. Weikove: *On the Regularity of Geodesics in the Space of Kähler Metrics*. Annals of PDE **3** (2017) no. 2. 6
- [CTW18] J. Chu, V. Tossatti and B. Weikove: *$C^{1,1}$ -regularity for degenerate complex Monge-Ampère equations and geodesic rays*. Communications in Partial Differential Equations **43** (2018), no. 2, 292–312. 21
- [CHT22] T. C. Collins, T. Hisamoto, R. Takahashi: *The inverse Monge-Ampère flow and applications to Kähler-Einstein metrics*. J. Differential Geom. **120** (2022), no. 1, 51–95. 2, 5
- [D15] T. Darvas: *The Mabuchi geometry of finite energy classes*. Adv. Math. **285** (2015), 182–219. 7
- [D17a] T. Darvas: *Weak geodesic rays in the space of Kähler potentials and the class $\mathcal{E}(X, \omega)$* . J. Inst. Math. Jussieu **16** (2017), no. 4, 837–858. 7
- [D17b] T. Darvas: *The Mabuchi completion of the space of Kähler potentials*. Amer. J. Math. **139** (2017), no. 5, 1275–1313.
- [DL20] T. Darvas and C. H. Lu: *Geodesic stability, the space of rays, and uniform convexity in Mabuchi geometry*. Geom. Topol. **24** (2020), no. 4, 1907–1967. 22
- [DR17] T. Darvas and Y. A. Rubinstein: *Tian’s properness conjectures and Finsler geometry of the space of Kähler metrics*. J. Amer. Math. Soc. **30** (2017), no. 2, 347–387. 15
- [DS16] V. Datar and G. Székelyhidi: *Kähler-Einstein metrics along the smooth continuity method*. Geom. Funct. Anal. **26** (2016), no. 4, 975–1010. 2, 25
- [D92] J. P. Demailly: *Regularization of closed positive currents and intersection theory*. J. Algebraic Geom. **1** (1992), no. 3, 361–409.
- [Der16a] R. Dervan: *Uniform stability of twisted constant scalar curvature Kähler metrics*. Int. Math. Res. Notices **15** (2016), 4728–4783. 3
- [Der16b] R. Dervan: *Alpha invariants and coercivity of the Mabuchi functional on Fano manifolds*. Ann. Fac. Sci. Toulouse Math. (6) **25** (2016), no. 4, 919–934.
- [DS20] R. Dervan, G. Székelyhidi: *The Kähler-Ricci flow and optimal degenerations*. J. Differential Geom. **116** (2020), no. 1, 187–203.
- [D88] W. Y. Ding: *Remarks on the existence problem of positive Kähler-Einstein metrics*. Math. Ann. **282** (1988), no. 3, 463–471. 2
- [DT92a] W. Y. Ding and G. Tian: *Kähler-Einstein metrics and the generalized Futaki invariant*. Invent. Math. **110** (1992), no. 2, 315–335.
- [DT92b] W. Y. Ding and G. Tian: *The generalized Moser-Trudinger inequality*. Nonlinear Analysis and Microlocal Analysis: Proceedings of the International Conference at Nankai Institute of Mathematics. World Scientific, 57–70, 1992.
- [D02] S. K. Donaldson: *Scalar curvature and stability of toric varieties*. J. Differential Geom. **62** (2002), no. 2, 289–349. 3, 4

- [D05] S. K. Donaldson: *Lower bounds on the Calabi functional*. J. Differential Geom. **70** (2005), no. 3, 453–472.
- [D15] S. K. Donaldson: *The Ding functional, Berndtsson convexity and moment maps*. arXiv:1503.05173v1. 2, 4, 5
- [EGZ11] P. Eyssidieux, V. Guedj, and A. Zeriahi: *Viscosity solutions to degenerate Complex Monge-Ampère equations*. Communications on Pure and Applied Mathematics **64**, 8 (2011) 1059–1094. 16
- [F16] K. Fujita: *A valuative criterion for uniform K-stability of \mathbb{Q} -Fano varieties*. arXiv:1602.00901v2. to appear in J. Reine Angew. Math. 20
- [F83] A. Futaki: *An obstruction to the existence of Einstein Kähler metrics*. Invent. Math. **73** (1983), no. 3, 437–443. 4, 9, 10
- [FM95] A. Futaki and T. Mabuchi: *Bilinear forms and extremal Kähler vector fields associated with Kähler classes*. Math. Ann. **301** (1995), no. 2, 199–210. 9, 25
- [GZ07] V. Guedj and A. Zeriahi: *The weighted Monge-Ampère energy of quasipsh functions*. J. Funct. An. **250** (2007), 442–482. 7
- [GZ17] V. Guedj and A. Zeriahi: *Degenerate complex Monge-Ampère equations*. EMS Tracts in Mathematics **26** (2017). 7, 16
- [Gui94] V. Guillemin: *Moment maps and combinatorial invariants of Hamiltonian T^n -spaces*. Birkhäuser Boston, 1994. ISBN: 0-8176-3770-2.
- [HL20] J. Han and C. Li: *On the Yau-Tian-Donaldson conjecture for generalized Kähler-Ricci soliton equations*. arXiv:2006.00903. To appear in Comm. Pure Appl. Math. 18
- [H16a] T. Hisamoto: *On the limit of spectral measures associated to a test configuration of a polarized Kähler manifold*. J. Reine Angew. Math. **713** (2016), 129–148. 3, 24
- [H16b] T. Hisamoto: *Orthogonal projection of a test configuration to vector fields*. arXiv:1610.07158. 3, 26
- [H18] T. Hisamoto: *Stability and coercivity for toric polarizations*. arXiv:1610.07998v2. 2, 3, 4, 13, 28
- [H19] T. Hisamoto: *Geometric flow, Multiplier ideal sheaves and Optimal destabilizer for a Fano manifold*. arXiv:1901.08480. 2, 10
- [I18] E. Inoue: *The moduli space of Fano manifolds with Kähler-Ricci solitons*. arXiv:1802.08128v3.
- [Li22] C. Li: *G-uniform stability and Kähler-Einstein metrics on Fano varieties*. Invent. Math. **227** (2022), no. 2, 661–744. 3, 31, 32
- [LX11] C. Li and C. Xu: *Special test configurations and K-stability of Fano varieties*. Ann. of Math. **180** (2014) no. 1, 197–232.
- [LZ17] Y. Li and B. Zhou: *Mabuchi metrics and properness of the modified Ding functional*. Pacific J. Math. **302** (2019), no. 2, 659–692. 15, 16
- [M86] T. Mabuchi: *K-energy maps integrating Futaki invariants*. Tohoku Math. J. (2) **38** (1986), no. 4, 575–593. 4
- [M87] T. Mabuchi: *Some symplectic geometry on compact Kähler manifolds I*. Osaka J. Math. **24** (1987), no. 2, 227–252. 6
- [M01] T. Mabuchi: *Kähler-Einstein metrics for manifolds with nonvanishing Futaki character*. Tohoku Math. J. (2) **53** (2001), no. 2, 171–182. 2
- [M03] T. Mabuchi: *Multiplier Hermitian structures on Kähler manifolds*. Nagoya Math. J. **170** (2003), 73–115. 3, 17
- [M57] Y. Matsushima: *Sur la structure du groupe d'homéomorphismes analytiques d'une certaine variété kählérienne*. Nagoya Math. J. **11** 1957 145–150. 17
- [N19] S. Nakamura: *Generalized Kähler-Einstein metrics and uniform stability for toric Fano manifold*. Tohoku Math. J. (2) **71** (2019), no. 4, 525–532. 2
- [N19] S. Nakamura: *Hessian of the Ricci Calabi functional*. Proc. Amer. Math. Soc. **147** (2019), 1247–1254. 3, 5, 17

- [NSY17] Y. Nitta, S. Saito, and N. Yotsutani *Relative GIT stabilities of toric Fano manifolds in low dimensions*. [arXiv:1712.01131v1](#). 3, 18
- [OT87] T. Ohsawa and K. Takegoshi: *On the extension of L^2 holomorphic functions*. *Math. Z.* **195** (1987), no. 2, 197–204. 31
- [PS07] D. H. Phong and J. Sturm: *Test configurations for K -stability and geodesic rays*. *J. Symplectic Geom.* **5** (2007), no. 2, 221–247. 23
- [PS10] D. H. Phong and J. Sturm: *Regularity of geodesic rays and Monge-Ampère equations*. *Proc. Amer. Math. Soc.* **138** (2010), no. 10, 3637–3650.
- [RT07] J. Ross and R. Thomas: *A study of the Hilbert-Mumford criterion for the stability of projective varieties*. *J. Algebraic Geom.* **16** (2007), no. 2, 201–255. 25
- [RWN14] J. Ross and D. Witt Nyström: *Analytic test configurations and geodesic rays*. *J. Symplectic Geom.* **12** (2014), no. 1, 125–169.
- [ST19] S. Saito and R. Takahashi: *Regularity of weak multiplier hermitian metrics and applications*. In preparation. 16
- [S92] S. Semmes: *Complex Monge-Ampère and symplectic manifolds*. *Amer. J. Math.* **114** (1992), no. 3, 495–550. 6
- [Szé08] G. Székelyhidi: *Optimal test-configurations for toric varieties*. *J. Differential Geom.* **80** (2008), no. 3, 501–523.
- [ST09] G. Székelyhidi and V. Tosatti: *Regularity of weak solutions of a complex Monge-Ampère equation*. *Anal. PDE* **4** (2011), no. 3, 369–378. 16
- [T97] G. Tian: *Kähler-Einstein metrics with positive scalar curvature*. *Inv. Math.* **130** (1997), 239–265. 4
- [Tia15] G. Tian. *K -stability and Kähler-Einstein metrics*. *Comm. Pure Appl. Math.* **68** (2015), 1085–1156. 1
- [TZ02] G. Tian and X. Zhu: *A new holomorphic invariant and uniqueness of Kähler-Ricci solitons*. *Comment. Math. Helv.* **77** (2002), no. 2, 297–325. 17
- [WN12] D. Witt Nyström: *Test configurations and Okounkov bodies*. *Compositio Math.* **148** (2012), 1736–1756. 23, 24
- [WZ04] X. Wang and X. Zhu: *Kähler-Ricci solitons on toric manifolds with positive first Chern class*. *Adv. Math.* **188** (2004), no. 1, 87–103. 3
- [X21] M. Xia: *On sharp lower bounds for Calabi type functionals and destabilizing properties of gradient flows*. *Anal. PDE* **14** (2021), no. 6, 1951–1976. 2, 10
- [Y17] Y. Yao: *Mabuchi Solitons and Relative Ding Stability of Toric Fano Varieties*. [arXiv:1701.04016](#). To appear on *Int. Math. Res. Not. IMRN*. 2, 29
- [Y22] Y. Yao: *Relative Ding stability and an obstruction to the existence of Mabuchi solitons*. *J. Geom. Anal.* **32** (2022), no. 4, 105. 28

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, FUROCHO, CHIKUSA, NAGOYA, JAPAN

E-mail address: hisamoto@math.nagoya-u.ac.jp